

Full Length Research Paper

QRS fragmentation as a prognostic test in acute coronary syndrome

Tarek M Abdelrahman

MD Cardiology Department, Minia University Hospital, Egypt.

Accepted 30 April, 2013

Among several invasive and non-invasive tests for risk stratification of acute coronary syndromes (ACS), fewer markers can be utilized in clinical practice. Our rationale is to validate the use of QRS-fragmentation as a promising bed-side test for assessment of prognosis in those patients. Collection of two-hundred and twenty patients with ACS was done in two years (from January 2011 till January 2013). Significant organic valve disease and QRS duration ≥ 120 ms as well as patients with permanent pacemakers were excluded. Patients were subjected to full clinical examination, ECG and Echocardiography in the first day of admission followed by diagnostic coronary angiography before discharge and a nuclear study was done for Randomized sample from each group. 12-leads ECG revealed fragmentation of QRS in 74 patients and 146 patients with no QRS fragmentation. Localization of the infarct site revealed NS difference between percentages in both groups. Echocardiography revealed a significant deterioration of LV functions in group-A than group-B. Also, MR jet area was significantly higher in group-A. Coronary angiography revealed severer lesions in group-A more than group-B. Nuclear study revealed higher percentages of irreversible scars in group-A (30%) and higher reversibility in group-B (80%). In-hospital Occurrence of complications from ACS revealed a significant higher incidence of MACE in group-A. Presence of fragmented QRS in surface ECG during ACS represents myocardial scar or fibrosis and reflects the severity of coronary lesions and a correlation between fQRS and depression of LV function is established. Indeed, occurrence of MACE is suspected.

Key words: Fragmented QRS (fQRS), acute coronary syndrome (ACS), major Adverse Cardiac Events (MACE).

INTRODUCTION

Observation of slurring and changes in the morphology of the QRS complex has been investigated since 1960's. Flowers et al. (1969) discovered the presence of fragmented QRS (fQRS) complexes among patients with prior myocardial infarction and patients with either right or left ventricular (LV) enlargement and reported it as high-frequency components.^[1] Friedman et al., described this fragmentation as persistent changes in Purkinje fibers and myocardial fibrosis which may cause slow and inhomogeneous myocardial activation after analyzing canine heart with induced myocardial infarction.^[2] Chryssos suggested that RSR' complex unrelated to right bundle branch block (RBBB) or left bundle branch block (LBBB) could be associated with impaired depolarization within tissue surrounding the myocardial scar.^[3] Bayes de Luna suggested that abnormalities of the second half of the QRS complex (terminal slurring, sometimes with R' in lead V1) during MI might represent necrosis in late depolarized basal zones.^[4]

Contemporary definition of the fQRS was defined by Das et al. in 2006, as presence of an additional R wave (R') or notching in the nadir of the S wave, or the presence of > 1 R' in 2 contiguous leads, corresponding to a major coronary artery territory on the resting 12-lead ECG with filter range 0.16–100 Hz, AC filter 60 Hz, paper speed 25 mm/s and 10 mm/mV.^[5] It has been shown in some studies that abnormality within the QRS complex can represent conduction disturbance and myocardial scar. Injured tissue around an infarct scar resulted in the RSR' pattern of the QRS complex. However, the diagnostic and prognostic values of these subtle abnormalities within the QRS complex were not clarified in prior studies.^[5,6] In 2006, Das et al. proved that, fragmented QRS complex in patients with coronary artery disease (CAD) was associated with myocardial conduction block due to myocardial scar detected by myocardial single photon emission tomography (SPECT). fQRS improved identification of prior myocardial infarction in patients who are being evaluated for CAD.^[5] As fQRS represents prior occurrence of myocardial infarction, further studies are directed to its role in identifying the risk of subsequent occurrence of ischemic events, its correlation to vent-

*Corresponding author. E-mail: tarekabelrahman2@yahoo.com.

ricular dysfunction and occurrence of congestive heart failure. Das et al. demonstrated that fQRS is an independent predictor of cardiac events in patients with CAD [6]. Other studies confirmed a relation between fQRS and LV dysfunction [7, 8] myocardial scar is also a substrate for reentrant ventricular tachyarrhythmia. Signal averaged electrocardiogram (SAECG) reveals the presence of late potential that indicates low-amplitude high-frequency potentials outside the terminal QRS complex. Abnormal late potential represents a slow conduction zone with damaged myocardium around the fibrosis of healed myocardial infarction.[9] The presence of late potential has been used for risk stratification of sudden cardiac death or lethal arrhythmic events.[10] As well as SAECG, fQRS also can reflect intra-cardiac conduction abnormality and will represent a substrate for ventricular arrhythmia. There has been only one study in which the correlation between fQRS and late potential detected by SAECG was investigated. This study demonstrated that the existence of fQRS appeared independently from the existence of late potential in patients with Brugada syndrome, but it is still unknown whether there is a correlation between fQRS and late potential in other diseases such as CAD and various cardiomyopathies. [11]

FQRS in various diseases

Coronary artery disease (CAD)

It was demonstrated that the fragmentation of QRS complexes represent myocardial scar in patients with suspected or known CAD in two major studies, one for narrow (QRS duration <120 ms)[5] and the other for wide (QRS duration ≥120 ms). [12] Sensitivity, specificity, and the negative predictive value for myocardial scar were 86%, 89%, and 93%, respectively, for the fQRS in the first study. And the second study was on 879 patients showed a sensitivity, specificity, positive predictive value and negative predictive value of f-wQRS for myocardial scar were 87%, 93%, 92% and 88%, respectively. [12] Gardner et al., proposed that fQRS complex in infarcted canine heart is caused by slow and inhomogeneous activation related to healed myocardial scar rather than changes in transmembrane resting or action potentials. [13] In the analysis of data from an epicardial and endocardial mapping from the patients undergoing LV aneurysm incision, Wiener et al. [14] demonstrated that, the fragmented electrical activity was present in all patients with LV aneurysms. Indeed he found that, patients with ventricular tachycardia had fragmented electrical activity than patients without ventricular tachycardia. The surgical excision of the area, which demonstrated fragmented activity, was associated with better survival and freedom from ventricular arrhythmia. Wiener et al., proposed that endocardial electrical activity mapping and detection of fragmented activity was a useful tool for surgically guided therapy for ventricular aneurysm and ventricular tachycardia. [15]

Non-coronary artery disease (Non-CAD)

Dilated Cardiomyopathy (DCM)

In patients with DCM, QRS fragmentation has been recorded during wavelet ECG analysis recorded from 6 unipolar left precordial leads, using a high-precision amplifier. When compared with the control subjects with no heart disease, these wavelets predicted frequent premature ventricular complexes and SCD. [16] In DCM, fQRS signifies myocardial scar defined by gadolinium delayed enhancement on cardiac magnetic resonance imaging. fQRS also signifies myocardial scar in non-CAD patients. [17]

Cardiac Sarcoidosis

Homsy et al. (2009), demonstrated that fQRS is a risk marker of cardiac sarcoidosis. The sensitivity and specificity of fQRS for detecting abnormal gadolinium-delayed enhancement images in cardiac magnetic resonance imaging were 100% & 80%, respectively. [18]

ARVD/C

fQRS has been identified as a risk marker of ARVD/C by Peters et al. (2008), they studied the value of QRS fragmentation in a standard 12-lead ECG in 360 patients with ARVD/C and compared its presence with the detection of the epsilon wave in highly amplified right precordial and modified limb leads in a subgroup of 207 patients. Other ECG signs of ARVD/C include QRS prolongation, prolonged S-wave upstroke, terminal activation delay, and epsilon potentials. Most of these signs are incorporated in typical fQRS. fQRS will probably simplify the ECG diagnosis of ARVD/C in patients with a high probability of the disease. [19].

Electrical diseases

Brugada Syndrome

This is characterized by cove-type ST elevation in right precordial leads and episodes of ventricular tachyarrhythmia. Patients with Brugada syndrome often had fQRS and it was more frequently observed in the VF group.[20] Recently, the PRELUDE study has shown that fQRS was useful for identify candidates for a prophylactic ICD implantation in patients with Brugada syndrome.[21]

Acquired long QT syndrome

(ALQTS) is a disease due to secondary repolarization abnormality. Repolarization abnormality, such as prolongation of QT interval and prolongation of the peak

to the end of T wave represented transmural or intraventricular dispersion of repolarization, and was associated with occurrence of torsades de pointes.^[22] It was found that; fQRS was present in a large percentage of ALQTS patients with syncope/torsades de pointes (81 %). Although onset of torsades de pointes is triggered activity due to early after depolarization, myocardial scar that appears as fQRS could be a substrate for subsequent reentrant arrhythmia.^[23]

Congenital and Rheumatic HD

In adult patients with repaired tetralogy of Fallot, fQRS predicted ventricular fibrosis detected by late Ga-MRI. Laterality of fQRS was associated with operative scar of the RV, and patients with fQRS had larger RV volume and lower RV EF. fQRS was closely associated with more extensive RV fibrosis and dysfunction.^[24] fQRS was frequent in patients with mitral stenosis caused by rheumatic fever. Rheumatic fever induces inflammation and degeneration of the cardiac valves and also injury of the myocardium. fQRS was associated with low EF, pulmonary hypertension, poor NYHA functional class, and decreased mitral valve area.^[7]

Miscellaneous Diseases

There has been an interest in fQRS as a screening tool in a systemic disease with cardiac involvement. Kadi et al., assessed the frequency of fQRS complexes in patients with rheumatoid arthritis (RA) and in control group of patients with fibromyalgia rheumatica. Patients with BBB and cardiovascular disease were excluded. Fragmented QRS were more often found among the patients with RA than a control group.^[25]

PATIENTS AND METHODS

In Cardiology department of Minia University, hospital two-hundred and twenty eligible patients with ACS during two years (from January 2011 till January 2013) were enrolled consecutively in this study. Patients with significant organic valve disease and those with any QRS morphology that had a QRS duration ≥ 120 ms as well as patients with permanent pacemakers were excluded from this study. All patients in the first day were subjected to full clinical examination, ECG using Fukuda Denshi Climax machine. A 12-lead admission ECG was obtained from all patients. All ECGs (filter range 0.5 Hz to 150 Hz, AC filter 60 Hz, 25 mm/s, 10 mm/mV) were analyzed by two independent clinicians who were blinded to the study design. The fQRS was defined by the presence of various RSR' patterns (QRS duration < 120 ms) with or without Q wave, which included an additional R wave (R' prime) or notching of the R wave or S wave, or the presence of more than one R prime (fragmentation) without typical bundle branch block in two contiguous leads corresponding to

a major lead set for major coronary artery territory. The presence of fQRS was detected by inspection of tracings with the naked eye. Analysis of the standard 12-lead ECG was performed without using any magnification. As shown in Figure.1, Fragmented QRS (RSR2 pattern and its variants). The different morphologies of fragmented QRS (fQRS) which includes various RSR2 patterns are shown in this figure. Note that if RSR2 patterns are present in right precordial leads (leads V1 and V2) with QRS > 100 msec (incomplete right bundle branch block), or QRS > 120 msec (complete right bundle branch block) and in left precordial lead (RSR2 in lead I, V5, and V6) with QRS > 120 msec (left bundle branch block), they are defined as complete or incomplete bundle branch block and excluded from the definition of fQRS, whereas if the RSR2 pattern is present in the mid-precordial lead or in inferior leads, they are defined as fQRS.^[6] Echocardiography using a commercially available GE vivid 3 echo machine equipped with 2.5 – 3.5 MHz transducer to examine for LV volumes including (end-diastolic EDV, end-systolic ESV) through which calculation of LVEF% is made by Simpson's Biplane method. Also, recording wall motions abnormalities and presence of aneurysm was done. Laboratory analysis of troponin-I and other routine investigations were recorded. Patients are then divided into two groups (group: A) 74 patients having fQRS in their resting ECG represents 33.6% and (group-B) were 146 patients having either pathological 'q' waves or repolarization changes only with no fQRS and represents 66.4%. All cases underwent diagnostic coronary angiography (CA) before discharge using a flat-panel GE-imaging system. All subjects were in the fast sedated state. CA was performed from the arterial femoral approach after local groin infiltration of 10-20 ml xylocaine 2% using modified seldinger's technique after injection of 5000 IU of Heparin, 6F JL then JR coronary catheters were used to engage the corresponding arteries. The study was conducted with a General Electric Innova 2000 angiographic unit (GE medical system Milwaukee, WI, USA). Due to lack of nuclear scan in our hospital, a randomized selection of 20 patients in each group was sent to nuclear study in specialized centers. From group (A): 10 patients having anterior MI (anteroseptal and anterolateral MI were included) and 10 patients having inferior wall MI (inferoposterior and inferolateral MI were included). The same sample was taken from group (B) also as to compare their scan results. As shown in Figure (2) Clinical characteristics, which consisted of multiple descriptors from each patient's history and physical examination, were collected by physicians from cardiology clinic for each patient before cardiac catheterization and were stored in the database of coronary angiography laboratory at our institution. We recorded the baseline characteristics, which includes hypertension, diabetes mellitus, smoking history, family history for coronary artery disease (CAD) and lipid parameters. Patients with significant organic valvular heart disease and BBB (Left BBB, incomplete or complete Right BBB or duration QRS > 120 ms), and patients with permanent pacemakers were excluded

Figure 1.





Figure 2. Nuclear imaging (99m Tc-TF) showed a fixed inferior wall myocardial perfusion defect in patient with recent inferior Myocardial infarction.

Table 1. Baseline characteristics of the study population.

Parameter	Fragmentation of QRS in admission ECG		P- Value
	Fragmented QRS	Non-fragmented QRS	
	Group (A): N= 74 (33.6%)	Group (B): N= 146 (66.4%)	
Age (yrs)	55 ± 9	56 ± 8	NS
Height (cm)	167 ± 4	166 ± 8	NS
Weight (kg)	85 ± 2	84 ± 6	NS
BMI (kg/m ²)	33.2 ± 4	32.8 ± 8	NS
Gender male, n (%)	61 (82%)	122 (83%)	NS
Hypertension, n (%)	38 (51%)	77 (52%)	NS
Diabetes mellitus, n (%)	40 (53%)	76 (52%)	NS
Hyperlipidemia, (n %)	52 (70%)	102 (69%)	NS
Serum creatinine (mg%)	1.5±0.2	1.3±0.3	NS
Medications			
ASA, n (%)	62 (83%)	119 (81%)	NS
Clopidogrel , n (%)	18 (24%)	32 (22%)	NS
ACEI, n (%)	33 (44%)	62 (42%)	NS
ARB, n (%)	22 (29%)	46 (31%)	NS
B-Blockers, n (%)	66 (89%)	128 (87%)	NS
CCB, n (%)	12 (16%)	21 (14%)	NS
Statin, n (%)	28 (37%)	56 (38%)	NS
Diuretics, n (%)	8 (10.8%)	18 (12%)	NS
OAD/Insulin, n (%)	17 (12%)	19 (13%)	NS

from the study. MACE was defined as cardiac death, re-infarction, heart failure, cerebrovascular event, sustained ventricular tachycardia or fibrillation. Low cardiac output syndrome was defined as usage of inotropic or intraaortic balloon pump support for more than 30 minutes in the coronary care unit in order to maintain adequate hemodynamics after appropriate adjustment of preload and after load.

Statistical Methods

Data were collected and analyzed by Statistical Packa-


ge for Social Sciences (SPSS) version 13 Data are given as mean (SD) for both numbers and percentages and compared using Chi-square and the Student t test . Pearson's correlation was made for the relations between MACE and myocardial scars in nuclear scan to fQRS positive patients. A probability level of p-value ≤ 0.05 was considered as statistically significant in all tests

RESULTS

Patients characteristics as regard Age, Gender, BMI,

Table 2. ECG Localization of Infarct site.

Site of MI	Group-A (N=74)	Group-B (N=146)	P value
Anterior MI	36 (48.6 %)	68 (46.5%)	NS
Inferior MI	12 (16.2 %)	34 (23.3%)	NS
Lateral MI			
- anterolateral	9 (12.3 %)	18 (12.4%)	NS
- inferolateral	6 (8.1 %)	12 (8.3%)	
Posterior MI			
- inferoposterior	6 (8.1 %)	8 (5.3 %)	NS
- posterolateral	5 (6.7 %)	6 (4.2 %)	

Figure 3. Localization of infarct-site in Group-A

Concomitant medical diseases (hypertension, diabetes mellitus and Hyperlipidemia), Renal functions and medications used inside CCU are shown in table (1) and revealed no significant statistical differences between both fragmented and non-fragmented groups. 12-lead ECG revealed presence of fragmentation of QRS in 74 patients named as (Group-A) and 146 patients with no QRS fragmentation and named as (Group-B). Localization of the infarct site in each group revealed presence of anterior MI in 36 patients in group-A represents (48.6%) of the total group and 68 patient in group-B (46.5%), inferior MI were 12 patient in group-A (16.2%) and 34 patients in group-B (23.3%), anterolateral MI were 9 patients in group-A (12.3% and 18 patients (12.4%) in group-B, inferolateral MI were 6 patients in group-A (8.1%) and 12 patients (8.3%) in group-B, inferoposterior MI were 6 (8.1%) in group-A versus 8 patients (5.3%) in group-B and finally, posterolateral MI were 5 (6.7%) in group-A versus 6 (4.2%) in group-B. As shown in table (2),

figure (3) and (4). Chi square analysis for descriptive values revealed non-significant differences between both groups at any site. Echocardiographic findings in both groups were tabulated and revealed a significant increase in Lv volumes and significant decrease in EF% and FS % in group-A than group-B. Also, MR jet area was significantly high in group-A which concludes deterioration in LV functions in group-A. As shown in Table (3).

Coronary Angiographic Findings

Semi-quantitative measurement of lesions severity revealed more significant lesions in group-A than in group-B as shown in Table (4).

Nuclear Findings

Twenty patients were randomly selected from each




Table 3. Echocardiographic findings.

Echocardiographic Finding	Group-A (N=74)	Group-B (N=146)	P value
Lv end diastolic volume (LVEDV) ml	131±18	98±12	0.001
Lv end systolic volume (LVESV) ml	88±15	65±5	0.001
Ejection Fraction (EF %)	49±4	61±4.4	0.001
Fractional shortening (FS %)	20±2.1	27±2.8	0.001
MR jet area width (Cm ²)	2.5±0.4	0.7±0.2	0.001

Table 4.

Coronary artery	Lesion severity		P value
	Group-A (N=74)	Group-B (N=146)	
Left Main artery (LM)	40% ±10	20%	0.001
Left anterior descending A (LAD)	80%±20	40%±10	0.001
Diagonal branch artery (DA)	90%±10	50%±10	0.001
Left circumflex artery (LCx)	60%±20	40%±10	0.001
Obtuse marginal artery (OM)	80%±10	50%±10	0.001
Right coronary artery (RCA)	70%±30	40%±10	0.001

group and detection of irreversible scar tissue and reversible ischemia in every group was calculated. Results revealed a significantly higher percentages of

irreversible scars in group-A (30%) and higher percentages of reversibility in group-B (80%). As shown in Figure (5).





Figure 5. showed percentages of scar tissue and reversible ischemia in both groups.

Figure 6. MACE in Group-A versus Group-B.



MACE

In-hospital Occurrence of complications from ACS were studied and revealed a significant ($p < 0.05$) higher incidence of MACE in group-A in the form of 6 cardiac deaths in group-A versus no cardiac deaths in group-B, Re-infarction in 8 patients in group-A versus 2 patient in group-B, symptomatic heart failure in 26 patient in group-A 5 cases from them needed a circulatory support in relation to only 8 cases developed heart failure in group-B, none of them needed a circulatory support. Cerebrovascular stroke was found in 4 cases in group-A versus no cases in group-B, sustained VT or VF needed cardioversion occurred in 18 patient in group-A versus 6 cases in group-B most of them was self-terminated. As shown in the figure (6):

DISCUSSION

Presence of fragmented QRS represents distortion of signal conduction and depolarization process within the

ventricles which is related to myocardial scar/myocardial ischemia or myocardial fibrosis. The short-term and in-hospital prognosis is still dispute. [26] About the methodology used different questions related to the technique of ECG used are raised. Das et al., in 2008 proved that, ECG recording used to detect fQRS is not a specific setting and is the same as routine 12-lead ECG recording: high-pass filter: 0.05-20 Hz (usually 0.15 Hz), low-pass filter: 100-150 Hz, AC filter: 50 or 60 Hz, paper speed: 25-50 mm/sec (usually 25mm/sec) and voltage 1mm/mV. [16] Our study detect a high percentage (30%) of irreversible scar tissues in group-A (fQRS) and this is supported by many studies where Myocardial single photon emission tomography (SPECT) was used to identify regional perfusion abnormalities from a scar by a prior myocardial infarction and significantly greater perfusion and functional abnormalities than was the Q wave was proved. [5, 16] Also, the combination of Q wave and fQRS improved the sensitivity (91.4%), specificity (89%), and negative predictable value (92.4%) for detection of scared myocardium. Wang et al., 2010 failed to show

significance for fQRS in detection of myocardial scar in patients with CAD. The reason for the different result from previous studies is unclear, and the authors stated that larger study is required to confirm the diagnostic value of fQRS.^[28] Our study correlates fQRS to the coronary angiographic findings and detects a higher percent and severer lesions in fQRS-group than Non fQRS patients. The findings of Yutaka et al. 2012, support this idea, they found that fQRS in acute coronary syndrome appeared within 48 hours (especially within 24 hours) from the onset of symptoms and persisted thereafter. fQRS on 12-lead ECG developed in 55% of patients with ST elevation myocardial infarction (STEMI) and in 50% of patients with non-ST elevation myocardial infarction (NSTEMI), but in only 3.7% of patients with unstable angina pectoris. Although the sensitivities of fQRS for STEMI and NSTEMI were 55% and 50% respectively, the specificity of fQRS for AMI was 96%.^[29] Our study proved that, patients with fQRS showed a lower LV functions in the form of significant increased LV volumes, lower EF% and higher MR degree than non-fQRS group. Yan et al., in 2012 demonstrated a strong relation between fragmentation of QRS and depression of LV functions even in a sub-clinical situations.^[30] Also, many studies support a high sensitivity of fQRS to the presence of symptomatic LV failure in CAD patients.^[8,31] Since fQRS represents myocardial scar, fQRS may be associated with heart failure and ventricular tachyarrhythmia. Major adverse cardiac events (MACE) were studied in our research and revealed a significant ($p < 0.05$) higher incidence of cardiac deaths in group-A than group-B. Some studies have shown a strong relationship between existence of fQRS in patients with CAD and prognosis. In a study made by Das et al., 2007 patients with narrow QRS complexes that showed a fragmentation of QRS have higher all-cause mortality (34.1% vs. 25.9%) and cardiac event rate (49.5% vs. 27.6%) than patients without fQRS.^[6] A study made by Korhonen et al., 2010 in which fQRS in patients with prior myocardial infarction was evaluated by a body surface mapping system also showed that fQRS was an independent predictor for cardiac death and hospitalization due to heart failure.^[31] Torigoe et al. 2012 evaluated fQRS by 12-lead ECG and showed that the number of the leads with fQRS was a predictor for cardiac death and hospitalization for heart failure in patients with prior myocardial infarction. Although prior criteria for positive fQRS included the presence of fQRS in 2 or more contiguous leads, they showed that the presence of fQRS in 3 or more leads was the most useful for distinguishing between patients with and without risk for cardiac death or hospitalization.^[32] An increase in the number of leads with fQRS would represent a wide scar area, which would result in an adverse outcome. fQRS also predicted intra- and post-operative hemodynamic instabilities and adverse cardiovascular events in patients undergoing coronary artery bypass graft surgery. Mortality of patients with f-wQRS was significantly higher than that of patients without f-wQRS, and subgroups of f-wQRS (fragmented left

BBB, fragmented PVC and fragmented paced QRS) were also associated with short time to death compared to patients without f-wQRS. Multivariate analysis showed that reduced EF ($\leq 35\%$) and f-wQRS were independently associated with mortality.^[33] For more confirmation of the effect of fragmentation of QRS in ECG, a number of follow-up studies^[32,34] discovered a strong relation between fragmentation of QRS and cardiac death. The findings of Pietrasik et al., in 2010, who studied the appearance of fQRS in ECG of patients with first Q-wave MI, resolved Q wave, and persistent Q wave on 2 month follow-up and its relation to the risk of recurrent adverse cardiac events defined as, unstable angina, recurrent MI or cardiac death. They found a significant correlation between fQRS with the higher risk of recurrent adverse cardiac events among patients with resolved Q wave. It was proposed that fQRS may identify ischemic myocardium.^[34] Another researches found a strong correlations between fQRS and cardiac mortality in long-term periods especially in those need ICD or CRT therapy. Das et al., studied 361 patients in a follow up period from 6 months to more than 2 years and detected a significant increased risk of arrhythmic events but no effect on mortality.^[17] Brenyo et al.,^[35] and Pietrasik et al.,^[34] during (19 ± 12 months) discover no effect in total population, however, increased risk of death only among patients with LBBB and inferior fQRS. Cheema et al., evaluated the association between fQRS complex and primary end-point of all cause mortality and secondary end-points specific cause mortality and appropriate ICD shocks in the cohort of 842 patients with LV dysfunction (LVEF $\leq 35\%$), both ischemic and non-ischemic etiology.^[8] Fragmented QRS was present in 32.5% of patients. Mean follow-up of the study was 40 ± 17 months for entire cohort. There was no significant difference in event-free survival between patients with fQRS and without fQRS. Fragmented QRS was not associated with increased risk of all-cause mortality, arrhythmic mortality and non-arrhythmic mortality. Similar conclusions were drawn from study by Forleo et al.,^[36]

Limitations of the study

FQRS on a 12-lead ECG requires an optimal low pass filter setting (100 or 150 Hz). Fragmentation may be missed with a filter setting of 40 or 60 Hz. Apart from notching, slurring of fQRS may also represent myocardial scar.

The present definition of fQRS is being qualitative, prevents the incorporation of slurring in this phenomenon. A quantitative definition of fQRS with development of specific software is underway.

It is to be emphasized that fQRS is a nonspecific finding and should only be interpreted in the presence of pertinent clinical evidence of myocardial scar as in CAD or primary electrical abnormalities of depolarization such as Brugada syndrome.

The effects of different therapeutic strategies (thrombo-

lysis or revascularization) on fQRS and prognosis of CAD patients need a further study.

Correlation between QRS fragmentation and the novel selvester QRS-score has to be established.

A further follow up should be evaluated well as many researches did.

A larger population sampling supports such tools to increase its validity.

ACKNOWLEDGMENT

To the (Team worker) colleagues in Non-invasive unit of Cardiology Department of El Minia University Hospital and the secretary, employees of the Department for easy call back and the archiving process.

Funding sources

All the study fees were sponsored by the governmental policies to support the research work in my university according to the established financial protocols

Disclosures: None

Ethics Approval: All patients wrote a written approval to be included in the research work

List of Abbreviations

EF: Ejection fraction

NYHA: New York Health Association Classification

MACE: Major Adverse Cardiac Events

ACS: Acute Coronary Syndrome

fQRS: Fragmented QRS-complex in ECG

ECG: Electrocardiography

GE: General Electric

ICD: Intracardiac defibrillation device

CRT: Cardiac Re-synchronization Therapy

REFERENCES

Flowers NC, Horan LG, Thomas JR, Tolleson WJ (1969). The anatomic basis for high-frequency components in the electrocardiogram. *Circulation*.39.pp. 531–539.

Friedman PL, Fenoglio JJ, Wit AL (1975). Time course for reversal of electrophysiological and ultrastructural abnormalities in subendocardial Purkinje fibers surviving extensive myocardial infarction in dogs. *Circ. Res.* (36). pp. 127–144.

Varriale P, Chryssos BE (1992). The RSR' complex not related to right bundle branch block: Diagnostic value as a sign of myocardial infarction scar. *Am. Heart. J.*(123). Pp. 369–376.

Bayes de Luna A (1998). Electrographic patterns of ischemia, injury and necrosis. In: Bayes de Luna A ed. *Clinical electrocardiography: A textbook*. 2nd Ed.

Futura Publishing Company, Inc. Armonk, NY. Pp. 141–142.

Das MK, Khan B, Jacob S, Kumar A, Mahenthiran J (2006). Significance of a fragmented QRS complex versus a Q wave in patients with coronary artery disease. *Circulation*. (113). Pp. 2495–2501.

Das MK, Saha C, El Masry H, (2007). Fragmented QRS on a 12-lead ECG: a predictor of mortality and cardiac events in patients with coronary artery disease. *Heart Rhythm*. (4).pp.1385-92.

Yuce M, Davutoglu V, Ozbala B (2010). Fragmented QRS is predictive of myocardial dysfunction, pulmonary hypertension and severity in mitral stenosis. *Tohoku. J. Exp. Med.* (220). pp. 279-83

Cheema A, Khalid A, Wimmer A (2010). Fragmented QRS and mortality risk in patients with left ventricular dysfunction. *Circ. Arrhythm. Electrophysiol.* (3). Pp.339-44.

Breithardt G, Cain ME, el-Sherif N (1991). Standards for analysis of ventricular late potentials using high-resolution or signal-averaged electrocardiography. A statement by a Task Force Committee of the European Society of Cardiology, the American Heart Association, and the American College of Cardiology. *Circulation*.(83).pp.1481-8.

Gomes JA, Cain ME, Buxton AE, Josephson ME, Lee KL, Hafley GE (2001). Prediction of long term outcomes by signal-averaged electrocardiography in patients with unsustained ventricular tachycardia, coronary artery disease, and left ventricular dysfunction. *Circulation*. (104). pp.436-41.

Morita H, Kusano KF, Miura D (2008). Fragmented QRS as a marker of conduction abnormality and a predictor of prognosis of Brugada syndrome. *Circulation*. (118). pp.1 697-704.

Das MK, Suradi H, Maskoun W, Michael MA, Shen C, Peng J, et al (2008). Fragmented Wide QRS on a 12-Lead ECG: A Sign of Myocardial Scar and Poor Prognosis. *Circ. Arrhythmia. Electrophysiol.* (1).pp.258-68.

Gardner PI, Ursell PC, Fenoglio JJ Jr, Wit AL (1985). Electrophysiologic and anatomic basis for fractionated electrograms recorded from healed myocardial infarcts. *Circulation*. (72). Pp.596– 611.

Wiener I, Mindich B, Pitchon R (1982). Determinants of ventricular tachycardia in patients with ventricular aneurysms: Results of intraoperative epicardial and endocardial mapping. *Circulation*. (65). pp.856–861.

Wiener I, Mindich B, Pitchon R (1984). Fragmented endocardial electrical activity in patients with ventricular tachycardia: A new guide to surgical therapy. *Am. Heart. J.*(107). Pp. 86–90.

Das MK, Suradi H, Maskoun W (2008). Fragmented wide QRS on a 12-lead ECG: A sign of myocardial scar and poor prognosis. *Circ Arrhythm Electrophysiol.* (1).pp.258–268.

Das MK, Maskoun W, Shen C, Michael MA, Suradi H, Desai M, et al (2010). Fragmented QRS on twelve-lead electrocardiogram predicts arrhythmic events in patients with ischemic and nonischemic cardiomyopathy. *Heart Rhythm*. (7). pp.74-80.

- Homsy M, Alsayed L, Safadi B, Mahenthiran J, Das MK (2009). Fragmented QRS complexes on 12 lead ECG: a marker of cardiac sarcoidosis as detected by gadolinium cardiac magnetic resonance imaging. *Ann Noninvasive Electrocardiol.*(14).pp.319-26.
- Peters S, Trümmel M, Koehler B (2008). QRS fragmentation in standard ECG as a diagnostic marker of arrhythmogenic right ventricular dysplasia-cardiomyopathy. *Heart Rhythm* (5).pp.1417- 21.
- Takagi M, Yokoyama Y, Aonuma K, Aihara N, Hiraoka M (2007). Clinical characteristics and risk stratification in symptomatic and asymptomatic patients with brugada syndrome: multicenter study in Japan. *J Cardiovasc Electrophysiol.* (18).pp.1244-51.
- Priori SG, Gasparini M, Napolitano C, et al (2012). Risk stratification in Brugada syndrome: results of the PRELUDE (PRogrammed ELectrical stimUlation preDICTive valuE) registry. *J. Am. Coll. Cardiol.* (59).pp.37-45.
- Antzelevitch C (2005). Role of transmural dispersion of repolarization in the genesis of drug-induced torsades de pointes. *Heart Rhythm.* (2).pp.9-15.
- Haraoka K, Morita H, Saito Y, et al (2010). Fragmented QRS is associated with torsades de pointes in patients with acquired long QT syndrome. *Heart Rhythm.* (7).pp.1808-14.
- Park SJ, On YK, Kim JS, et al (2012). Relation of Fragmented QRS Complex to Right Ventricular Fibrosis Detected by Late Gadolinium Enhancement Cardiac Magnetic Resonance in Adults With Repaired Tetralogy of Fallot. *Am. J. Cardiol.* (109). pp.110-5.
- Kadi H, Inanir A, Habiboglu A et al (2011). Frequency of fragmented QRS on ECG is increased in patients with rheumatoid arthritis without cardiovascular disease: A pilot study. *Mod. Rheumatol.* DOI: 10.1007/s10165-011-0493. P.9.
- Grzegorz P, Wojciech Z (2012). QRS fragmentation: Diagnostic and prognostic significance. *Cardiol. J.* Vol. 19, No. 2, pp.114–121.
- Wang DD, Buerkel DM, Corbett JR, Gurm HS (2010). Fragmented QRS complex has poor sensitivity in detecting myocardial scar. *Ann Noninvasive Electrocardiol.* (15).pp.308-14.
- Yutaka T, Hiroshi M (2012). *Indian. Pacing. Electrophysiol. J.* (ISSN 0972-6292). 12. (5). pp. 213-225.
- Yan GH, Wang M, Yiu KH (2012). Subclinical Left Ventricular Dysfunction Revealed by Circumferential 2D Strain Imaging in Patients with Coronary Artery Disease and Fragmented QRS Complex. *Heart Rhythm.* (9). pp.928-935.
- Korhonen P, Husa T, Konttila T (2010). Fragmented QRS in prediction of cardiac deaths and heart failure hospitalizations after myocardial infarction. *Ann. Noninvasive. Electrocardiol.* (15).130-7.
- Torigoe K, Tamura A, Kawano Y, Shinozaki K, Kotoku M, Kadota J (2012). The number of leads with fragmented QRS is independently associated with cardiac death or hospitalization for heart failure in patients with prior myocardial infarction. *J. Cardiol.* (59).pp.36-41.
- Erdogan T, Cetin M, Kocaman SA (2012). Relationship of fragmented QRS with prognostic markers and in-hospital MACE in patients undergoing CABG. *Scand Cardiovasc. J.*(46). pp. 107-13.
- Pietrasik GM, Polonsky S, Moss AJ, Zareba W (2010). Presence of fragmented wide-QRS complex and the risk of death and sudden cardiac death among MADIT-II patients with left bundle branch block. ACCPoster Atlanta, USA.
- Brenyo AJ, Pietrasik G, McNitt S, Zareba W (2010). QRS fragmentation: Lack of association with cardiac events in patients with a normal QRS duration. MADIT II Heart Rhythm Denver, CO, USA.
- Forleo GB, Della Rocca DG, Papavasileiou LP (2011). Predictive value of fragmented QRS in primary prevention implantable cardioverter defibrillator recipients with left ventricular dysfunction. *J. Cardiovasc. Med. (Hagerstown)*,(12). pp. 779–784.