

*Full Length Research Paper*

# Evaluation of seedling emergence of *Senna siamea* as influenced by crude oil in soil

Egwu .E.A, Eric Nwachukwu and Emmanuel Ugoh

Department of Forestry and Wildlife, Faculty of Agriculture, Delta State University, Asaba Campus, Asaba, Delta State, Nigeria.

Accepted 06 July, 2014

A study was carried out in 2011 at the Nursery site of the Department of Forestry and Wildlife, Faculty of Agriculture, Delta State University, Asaba Campus, Nigeria to evaluate the seedling emergence of *Senna siamea* as influenced by crude oil in soil. 0.0, 2.0, 4.0, 6.0, 8.0 and 10.0% (w/w) of the oil constituted treatments 10 seeds were sown in the soil crude oil mixture. The experiment was laid out in a randomized complete block design with four replicates. The results showed that contamination of soil with crude oil significantly ( $P \geq 0.05$ ) reduced percentage germination, delayed germination by increasing the days to 50% germination and reducing the rate of germination in *S siamea* when compared with the uncontaminated plots. The effects were generally oil level-dependent. Only two seeds planted in 10.0% (w/w) of soil contaminated germinated while the majority did not sprout. This study has shown that oil in soil has a significant effect of reducing seedling emergence in *Senna siamea*. This study has a great ecological significance especially in the Niger Delta.

**Key words:** Seedling emergence, *Senna siamea*, crude oil, contaminated soil.

## INTRODUCTION

*Senna siamea* belongs to the plant family Fabaceae and order Fabales. It is commonly called black wood cassia, Bombay black cassia, iron wood, kassod tree. It is a medium sized, evergreen tree growing up to 18m fall with a straight trunk of up to 30 cm in diameter. The bowl is short, crown usually dense and rounded at first later becoming irregular and spreading with dropping branches (Okafor, 1980). The root system consists of a few thick roots, growing to a considerable depth and a dense mat of rootlets in the top 10 - 20 cm of soil, which may reach a distance of 7 m from the stem in one year and eventually a distance up to 15 m (Keay et al., 1989). The leaflets 6 - 12 pairs are on short stalks of 3 mm. The pods are numerous,

long; narrow, between 5 - 25 cm long, 12 - 20 mm broad with many bright yellow flowers (Dalziel, 1987). The seeds are bean - shaped, shining, dark brown, 8 mm long with distinct areole (small clearly defined space) (Opeke, 1992). *S. siamea* is a native to South and Southern Asia but it has been cultivated for so long in other places. Okafor (1980) reported that plantations of *S. siamea* were established in the 1920s in Ghana, Nigeria and Sierra Leone mainly for fuel wood. Ecologically, *S. siamea* will grow in a range of climatic condition but is particularly suited to lowland tropics with a monsoon climate. The plant grows only when its roots have access to groundwater and the maximum length of dry period should not exceed 4 - 8 months. It is susceptible to cold and frost and does not do well at altitudes above 1:300 m. The plant starts flowering and fruiting at the age of 2 - 3 years (Dalziel, 1987). Once established, it flowers precociously and abundantly throughout the year.

\*Corresponding Author. E-mail: [egwu\\_ekanem@hotmail.com](mailto:egwu_ekanem@hotmail.com)

The seeds remain viable for 2 - 3 years under Sahel conditions. Seeds germinate readily without treatment but older seeds may receive some forms of pre-treatments.

Economically, *S. siamea* is a rather fast growing species for the production of pole timber and fuel wood in sudation ecozones. It regenerates vigorously by coppicing, rotation of 4 - 7 years for pole production. Throughout the Sahel, it is used as fuel wood. The wood is fairly heavy with a specific gravity of 0.6 - 0.8 which makes it appropriate size suitable for furniture, turnery and cabinet work. The use of poles-posts in various endeavours cannot be over emphasized. Leaves and pods are widely browsed by ruminants but highly toxic to pigs, poultry and possibly to other monogastrics (Gutteridge, 1997). The young leaves and flowers of *S. siamea* are used in curry dishes. The species is also used for the production of honey and tannins. It is suitable for windbreaks, shelterbelt plantation but not as shade tree in agro forestry because of root competition. It can be used as shade tree in cocoa, coffee and tea plantations roadside and court yard planting in the south planting erosion control, villages and squares laxative properties and used for a variety of ailments, of blood forming organs, genitourinary tract, also for herpes and rhinitis (F/FRED, 1994).

Petroleum and its refined products account for over 90% of Nigeria source of income and during its exploration and exploitation, crude oil is inevitably discharged into the environment through transportation, oil well blowouts, rupture/leakages of oil storage tanks and pipelines terminal loading, operational discharge and accidental discharge (Terje, 1984; Ogri, 2001; Agbogidi and Dolor, 2007). Oil pollution effects vary from growth reduction, yield reduction, growth stagnation wilting of plant and total death. Oil pollution effects depends on the type of oil, plant types, oil level, time of application, adequacy of plant responses and other environmental factors (Agbogidi et al., 2007). Crude oil contamination is a serious environmental problem in the Niger Delta area of Nigeria because it affects agricultural activities of the rural inhabitants as well as the socio-economic (Agbogidi et al., 2001). Studies have been conducted on the effects of oil pollutions using various plant species including African star apple (Agbogidi and Ejemete, 2005), castor oil (Vwioko and Fashemi, 2005), native pear (Agbogidi and Eshgebeyi, 2006), dikanut (Agbogidi and Dolor, 2007), information on soil contamination and *S. siamea* is however, scarce. The aim of this study was to evaluate the seedling emergence in *S. siamea* as affected by crude oil in soil with a view to recommending the critical level of crude oil to this multi-purpose tree species.

## MATERIALS AND METHODS

The experiment was conducted in 2011 for a period of

two weeks at the Forestry and Wildlife Research Farm of the Department of Forestry and Wildlife, Faculty of Agriculture, Delta State University, Asaba Campus, Delta State, Nigeria (Asaba Meteorological Station, 2010). The seeds of *S. siamea* were collected from a *S. siamea* tree behind Hall B in Delsu, Asaba Campus. Crude oil was with specific gravity  $0.8334 \text{ cm}^{-3}$  was sourced from NNPC, Warri, Delta State. The soil samples collected was dir-dried and passed through a 2 mm sieve. The oil was applied at 0.0, 2.0, 4.0, 8.0 and 10.0% w/w (based on weights of oil and soil). The soil/oil samples were thoroughly mixed together by hand. The mixtures were poured into bottom perforated polypots (15 × 10 cm in dimension). The seeds (10) were later planted in the polypots, watered to field capacity immediately after planting and later, once in two days. The composite soil samples were collected from 0 - 20 cm depth prior to treatment application and at the end of experiment and used to determine soil physico-chemical properties such as soil pH, texture, nutrients water holding capacity, moisture, bulk density and structure. Similarly, the characteristics of the crude oil used were determined in NNPC laboratory, Warri prior to experimentation. The set up was arranged in randomized complete block design with four replications. The polypots were kept in the nursery for subsequent examination. Parameters measured were germination percentage, days to germination and rate of germination. Data collected were subjected to analysis of variance while significant means were separated with the Duncan's multiple range tests using SAS (2005).

## RESULTS AND DISCUSSION

Table 1 shows the soil physico-chemical properties before experimentation. The soil is sandy loam in texture with characteristics of 94.5% (sand), 2.1% (silt) and 3.4% (clay). The pH 5.60 indicated that the experimental site is slightly acidic. Both the organic matter (2.64g/kg) total nitrogen (0.06%) is relatively low. The characteristics of the crude oil used are presented in Table 2. The oil contains the major nutrients including carbon, nitrogen, sulphur, hydrogen, oxygen and trace metals in varying quantities.

The germination characteristics of *S. siamea* as influenced by crude oil in soil are presented in Table 3. Over 98% of the seeds of *S. siamea* planted in the uncontaminated soils germinated as from the 3<sup>rd</sup> day after sowing. The percentage germination of seeds sown in soils with 2.0%w/w crude oil contamination was 94.87. No significant ( $P \leq 0.05$ ) differences existed in the seeds sown in the control plots and those exposed to 2.0%w/w oil treatment. Significant reductions ( $P \geq 0.05$ ) were however, observed as the level of oil in soils increased. As low as 22.67% germination was recorded for *S. siamea* seeds planted in soils that received 6.00%w/w

**Table 1.** Physico-chemical properties of soil before experimentation.

Parameters	Asaba
Sand (%)	94.5
Silt (%)	2.1
Clay (%)	3.4
Soil pH	5.60
Textural class	Sandy loam
Organic carbon (%)	0.91
Organic matter (gkg <sup>-1</sup> )	2.64
Total N (%)	0.06
Available P (mg/kg)	30.00
Ca <sup>2+</sup> (cmol/kg)	1.31
Mg <sup>2+</sup> (cmol/kg)	0.16
Na <sup>+</sup> (cmol/kg)	0.25
K <sup>+</sup> (cmol/kg)	0.17
H <sup>+</sup> (cmol/kg)	0.45
Al <sup>3+</sup> (cmol/kg)	0.08
ECEC (cmol/kg)	2.42
Base saturation (%)	78.10

**Table 2.** Characteristics of the crude oil used.

Parameters	Asaba
Specific gravity (cm <sup>-3</sup> )	0.8334
API gravity	34.2897
Bottom temperature (°C)	98.0
Viscosity (cp)	0.27
Gas oil ratio	881
Reservoir depth (m)	2556
Carbon (%)	84.8
Hydrogen (%)	12.61
Sulphur (%)	1.48
Nitrogen (%)	0.48
Oxygen	0.50
Trace metals (%)	0.13

crude oil treatment (Table 3). In the same vein, it took 3.4 days for *S. siamea* seeds sown in the control plots to sprout when compared to seeds exposed to 8.0%w/w of crude oil that sprouted at about 7 days after sowing indicating growth reduction and stagnation. The observed reduction in growth of crops in crude oil contaminated soils may be attributed to one or a combination of the following factors: slight acidity of the soil, poor wetability and aeration of the soil, loss of seed viability, penetration of the oil components into the embryo and effect of the oil on the metabolic activities of the seeds. The radicle could have emerged and the oil affected subsequent events relating to seedling growth. If a seed is dehydrated at the phase of seedling growth, such a seed is killed (Agbogidi

et al., 2006). Since increasing respiratory activities are associated with germination, there could be a correlation between any treatments that affects respiration in the soil. The physical surface characteristics of the oil which make it function as a physical barrier to water and oxygen uptake thus adversely affecting gaseous exchange as well as a delay in days to seedling emergence with increasing oil level. The rate at which the seeds of the test plant germinated were also, significantly ( $P \geq 0.05$ ) affected. Almost all the seeds (9.8) in the soil without crude oil treatment germinated but only 3.4 of the 10 seeds subjected to 10.0% w/w of the oil germinated showing a significant reduction with increasing oil level in soil. These observations are in agreement with prior reports

**Table 3.** Germination characteristics of *S. siamea* as influenced by crude oil in soil

Oil in soil (% w/w)	% germination	Days to germination	Rate of germination
0.01	98.42	5.4	9.8
2.0	94.87	2.8	8.6
4.0	63.58	5.7	6.7
8.0	38.70	6.9	5.1
10%	22.67	7.5	3.4

Means with same letters within the same column are not significantly different from each other at  $P \geq 0.05$  using the Fisher's least significant Difference (LSD).

of oil and other crops including *Leucaena leucocephala* by (Agbogidi et al., 2005); *Ricinus cummunis* (Vwioko and Fashemi, 2005) and Sharfi et al. (2007) on six plant species. According to Agbogidi and Ejemete (2005); Ogbo (2009); Vwioko et al. (2006) and Agbogidi and Egbuchua (2010), reduced germination percentage, delayed germination, growth reduction could be due to oxygen tension that has been created following the high biological oxygen demand (BOD) as a result of the crude oil addition to soil. The presence of heavy metals and other toxic oil components could have also endangered the life and growth activities of plant. Reduction in seedling emergence of plants grown in crude oil contaminated soils had been reported by Anoliefo and Edegbai (2000); Anoliefo and Vwioko (2001) and Agbogidi and Dolor (2007).

## Conclusion

This study has shown that soil contamination with crude oil has highly significant effects of reducing the seedling emergence in *S. siamea*. This study has important ecology relevance especially in the Niger Delta areas of Nigeria. *S. siamea* can serve as a bioindicator in crude oil contaminated soil and could be useful species for phytoremediation studies.

## REFERENCES

- Agbogidi OM, Ejemete OR (2005). An assessment of the effects of crude oil pollution on soil properties, germination and growth of *Gambaya albida* (L.). *Uniswa Res, J. Agric, Sci. Technol.*, 8 (2): 148-155.
- Agbogidi OM, Nweke FU, Eshegbeyi OF (2005). Effects of soil pollution by crude oil on seedling growth of *Leucaena leucocephala* (Lam.) de Witt. *Global J. Pure and Applied Sci.*, 4 (2): 453-456.
- Agbogidi OM, Eshegbeyi OF (2006). Performance of *Dacryodes edulis* (Don. G. Lam H.J.) seeds and seedlings in a crude oil contaminated soil. *J. Sustainable Forestry* 22 (3/4): 1-14.
- Agbogidi OM, Eruotor PG, Akparobi SO (2006). Effects of soil contaminated with crude oil on the germination of maize (*Zea mays* L). *Nig. J. Sci. Environ.*, 5: 1-10.
- Agbogidi OM, Dolor DE (2007). An assessment of the growth of *Irvingia gabonensis* (Aubry-Lecomte Ex O' Porte) Bail seedlings as influenced by crude oil contamination of soil. *Asian J Plant Sci.*, 2: 1287-1292.
- Agbogidi OM, Eruotor PG, Akparobi SO, Nnaji GU (2007). Evaluation of crude oil contaminated soil on the mineral nutrient elements of maize (*Zea mays* L). *J. Agronomy* 6(1): 188-193.
- Agbogidi OM, Egbuchua CO (2010). Heavy metal concentrations of soil contaminated with spent engine oil in Asaba, Delta State. *Acta Agronomica Nigeriana* 10 (1): 65-69.
- Anoliefo GO, Vwioko DE (1995). Effect of spent lubricating oil on the growth of *Capsicum annum* L. *Lycopersicon esculentum* Miller. *Environmental Pollution* 88:361-364.
- Anoliefo GO, Edegbai BO (2000). Effect of spent engine oil as a soil contaminant on the growth of two egg plant species, *Solanum melongena* L. and *S. incanum* L. *J. Agric., Forestry Fish.*, 1:21-25.
- Anoliefo GO Vwioko DE (2001). Tolerance of *Chromolaena odorata* (L) K. and R. grown in soil contaminated with spent lubricating oil. *J. Tropical Biosci.*, 1(1):20-24.
- Asaba Meteorological Station (2010). National Meteorological Bulletin 2007 Lagos, Nigeria.
- Dalziel JM (1987). The useful plants of west tropical Africa. Longman Publishers, United Kingdom.
- Module 9: Species fact sheets (2<sup>nd</sup> edition), Bangkok, Thailand: Winrock International. Pp.127
- Gutteridge RC (1997). *Senna siamea* (Larnk) Irwin and Bameby. In: Faridah, H.I. and Van der Maesen, J.G. (eds.). Plant resources of South-East Asia. No 11. Auxiliary plants. Backhuys Publishers. Leiden, the Netherlands, Pp 232-236.
- Keay RWJ, Onochie CFA, Stanley OP (1989). Nigerian trees. Oxford University Press, New York
- Ogbo EM (2009). Effects of diesel fuel contamination on seed germination of four crop plants—*Arachis hypogaea*, *Vigna unguiculata*, *Sorghum bicolor* and *Zea mays*. *Afri J. Biotechnol.*, 8 (2): 250-253.
- Okafor JC (1980). Edible indigenous woody plants in the rural economy of the Nigerian rainforest zone. *Forest*

Ecol. Manag., 3:45-55.

Ogri OR (2001). A review of the Nigerian petroleum industry and the associated environmental problems. *The Environmentalist* 21:11 – 21.

Sharifi M, Sideshow Y, Akharpour M (2007). Germination and growth of six plant species on contaminated soil with spent oil. *Int. J. Environ. Sci. Technol.*, 4(4): 463 – 470.

Siddiqui S, Adams WA (2002). The fate of diesel hydrocarbons in soils and their effects on the germination of perennial ryegrass. *Environ. Toxicol.*, 17(1): 49-62.

Statistical software (SAS) (2005). Hergen and enhanced SAS Inst. Inc. USA.

Terge K (1984). Effect of oil pollution in the germination and vegetative growth of five species of vascular plants. *Oil Petroleum J.*, 2:25-30.

Vwioko DE, Fashemi DS (2005). Growth response of *Ricinus cummunis* L. (castor oil) in spent lubricating oil polluted soil. *J. Applied Sci. Environ Manage.*, 9(2): 73 – 79.

Vwioko DE, Anoliefo GO, Fashemi SD (2006). Metal concentration in plant tissue of *Ricinus communis* L. (Castor oil) grown in soil contaminated with spent lubricating oil. *J. Applied Sci. Environ. Manage.* 10(3): 127 - 134.