

*Full Length Research Paper*

# Sera from camels slaughtered in abattoirs using the Rose Bengal plate test (RBPT) for the presence of *Brucella* antibodies

Osinbajo David-West<sup>1</sup>\*, Benjamin Elliot<sup>2</sup>, Aaron Owologba<sup>1</sup>, Ogheneborhie Micheal<sup>3</sup>,  
Chukwudi Olubuchi<sup>3</sup> and Atasariba Jackson<sup>3</sup>

<sup>1</sup>Brucella Research Section, National Veterinary Research Institute, Vom, Nigeria. <sup>2</sup>Quality Control Division, National Veterinary Research Institute, Vom, Nigeria.

<sup>3</sup>National Veterinary Research Institute, Vom, Nigeria.

Accepted 16 July, 2016

A sero prevalence study of camel brucellosis was carried out in three abattoirs of Northern Nigeria during the period of October to December, 2013. A total of three hundred and eleven (311) serum samples were collected from Kano, Sokoto and Maiduguri municipal abattoirs. The serum samples were screened using the Rose Bengal plate test with positive samples further tested with the lateral flow immunoassay. Out of 180 camel sera collected from Kano Municipal abattoir, 4 (2.2%) were positive for *Brucella* antibodies by Rose Bengal plate test of which one was confirmed by lateral flow immunoassay. Moreover, on sex distribution 3 (3.5%) female camels and one (1.1%) male camel were positive. There was no significant association with sex and prevalence of *Brucella* antibodies ( $P>0.05$ ). On age distribution, 3 (12.5%) adult camels were positive by only one young camel (0.6%) was positive. The result was statically significant ( $P<0.05$ ). From the 32 camel sera collected from Sokoto Municipal abattoir, 2 (6.3%) were positive for *Brucella* antibodies by Rose Bengal plate test of which one was confirmed by lateral flow immunoassay. All positive samples were adult (16.7%) female (11.8%) camels. There was no significant association with sex and age of camel ( $P>0.05$ ). None of the 99 serum samples collected from Maiduguri Municipal abattoir tested positive by Rose Bengal plate test. The study concluded that *Brucella* antibodies are present in camel and they were probably infected due to contact with infected cattle. The importance of these finding are discussed.

**Key words:** Sero-prevalence, brucellosis, camel, Nigeria.

## INTRODUCTION

The world's camel population was estimated to be 19 million (FAO, 2010) with 80% found in Africa. The estimated camel population in Nigeria varied from 25,

000 to 90,000 (Adamu and Ajogi, 1999) with most animals kept around Borno, Kano and Sokoto States in the Northern part of the country. Most camels in Nigeria

\*Corresponding author. E-mail: [david.west55@gmail.com](mailto:david.west55@gmail.com)

are imported from neighbouring Chad and Niger Republics where camel breeding is very common (FDLPCS, 1992; Ducrotoy et al., 2014). Due to their physiological features, camels have the ability to survive under harsh environmental conditions. They are important sources of milk, meat; leather and wool in many parts of the world (Gwida et al., 2012). They have been used as a source of investment and long-time saving, in sports, transportation and tourism (Wilson 1984; Rollefson, 2000). As a result of their unique qualities, camels were in the past thought to be resistant to diseases affecting livestock (Gauthier-Pilters and Dagg, 1981; Zaki, 1948; Bitter, 1986, Dalling et al., 1988). However, recent studies have shown that camels are susceptible to many diseases affecting livestock species including brucellosis (Abbas and Tilley, 1990; Abbas and Agab, 2002; Musa, 2008; Gwida et al., 2012).

Brucellosis is one of the most important zoonotic diseases in the world. It is an economically important disease in livestock resulting in abortion, still births, retained placenta, reduced milk yield and infertility (Godfroid et al., 2011). Although the disease has been eradicated in some developed countries, it is still a problem in developing countries. There are also reports of its re-emergence in some developed countries where the disease has been previously eradicated (Seleem et al., 2011). The disease is caused by bacteria belonging to the genus *Brucella*. Eleven species are currently recognised belonging to the genus *Brucella*; with seven species found in terrestrial animals; *Brucella abortus* (in cattle and buffalos), *Brucella melitensis* (in sheep and goats), *Brucella ovis* (in sheep), *Brucella suis* (in pigs), *Brucella canis* (in dogs), *Brucella neotomae* (in desert wood rats) and *Brucella microti* found in the common vole (Verger et al., 1987). Two species that affect marine mammals are *Brucella pinnipedialis* found in pinnepeds and *Brucella ceti* in cetaceans (Foster et al., 2007); a novel *Brucella* species *Brucella inopinata*, has been isolated in a breast implant infection (Scholz et al., 2010). Recently, *Brucella* species has been isolated from baboons and was named *Brucella papionis* (Whatmore et al., 2014). Brucellae are usually host specific meaning a particular species infect specific animal host; however, cross infection have been reported where infection can occur in animals that are not the primary host of a particular species. *B. abortus* has also been reported in sheep, goats, dogs, horses and camels (FAO, 2004) and *B. melitensis* in cattle, dogs, camels and pigs (Godfroid, 2004).

The earliest report of Brucellosis in camels was in 1931 (Solonitsuin, 1949). Since then the disease has been reported in camels in the middle East (Radwan et al., 1992; Dawood, 2008; Yawoz et al., 2012), North Africa (Musa and Shigidi, 2011; Sisay and Mekonnen, 2012) and East Africa (Wanjohi et al., 2001). In Nigeria, Brucellosis was first reported in camels by Okoh (1979) who carried out a serological survey on animals brought

to slaughter house by showing 1.0% prevalence of *Brucella* antibodies. Subsequent studies reported prevalence rates of 2% (Zaria et al., 1990) and 7.5% (Kudi et al., 1997) in an abattoir surveys carried in Maiduguri and Kano respectively and in a very recent study carried out in Sokoto municipal abattoir prevalence rate of 19.5 % was found (Junaidu et al., 2006). This study examined sera from camels slaughtered in abattoirs in three cities of Northern Nigeria for presence of *Brucella* antibodies using the Rose Bengal plate test (RBPT) as a screening test, and also used lateral flow immunoassay (LFIA) for confirmation test.

## MATERIALS AND METHODS

### Study areas and period.

The studies were carried out in 2013 between October and December in Kano, Maiduguri and Sokoto Municipal abattoirs in Northern Nigeria (Figure 1)

### Sample collection

Camels brought for slaughter were sampled based on consent and co-operation from butchers. Blood (5 ml) was collected from the jugular vein at the time of slaughter using sterile sample tubes. The tubes were labeled with the animal's sex and age (young 6 to 9 months and adult one year and above). The blood samples were allowed to clot in a slanting position, then transported to the laboratory in a leak-proof container with ice packs. They were centrifuged at 1000 rpm for 5 mins. Sera were then decanted into 5 ml plastic tubes and stored in the refrigerator at  $-20^{\circ}\text{C}$  until required for testing. The distribution of samples collected from the three abattoirs is presented in Table 1.

### Serological examination


Serum samples were tested for *Brucella* antibodies by RBPT as described by Alton et al. (1988). Samples positive by RBPT were further tested by *Brucella* LFIA done as described by Abdoel et al. (2008)

### Rose Bengal plate test

Briefly, 30  $\mu\text{L}$  of RBPT antigen (Procured from Veterinary Laboratory, United Kingdom) were added to an equal volume of serum on a ceramic tile. The sera and the antigen were mixed with an applicator stick and rocked gently. It was observed for four minutes for agglutination. Result was graded as +1, +2 or +3 based on degree of agglutination.

### *Brucella* lateral flow immuno assay

Lateral flow immunoassay was performed as described by Abdoel et al. (2008) using bovine *Brucella abortus* kit (Royal Tropical Institute, the Netherlands). The flow kit was removed from the packaging and placed on a bench top with the test window facing upwards. 5  $\mu\text{L}$  of test serum were pipetted into the round sample port on the sample pad. Immediately, 130  $\mu\text{L}$  of running fluid were added to the sample port. The sample pad was allowed standing for



**Figure 1.** Map of Nigeria showing the cities where the study was carried out.

**Table 1.** Distribution of samples collected from camel in three abattoirs of Northern Nigeria.

Variable	Total	Males	Females	Young	Adults
Maiduguri	99	39	60	85	14
Kano	180	94	86	156	24
Sokoto	32	15	17	20	12
<b>Total</b>	<b>311</b>	<b>148</b>	<b>163</b>	<b>261</b>	<b>50</b>

10 min before results were read. A positive result was indicated by the presence of a line at the test zone and a line at the control zone. A negative result is indicated by absence of a line in the test zone and the presence of a line in the control zone. Results were graded as +1 or +2 based on intensity of the red line

#### Statistical analysis

Descriptive statistics was calculated using open Epi version 3.03 to determine the level of *Brucella* sero-prevalence in relation to age

and sex of camels. P-value lower than 0.05, was considered statistically significant.

## RESULTS

The sero-prevalence of *Brucella* antibodies in camel sera collected from abattoirs in the three cities by RBPT is presented in Table 2. Out of 180 camel sera collected from Kano Municipal abattoir, 4 (2.2%) were positive for *Brucella* antibodies by RBPT of which one was confirmed by LFIa. However, on sex distribution 3 (3.5%) female camels and one (1.1%) male camel were positive. There was no significant association with sex and prevalence of *Brucella* antibodies ( $P > 0.05$ ). On age distribution, 3 (12.5%) adult camels were positive by only one young camel (0.6%) was positive. The result was statically significant ( $P < 0.05$ ). From the 32 camel sera collected from Sokoto Municipal abattoir, 2 (6.3%) were positive for *Brucella* antibodies by RBPT of which one was confirmed

**Table 2.** Sero-prevalance of brucellosis in Camel in three abattoirs of Northern Nigeria.

Variable	No	RBPT	Females	Males	Adults	Young	LFA*
Kano	180	4 (2.2%)	3 (3.5%)	1 (1.1%)	3 (12.5%)	1 (0.6%)	1
Sokoto	32	2 (6.3%)	2 (11.8%)	0 (0)	2 (16.7%)	0 (0)	1
Maiduguri	99	0	0	0	0	0	0
Total	311	6	5	1	5	1	2

\*All sample positive by LFA were adult female camels.

by LFIa. All positive samples were adult (16.7%) female (11.8%) camels. There was no significant association with sex and age of camel ( $P>0.05$ ). None of the 99 serum samples collected from Maiduguri Municipal abattoir tested positive by RBPT.

## DISCUSSION

The 2.2% sero-prevalence recorded in camel from samples collected from Kano Municipal abattoir compares with that by Okoh (1979) who reported a prevalence rate of 1% in camel slaughtered in the same area. Warsame et al. (2012) also reported a sero-prevalence rate of 2% in a study carried out in and around Dire Dawa, District of Ethiopia. However, the result was lower than 7.5% prevalence reported by Kudi et al. (1997), who carried out a similar study relating to this study in Kano. The variation in the prevalence rates could be due to the fact that Kudi et al. (1997) used micro serum agglutination test (MSAT) while this study used the RBPT. The MSAT has been reported to be a poor test in camel sera with a lot of false positive results (Abbas and Agab, 2002)

A sero-prevalence rate of 6.3% was recorded in this study in camels slaughtered at Sokoto municipal abattoir. This agrees with studies carried out by Junaidu et al. (2006) who reported a sero-prevalence rate of 11.42% in the same area. This shows that the prevalence rate has remained high in the area. This may probably be due to the fact that sick camels intended for slaughter may be easily imported into Nigeria through the Sokoto border due to its proximity to camels rearing region of Niger. This study is also in accord with Sisay and Mekonen (2012) who reported a sero-prevalence rate of 11.9% in a study carried out in Afar region of Ethiopia.

A zero percent sero-prevalence was reported in camel slaughtered in Maiduguri Municipal abattoir. The study result is quite surprising as sero-prevalence rates of 9.5 and 25.5% have been reported in the same area by Zaria et al. (1990).

The variations in the sero-prevalence rates recorded in the three cities when compared to other similar studies in the same areas could also be due to the fact that being an abattoir survey, the possibility of sampling an infected animal will depend on it being brought for slaughter.

Some infected camels may be slaughtered without being taken to the abattoir or sold out.

In Nigeria, the RBPT is widely used as a screening test for brucellosis because of its simplicity, rapidity and field suitability. However, result of RBPT requires an alternative test by complement fixation test (CFT) or enzyme linked immunosorbent assay (ELISA) as recommended by OIE (2009). These tests (CFT and ELISA), require specialized training and expensive laboratory equipment and reagents. The LFA has several practical advantages which include the fact that the use of the LFA does not require specific training, expertise, electricity or expensive equipment, and the test devices may be stored without the need for refrigeration. The test results are obtained almost instantaneously and by visual inspection with the unaided eye (Abdoel et al., 2008). The test has been compared to Competitive ELISA (cELISA), and was reported to be even more sensitive and specific than cELISA (Bronsvort et al., 2009). Its sensitivity and specificity was reported to be 87 and 97%, respectively (Bronsvort et al., 2009). It was suggested to be used as a confirmatory test.

In this study, the RBPT was used as a screening test, and LFIa as a confirmatory test. The result of the RBPT that were graded as +2 and above were all confirmed by LFIa. Since the kit had bovine specific conjugates, it is likely that the camels were infected with as a result of contact with cattle. The contact with infected cattle could be from countries where the camels were imported or with cattle within Nigeria, since camels and other livestock are usually reared together (Ducrottoy et al., 2014). Camels are not known to be primary host to *Brucella* species but infection has been reported due to contact with infected cattle or sheep (FAO, 2004). Protocol for bovine brucellosis has been used to screen camel sera used (Gwida et al., 2012). Although the LFIa produces specifically cattle, sheep, pigs and goats using host specific conjugates. It has not been used to test camel sera. The study recommends a large scale study using bovine LFIa kit in camel sera.

A high prevalence rate recorded in females in Kano (3.5%) and Sokoto (11.8%) than in males (1.1 and 0%). The results were not statistically significant in both cases ( $P<0.5$ ). The study findings are in accord with Sadiq et al. (2011) who reported a higher prevalence rate (5.5%) in female camels than (3.9%) in males in a study carried

out in Lake Chad region of Borno State, North Eastern, Nigeria and Warsame et al. (2012) who also reported a higher prevalence rate in females (1.7%) than males (1.4%) camels in another study carried out in and around Dire Dawa, Ethiopia. Dawood (2008) also reported a high prevalence rate in female (13.8%) than males (7.5%) camels in South province of Jordan. Female animals are known to be the main source and foci of *Brucella* infection (Godfroid et al., 2010). Brucellae are also known to have a high affinity to the alcohol D-erythritol found in higher volume in the gravid uterus than the seminal vesicles, making the infection more common in female than male cattle (Walker, 2004).

A higher prevalence rates were also recorded in adult camels in Kano (12.5%) and Sokoto (16.7%) than in young camels (0.6 and 0%). The results were statistically significant in both cases ( $P < 0.5$ ). This is in agreement with Dawood (2008) who reported a higher prevalence rate (64.8%) in adult than (35.2%) in young camels in southern province of Jordan, and Sisay and Mekonnen (2012) also reported a higher prevalence rate (13.8%) in adult than (0) in young camels in selected district of Afar region in Ethiopia. Musa and Shigidi (2001) also reported higher prevalence range of 6.8 to 9.2% in camels 21/2 to 4 years than 4.2% in camels of 6 to 9 months old. This is not surprising since Brucellosis is mainly a disease of the reproductive tract where the organisms have a higher affinity to the reproductive organs (Acha and Szyfres, 2003). Young animals are not sexually mature and may be resistant to *Brucella* infection.

## Conclusion

The study showed that *Brucella* antibodies are present in camels in Nigeria with the infection possibly as a result of contact with infected cattle. Although there were variations in the prevalence rates when compare to similar studies in the same area, the reason could due to the fact that the present study is an abattoir survey. The LFIa kit, though produced specifically for cattle, it was able to detect the presence of *Brucella* antibodies in camel sera. The LFIa may be a good test in camels. The study recommends a large evaluation of the test in camel sera. More studies need to be carried out especially in mixed herds (camels and other livestock) to highlight the risk factor of the disease in camel. The study also recommends an officially coordinated control programmes for brucellosis in Livestock as a means of controlling the diseases in camels, since camels are spill over host

## ACKNOWLEDGEMENT

The authors wish to acknowledge the Former Executive Director, NVRI Vom, Dr. M.S. Ahmed and the Director of Research, Dr David Shamaki, for approval and support

for the studies. They also wish to thank Dr. H.L. Smits of Royal Tropical Institute, The Netherland for supplying the LFIa kits free charge. They also acknowledge all the abattoir workers in the three cities where the studies were carried, without their cooperation the studies could not have been carried out.

## Conflict of interest

The authors have not declared any conflict of interest.

## REFERENCES

- Abbas B, Agab H (2002). A review of camel Brucellosis. *Prev. Vet. Med.* 55:47-56.
- Abbas B, Tilley P (1990). Pastoral management for protection of ecological balance in Halaib District, Red Sea Province, Sudan. *Nomad. People* 29:77-86.
- Abdoel T, Dias IT, Cardoso R, Smits HL (2008). Simple and rapid test for brucellosis diagnosis in livestock. *Vet. Microbiol.* 130:312-319.
- Acha NP, Szyfres B (2003). Zoonoses and communicable diseases common to man and animals. Vol 1, Third edition. Pan American Health Organization (PAHO) Washington, D.C.
- Adamu NB, Ajogi I (1999). Serological investigation of Camels (*Camelus dromedarius*) slaughtered at Kano municipal abattoir for evidence of brucellosis. *Trop. Vet.* 18:45-48.
- Alton GG, Jones, LM, Angus RD, Verger JM (1988). Techniques for the Brucellosis Laboratory. Institute National de la Vacherche Agronomique, (INRA) Paris. 1<sup>st</sup> Edition. pp. 63-129.
- Bitter H (1986). Disease resistance in dromedaries with particular reference to *Trypanosoma evansi* infection. Inaugural Dissertation. tierartliche Hochschule, Hanover, Germany. pp. 24-150
- Bronsvort BM, Koterwas B, Land F, Handel IG, Tucker J, Morgan KL, Tanya VN, Abdoel TH, Smits HL (2009). Comparison of a flow assay for brucellosis antibodies with the reference cELISA test in West African *Bos indicus*. *PLoS one* 4:e5221
- Dalling T, Robertson A, Boddie G, Spruell J (1988). Diseases of camels. In: *The Int. Encyclopedia for Vet. Med.* Edingburgh, U.K; W. Green and Son. P.585.
- Dawood HA (2008). Brucellosis in Camels (*Camelus dromedarius*) in the Southern province of Jordan. *Am. J. Agric. Biol. Sci.* 3(3):623-62.
- Ducrottoy MJ, Bertu WJ, Ochoi RA, Gusi AM, Bryssinckx, W, Sue W, Ignacio M (2014). Brucellosis as an Emerging Threat in Developing Economies: Lessons from Nigeria. *PLoS Negl. Trop. Dis.* 8(7).
- Food and Agriculture organization (2004). Bovine brucellosis. In: *Animal health and disease cards.* pp. 1-18.
- Food and Agricultural Organization (2010). FAO statistics Division. FAOSTAT. Available at: <http://www.fao.org/statistics/en/>.
- Federal Department of Livestock and Pest Control Services. Nigerian Livestock Resources (1992). National Synthesis. Report by Resource Inventory and Management Limited, to FDLPCS, Abuja, Nigeria. 2:440p
- Foster G, Osterman BS, Godfroid J, Jacques I, Cloeckert A (2007). *Brucella ceti* sp. nov. and *Brucella pinnipedialis* sp. nov. for *Brucella* strains with cetaceans and seals as their preferred hosts. *Int. J. Syst. Evol. Microbiol.* 57:2688-2693
- Gauthier-Pilters H, Dagg AI (1981). *The Camel.* University of Chicago Press, Chicago.
- Godfroid J, Scholz HC, Barbier T, Nicolas C, Wattiau P, Fretin, D, Whatmore AM, Cloeckert A, Blasco, JM, Moriyon I, Saegerman C, Muma JB, AlDahouk S, Neubauer H, Letesson JJ (2011). Brucellosis at the animal/ ecosystem/ human interface at the beginning of the 21<sup>st</sup> century. *Prev. Vet. Med.* 102:118-131s
- Godfroid J (2004). *Brucella melitensis* infection. In: Coetzer JAW, Tustin RC (Eds). *Infectious diseases of livestock.* Cape town, South Africa, Oxford University press. pp. 1535-1539.
- Godfroid J, Nielsen K, Saegerman C (2010). Diagnosis of Brucellosis in Livestock and wildlife. *Croatian Med. J.* 51:296-305

- Gwida M, El- Gohary, A, Melzer F, Khan I, Rösler U, Neubauer H (2012). Brucellosis in Camels. Res. Vet. Sci. 92:351-355
- Junaidu AU, Oboegbulem SI, Sharubutu GH, Daneji IA (2006). Brucellosis in Camels (*camelus dromedaries*) slaughtered in Sokoto North Western Nigeria. Anim. Prod. Res. Adv. 2(3):11-13.
- Kudi AC, Kalla DJU, Kudi MC, Kapio GI (1997). Brucellosis in Camels. J. Arid Environ. 37(2):413-417.
- Musa MT, Eisa MZM, El Sanousi EM, Abdel Wahab MB, Perret L (2008). Brucellosis in Camels (*camelus dromedarius*) in Dafur, Western Sudan. J. Comp. Pathol. 138:151-155.
- Musa MT, Shigidi MTA (2001). Brucellosis in camels in Intensive Animal breeding areas of Sudan. Implications in Abortion and early life infections. Revue. Elev. Med. Vet. Pays Trop. 54(1):11-15.
- OIE, (Office International de Epizootics) (2009). Bovine Brucellosis. Manual of Diagnostic Tests and Vaccines for Terrestrial Animals (Chapter 2.4.3).
- Okoh AEJ (1979). A survey of brucellosis in camels in Kano, Nigeria. Trop. Anim. Health Prod. 11:231-314.
- Radwan AI, Bekairi SI, Prasad PV (1992). Serological and bacteriological studies of brucellosis in camels in central Saudi Arabia. Rev. Sci. Technol. off. Int. Epiz 11(3):837-844.
- Rollefson K (2000). The Camel and human society. In: Gahlot TK (Ed). Selected topics on Camelids. The Camelids Publishers, Bikaner, India. pp. 1-17.
- Sadiq MA, Ajogi I, Bale JOO, Mosimabale FB, Tijjani AN, Kaikabo AA (2011). Serological Survey of Antibodies against *Brucella* organisms in One humped Camel (*Camelus dromedarius*) herds in the Lake Chad Area of Borno State, North Eastern Nigeria. Niger. Vet. J. 32(1):45-48.
- Scholz HC, Hubalek Z, Sedlacek I, Vergnaud G, Tomaso H, Al Dahouk S, Melzer F, Kampfner P, Neubauer H, Cloeckert A, Maquart M, Zygmunt MS, Whatmore AM, Falsen E, Bahn P, Gollner C, Pfeffer M, Huber B, Busse HJ, Nockler K (2010). *Brucella inopinata* sp. Nov.; isolated from a breast implant infection. Int. J. Syst. Evol. Microbiol. 60:801.
- Seleem MN, Boyle SM, Sriranganathan N (2011). Brucellosis: A re-emerging zoonosis. Vet. Microbiol. 140:392-398.
- Sissay WZ, Mekonnen H (2012). Seroprevalence of *brucella* infection in camel and its public health significance in selected district of afar region, Ethiopia. J. Environ. Occup. Sci. 1(2):91-98
- Solonitsuin MO (1949). Brucellosis in camels. Veterinariya Moscow 26:16-21.
- Verger JM, Grimont F, Grimont PA, Grayon, M1 (1987). Taxonomy of the genus *Brucella*. Ann. Pasteur. Microbiol. 138:235-238.
- Walker RL (2004). *Brucella*. In: Vet. Microbiol. by Hirsh DC Maclachlan NJ Walker RL (Eds). Blackwell publishing company. pp. 105-112
- Wanjohi M, Gitao CG, Bebora L (2011). The prevalence of *Brucella* species in camels Milk marketed from North Eastern Province, in Kenya. Res. Opin. Anim. Vet. Sci. 2(7):425-434.
- Warsame I, Alemu S, Temesgen W, Molla W (2012). Seroprevalence and Associated Risk factors of Camel (*Camelus dromedaries*) Brucellosis in and around Dire, Dawa, Ethiopia. Global Vet. 8(5):480-483
- Whatmore AM, Davison N, Cloeckert A, Al DS, Zygmunt MS, Brew SD, Perrett LL, Koylass MS, Vergnaud G, Quance C, Scholz HC, Dick EJ Jr, Hubbard G, Schlabritz-Loutsevitch NE (2014). *Brucella papionis* sp. nov., isolated from baboons (*Papio* spp.). Int. J. Syst. Evol. Microbiol. 64:4120-8
- Wilson TR (1984). The Camel. Longman, New York.
- Yawoz M, Jaafar SE, Salih, AI, Abdullah MH, (2012). A serological survey of brucellosis in Camels in South of Kirkuk, Iraq. Iraqi J. Vet. Sci. 26(2):105-107.
- Zaki R (1948). *Brucella* infection among ewes, camels and pigs in Egypt. J. Comp. Pathol. 58:145-151.
- Zaria LT, Egwu GO, Nawathe DR, Ojeamiren ME (1990). A preliminary report on the seroprevalence of brucellosis in Camels (*camelus dromedarius*) in Maiduguri. Zaria Vet. 5(1):64-65.