

Full Length Research Paper

Production and characterization of protease enzyme from *Bacillus laterosporus*

B. Usharani¹ and M. Muthuraj^{2*}

¹Department of Biotechnology, Madras University, Guindy, Chennai, Tamil Nadu, India.

²Department of Microbiology, Government Hospital for Chest Diseases, Puducherry, India.

Accepted 8 March, 2016

Production and partial purification of protease enzyme by *Bacillus laterosporus* was the aim of this study. *B. laterosporus* was allowed to grow in shake flask broth culture for purpose of inducing protease enzyme. The protease enzyme was purified by ammonium sulfate precipitation followed by dialysis and further concentrated by Amicon tubes. After concentration, the protein was subjected to 12% Zymogram gel with gelatin and the molecular weight of the protease enzyme was 15 kDa. The protease activity increased as there was increase in enzyme concentration; optimum substrate concentration (starch) was 1.0% (w/v); an optimum incubation temperature was 40°C. Purified protease enzyme had a maximum activity at pH 7.0 of phosphate buffer and the optimum incubation time was 24 h. The protease isolated from *B. laterosporus* is a mesophilic protease. It is stable at pH 7, at 40°C temperature, and this enzyme can be exploited commercially.

Key words: *Bacillus laterosporus*, PMSF, -mercaptoethanol, protease, fermentation.

INTRODUCTION

Proteases are essential constituents of all forms of life on earth including prokaryotes, fungi, plants and animals. Proteases are highly exploited enzymes in food, leather, detergent, pharmaceutical, diagnostics, waste management and silver recovery (Babu et al., 2005). Proteases [serine protease (EC. 3.4.21), cysteine (thiol) protease (EC 3.4.22), aspartic proteases (EC 3.4.23) and metalloprotease (EC 3.4.24)] constitute one of the most important groups of industrial enzymes, accounting for about 60% of the total enzyme market (Nunes et al., 2001; Singh et al., 2001; Zeikus et al., 1998). Among the various proteases, bacterial proteases are the most significant, compared with animal and fungal proteases (Ward et al., 1995) and among bacteria, *Bacillus* sp are specific producers of extra-cellular proteases (Priest et al., 1999). These enzymes have wide industrial application, including pharmaceutical industry, leather industry, manufacture of protein hydrolyzates, food industry and waste processing industry (Pastor et al., 2001).

Thermostable proteases are advantageous in some applications because higher processing temperatures can be employed, resulting in faster reaction rates, increase in the solubility of nongaseous reactants and products and reduced incidence of microbial contamination by mesophilic organisms. Proteases secreted from thermophilic bacteria are thus of particular interest and have become increasingly useful in a range of commercial applications (Rahman et al., 1994; Adams et al., 1998).

In this study, we describe the selection of medium components for the optimal production of extra cellular protease by thermophilic *Bacillus laterosporus* and characterization of the enzyme.

MATERIALS AND METHODS

Micro organism and Maintenance of Culture

The bacterial strain used in this study was the thermophilic *Bacillus laterosporus*, previously isolated from the palletized feed of prawn. The bacterial strain was cultured in 1% LB (Himedia) broth at 40°C for 24 h and then autoclaved at 121°C for 15 min. A loopful of autoclaved broth was plated on nutrient agar plates and incubated

*Corresponding author. E-mail: muthu_ggcd@yahoo.com.

at 37°C for 24 - 48 h. The culture was routinely maintained on Nutrient agar slants. The organism was subculture for every month. The inoculum was prepared by dispersing the loopful of bacterial culture from a week- old nutrient agar slant culture in 1% LB broth solution with a sterile inoculation loop.

Biochemical characterization of bacterial isolate

Sub culturing was done for single colony isolation. A loopful of seed culture from overnight sample was streaked on to nutrient agar plate and was incubated for 24 h at 37°C. Various biochemical study was done as per the Bergey's manual of determinative bacteriology (Holt et al., 1994) to identify the bacterial isolate.

Qualitative estimation of protease enzyme

A loopful of strain was dispersed in 9 ml of 1% LB broth medium and the diluted sample of *B.laterosporus* was plated onto pre sterilized skim milk agar plates (Hi media – M763). The plates were incubated at 37°C for 24 h to check the proteolytic activity. For maximum production of the protease enzyme, we checked the activities of extra cellular protease enzyme production in the culture supernatant at different stages of *B.laterosporus* growth.

Production of protease in shake flask fermentation

The bacterium strain was cultured in 2.5 l conical flask containing the following ingredients: Yeast extract (0.5 g), casein (0.1 g), KH_2PO_4 (0.2 g), Na_2CO_3 (1.0 g), distilled water (100 ml) and pH, 7.0. Inoculation was performed with 10% (v/v) seed culture of 1 day grown in 1% LB broth medium. The fermentation conditions were maintained at 37°C, 50 rpm agitation for 2 - 4 days. The culture broth was harvested and centrifuged at 8,000 rpm for 20 min at room temperature. Cell free supernatants were used for measuring protease activity.

Determination of protease activity

The activity of protease was assessed in triplicate by measuring the release of trichloroacetic-acid soluble peptides from 0.2% (w/v) azocasein in 50 mM HEPES/NaOH buffer (pH 7.5) at 50°C for 10 min. The 1 ml reaction was terminated by the addition of 0.5 ml of 15% trichloroacetic acid and then centrifuged at 10,000 g for 10 min, after cooling. One unit (U) enzyme activity was defined as the amount of enzyme required to produce an increase in absorbance at 420 nm equal to 1.0 in 60 min under the assay conditions (Janssen et al., 1994). Protein was measured by the method of Lowry, as modified by Petterson (Peterson et al., 1997).

Enzyme purification

The bacterial strain was grown for 48 h at 37°C and the culture broth was harvested and centrifuged at 8,000 rpm for 20 min at room temperature. Cell free supernatant was fractionated by ammonium sulfate precipitation. All subsequent steps were carried out at 4°C. The protein pellet obtained after precipitation with ammonium sulphate was resuspended in 0.1M Tris-HCl buffer, pH 7.8, and dialyzed against the same buffer. After dialysis, the contents were collected in the Amicon tubes Ultra-5 kDa (Millipore) and centrifuge at 10,000 rpm for 20 min at 27°C (Ellaiah et al., 2002). The purified enzyme was subjected to 12% Zymogram gel with gelatine.

Effect of culture conditions on enzyme production

The effects of carbon sources 1% (w/v) on enzyme secretion were investigated replacing trisodium citrate by glycerol, D(+) galactose, lactose, sucrose, maltose, starch, D(+) glucose, D(+) manose, L (+) arabinose, casein, D(+) xylose and citric acid. Different nitrogen sources including Soya oil, Casein, Wheat bran, Tryptone, Skim milk NH_4NO_3 , peptone, yeast extract, meat extract, casein, $(\text{NH}_4)_2\text{SO}_4$, $(\text{NH}_4)_2\text{HPO}_4$, NH_4Cl , KNO_3 , urea and ammonium citrate were employed in preliminary studies to determine growth and production of extra cellular protease.

Effect of pH on activity and stability of protease

The optimum pH was determined with azocasein 1% (w/v) as substrate dissolved in different buffers (citrate phosphate, pH 5-6, sodium phosphate, pH 7.0, Tris-HCl, pH 8.0 and glycine NaOH, pH 9 - 13). The effect of pH on enzyme stability was determined by pre-incubating the enzyme without substrate at different pH values (5.5 - 9.0) for 24 h at room temperature and measuring the residual activity at 40°C. Reaction mixtures were incubated at 40°C for 2 h and the relative activity of the enzyme was measured at standard assay conditions

Effect of temperature on activity and stability

The effect of temperature on the enzyme activity was determined by performing the standard assay procedure at pH 7.5 within a temperature range from 30 to 100°C. Thermostability was determined by incubation of crude enzyme at temperatures ranging from 30 - 100°C for 2 h in a constant-temperature water bath and the relative protease activities were assayed at standard assay conditions.

Effect of inhibitors and chelators on protease activity

The effect of various protease inhibitors (5 mM) such as serine inhibitors [phenylmethylsulphonyl fluoride (PMSF)] and -mercaptoethanol [- ME] and a chelator of divalent cations (ethylene diamine tetra acetic acid [EDTA]) were determined by the addition of the corresponding inhibitors and chelators at a final concentration of 1.0 mM to the reaction mixture and assayed under standard condition. The relative protease activity was measured (Rawling et al., 1994)

Effect of metal ions on protease activity

The effect of different metal ions on protease activity was determined by the addition of the corresponding ion at a final concentration of 1.0 mM to the reaction mixture, and assayed under standard conditions. The enzyme assay was carried out in the presence of KCl, CaCl_2 , MgSO_4 , FeSO_4 , CoCl_2 , ZnCl_2 , MnSO_4 , HgCl_2 , CuSO_4 , BaCl_2 and NaCl. The activity is expressed as a percentage of the activity level in the absence of metal ion. The enzyme was pre-incubated with metal ion (60°C, 5 min.) Separate blanks with individual metal ions were prepared.

RESULTS AND DISCUSSION

The isolated bacterial strain was identified as *B. laterosporus* using various biochemical tests (Table 1) as

Table 1. Biochemical characterization of *B. laterosporus*.

Gram staining	+
Rod shaped	+
Endospores produced	+
Mobility	-
Indole production	-
Methyl red	+
Voges Prokauer	-
Beta –galactosidase	-
Nitrate reduction	+
TSI	+
H ₂ S production (TSI)	-
Gas from TSI	-
Lysine decarboxylase	+
Ornithine decarboxylase	+
Arginine decarboxylase	-
Lecithinase activity	+
Acid formation from	
Arabinose	-
Xylose	-
Adonitol	-
Raminose	-
Cellobiose	-
Mellibiose	-
Saccharose	-
Raffinose	-
Trehalose	-
D-glucose	-
Lactose	-
Growth at	
37°C	+
40°C	+
45°C	+
Growth at	
5% NaCl	+
Starch hydrolysis	-
Gelatin hydrolysis	+
Catalase	+
Oxidase	+
Urease	-
Utilization of	
Esculin	-
Citrate	-
Malonate	-
Mannital	+
Tyrosine	+
Lecithinase activity	+
Resistance to Lysozyme	+

per the Bergey's manual of determinative bacteriology (Holt et al., 1994). The strain *B. laterosporus* hydrolyses the skim milk and the zone of hydrolysis on skim milk agar is shown in Figure 1. The highest activity of extra cellular protease enzyme was at 3 days as shown in Figure 2. The crude enzyme was concentrated using ultra 5kDa Amicon tube (Millipore product) and it was subjected to 12% Zymogram gel. The molecular weight was determined by interpolation from a linear semi logarithmic plot of relative molecular mass versus the Rf value (relative mobility). Depending on the relative mobility, the molecular weight of the protein band was calculated to be 15 kDa, which coincided with the band of - lactalbumin marker protein.

B. laterosporus was capable of utilizing a wide range of carbon sources. However, the best carbon sources in the present study, for protease secretion were soluble starch trisodium citrate, citric acid and glycerol (Table 2). In a similar study Johnvesly and Nailk (Adinarayana et al., 2004) showed that citric acid, soluble starch and trisodium citrate were the best carbon sources for protease production by *Bacillus* sp JB-99. According to these authors, culturing this organism in 1% glucose (w/v) repressed completely the synthesis of alkaline protease. However, in the present study, glucose was found to be a relatively good carbon source for enzyme production since moderate amount of protease activity was detected. The type of nitrogen sources also affected enzyme production. Among the various organic and inorganic nitrogen sources, the maximum enzyme activity (100%) was obtained when ammonium nitrate, wheat bran and ammonium chloride was used in the medium (Table 3). Moderate to good levels of enzyme activities were obtained when citric acid, ammonium citrate and potassium nitrate were used as nitrogen sources. When various organic nitrogen sources were tested for protease production, it was found to be increased.

The effect of different metal ions on protease is shown in Table 4. A stronger inhibitory effect was observed in the presence of KCl, BaCl₂, CuSO₄, ZnCl₂ and HgCl₂ inhibited completely the enzyme at 1 mM concentrations. The protease secreted by *Brevibacillus* (*Bacillus*) *brevis* was also inhibited by CuSO₄, ZnCl₂ and HgCl₂ (Banerjee et al., 1999). The inhibitory effect of heavy metal ions is well documented in the literature. It is known that the ions mercury, cadmium and lead react with the protein thiol groups (converting them to mercaptides), as well as with histidine and tryptophan residues. Moreover, by action of silver and mercury, the disulphide bonds were found to be hydrolytically degraded (Kumar et al., 1999). Protease activity was stimulated by MnSO₄ CoCl₂ and CaCl₂. These results suggest that these metal ions apparently protected the enzyme against thermal denaturation and played a vital role in maintaining the active conformation of the enzyme at higher temperatures (Beg et al., 2003). Similar effects of MnSO₄ on the activity of proteases were also observed by Rahman et al. (1994) and Manachinin et al. (1998).




Figure 1. Protein production in different time intervals.



Figure 2. Zone of hydrolysis of *B.laterosporus* on skim milk agar.

Table 2. Effect of the carbon sources on the production of the protease. Cells were grown in the basal medium containing 1% wheat bran and supplemented with each carbon sources at 37°C for 48 h.

S. no	Nitrogen source	Concentration (%)	Relative activity (%)
1	Glycerol	1	100
2	Galactose	1	75
3	Lactose	1	86
4	Sucrose	1	100
5	Maltose	1	89
6	Starch	1	100
7	Glucose	1	89
8	Manose	1	95
9	Arabinose	1	65
10	Casein	1	91
11	Xylose	1	76
12	Fructose	1	67
13	Trisodium citrate	1	100
14	Citric acid	1	95

Table 3. Effect of the nitrogen sources on the production of the extra cellular protease. Cells were grown in the basal medium containing 1% glucose and supplemented with each nitrogen source at 37°C for 48 h.

S. no	Nitrogen source	Concentration (%)	Relative activity (%)
1	Soya oil	1	86
2	Casein	1	82
3	Wheat bran	1	100
4	Tryptone	1	73
5	Skim milk	1	78
6	Yeast extract	1	65
7	Meat extract	1	61
8	Ammonium citrate	1	98
9	Peptone	1	45
10	NH ₄ NO ₃	1	67
11	(NH ₄) ₂ HPO ₄	1	72
12	NH ₄ Cl	1	100
13	KNO ₃	1	86
14	(NH ₄) ₂ SO ₄	1	49

Table 4. Effect of various metal ions on protease activity.

S. no	Nitrogen source	Relative activity (%)
1	KCl	7
2	CaCl ₂	153
3	MgSO ₄	69
4	FeSO ₄	57
5	CoCl ₂	109
6	ZnCl ₂	12
7	MnSO ₄	138
8	HgCl ₂	0
9	CuSO ₄	11
10	BaCl ₂	9
11	NaCl.	29
12	Control	100

The protease activities were assayed at different temperatures ranging from 30 to 90°C at a constant pH of 7.0 (Figures 3 and 4). Enzyme activity increased with temperature within the range of 30 to 40°C. A reduction in enzyme activity was observed at values above 40°C. The optimum temperature of this protease was 40°C. The thermo stability of the protease was examined by measuring the remaining activities at 40°C, after incubation of the enzyme without substrate at various temperatures between 30 and 90°C for 2 h. Thermo stability profile indicated that the enzyme was stable at 40°C for 1 h and gradually decreased after 1 h when incubation increased. A pH range between 6 and 11 was used to study the effect of pH on protease activity (Figure 5). Optimum pH was found to be 7.0. At pH 6.0, only 63% of the maximum enzyme activity was obtained, increasing to 87% at pH 7.0 and the enzyme activity declined

gradually after pH 7.0. Inhibition studies primarily give an insight into the nature of an enzyme, its cofactor requirements and the nature of the active centre. The effect of different inhibitors on the enzyme activity of the crude protease was studied (Figure 6). Of the inhibitors tested (at 5 mM concentration), EDTA was able to inhibit the protease considerably, while β mercaptoethanol and PMSF exhibited 18 and 13% inhibition, respectively. The protease isolated from *B. latrosporus* is a mesophilic protease. It is stable at pH 7, at 40°C temperatures and this enzyme can be exploited commercially.

ACKNOWLEDGMENTS

The authors are grateful to the authorities of Biocorporals, Chennai, Tamil Nadu, India for their timely




Figure 3. Effect of temperature on protease enzyme activity.




Figure 4. Stability of enzyme activity at 40°C.




Figure 5. Effect of pH on enzyme activity.




Figure 6. Effect of inhibitors on protease enzyme activity.

supply of materials all through the study.

REFERENCE

- Adams MWW, Kelly RM (1998). Finding and using thermophilic enzymes. *Trends Biotechnol.* 16: 329-332.
- Adinarayana K, Ellaiah P (2004). Investigations on alkaline protease production with *B subtilis* PE -11 immobilised in calcium alginate gel beads. *Biochem Process.* 39: 1331-1339.
- Babu NKS, Lakshmi KD (2005). Optimization of thermostable alkaline protease production from species of *Bacillus* using rice bran. *Afr. J. Biotechnol.* 4(7): 724-726.
- Banerjee UC, Sani RK, Azmi W, Soni R (1999). Thermostable alkaline protease from *Bacillus brevis* and its characterization as a laundry detergent additive. *Proc. Biochem.* 35: 213-219.
- Beg QK, Gupta R (2003). Purification and characterization of an oxidation stable, thiol-dependent serine alkaline protease from *Bacillus mojavensis*. *Enz. Microbial. Techn.* 32: 294-304.
- Ellaiah P, Adinarayana K, Pardhasaradhi SV, Srinivasulu B (2002). Isolation of alkaline protease producing bacteria from Visakhapatnam soil. *Ind. J. Microbiol.* 42: 173-175.
- Holt JG, Krieg NR, Sneath PHA, Staley JT (1994). *Bergey's Manual of Determinative Bacteriology*. Nineteenth edition, Williams and Wilkins company, Baltimore, MD, USA pp. 255-273.
- Janssen PH, Peek K, Morgan HW (1994). Effect of culture conditions on the production of an extracellular proteinase by *Thermus* sp. Rt41A. *Appl. Microbiol. Biotechnol.* 41: 400-406.
- Kumar CG, Tiwari MP, Jany KD (1999). Novel alkaline serine proteases from alkalophilic *Bacillus* spp: Purification and some properties. *Proc. Biochem.* 34: 441-449.
- Manachini PL, Fortina MG, Parini C (1988). Thermostable alkaline protease produced by *Bacillus thermoruber* a new species of *Bacillus*. *Appl. Microbiol.* 28: 409-413.
- Nunes AS, Martins MLL (2001). Isolation, properties and kinetics of growth of a thermophilic *Bacillus*. *Braz. J. Microbiol.* 32: 271-275.
- Pastor MD, Lorda GS, Balatti A (2001). Protease obtention using *Bacillus subtilis* 3411 and amaranth seed meal medium at different aeration rates. *Braz. J. Microbiol.* 32: 1-8.
- Peterson GL (1977). A simplification of the protein assay method of Lowry et al. which is more generally applicable. *Analytical Biochem.* 83: 346-356.
- Priest FG (1977). Extracellular enzyme synthesis in the genus *Bacillus*. *Bacteriol. Rev.* 41: 711-753.
- Rahman RNZA, Razak CN, Ampom K, Basri M, Yunus WMZW, Salleh AB (1994). Purification and characterization of a heat stable alkaline protease from *Bacillus stearothermophilus* F1. *Appl. Microbiol. Biotechnol.* 40: 822-827.
- Rahman RNZA, Razak CN, Ampom K, Basri M, Yunus WMZW, Salleh AB (1994). Purification and characterization of a heatstable alkaline protease from *Bacillus stearothermophilus* F1. *Appl. Microbiol. Biotechnol.* 40: 822-827.
- Rawling ND, Barret A (1994). Families of aspartic peptidases, and those of unknown mechanism. *Meth. Enzymol.* 248:105-120.
- Rawling ND, Barret A (1994). Families of cysteine peptidases. *Meth. Enzymol.* 244: 461-486.
- Rawling ND, Barret A (1994). Families of serine peptidases. *Meth. Enzymol.* 244: 18-61.
- Singh J, Batra N, Sobti CR (2001). Serine alkaline protease from a newly isolated *Bacillus* sp. SSR1. *Proc. Biochem.* 36: 781-785.
- Ward OP (1995). Proteolytic enzymes In: M. Moo-Young Editor, *Comprehensive Biotechnol.* 3: 789-818.
- Zeikus JG, Vieille C, Savchenko AT (1998). Biotechnology and structure-function relationship. *Extremophiles* 1: 2-13.