

Review

A review on trends in production of secondary metabolites from higher plants by *in vitro* tissue, organ and cell cultures

S. Karuppusamy

Department of Botany, The Madura College, Madurai, Tamil Nadu – 625 011, India. E-mail: ksamylin@yahoo.co.in

Accepted 15 December, 2021

Plant cell and tissue cultures can be established routinely under sterile conditions from explants, such as plant leaves, stems, roots, meristems etc for both the ways for multiplication and extraction of secondary metabolites. Strain improvement, methods for the selection of high-producing cell lines, and medium optimizations can lead to an enhancement in secondary metabolite production. However, most often trials with plant cell cultures fail to produce the desired products. In such cases, strategies to improve the production of secondary metabolites must be considered. One of the main problems encountered is the lack of basic knowledge of the biosynthetic routes, and mechanisms responsible for the production of plant metabolites. Where the productivity of the desired metabolites is limited by the lack of particular precursors, biotransformation using an exogenous supply of biosynthetic precursors, genetic manipulation and metabolic engineering may improve the accumulation of compounds. Feedback inhibition of metabolic enzymes as well as inhibition of membrane transport can be eliminated by the accumulation of synthesized products in a second phase introduced into the aqueous medium. Organ cultures and *in vitro* biomass production often have sites of synthesis and storage of secondary metabolites in separate compartments. Elicitors, compounds triggering the formation of secondary metabolites, can be abiotic or biotic. Natural elicitors include polysaccharides such as pectin and chitosan, which are also used in the immobilization and permeabilization of plant cells. Immobilization with suitable bioreactor system provides several advantages, such as continuous process operation, but for the development of an immobilized plant cell culture process, natural or artificially induced secretion of the accumulated product into the surrounding medium is necessary. The present review highlights the nature, applications, perspective and scale up methods for the production of valuable secondary metabolites *in vitro*.

Key words: Plant cell cultures, secondary metabolites, biotransformation, biosynthetic pathways, elicitation, immobilization, bioreactors.

INTRODUCTION

Since the early days of mankind, plants with secondary metabolites have been used by humans to treat infections, health disorders and illness (Wyk and Wink, 2004). Only during the last 100 years have natural products been partly replaced by synthetic drugs, for which plant structures were a lead in many instances, for example salicylic acid and aspirin. Many higher plants are major sources of useful secondary metabolites which are used in pharmaceutical, agrochemical, flavor and aroma industries. The search for new plant-derived chemicals should thus be a priority in current and future efforts towards sustainable conservation and rational utilization

of biodiversity (Philipson, 1990). Biotechnological approaches, specifically plant tissue culture plays a vital role in search for alternatives to production of desirable medicinal compounds from plants (Rao and Ravishankar, 2002). On a global scale, medicinal plants are mainly used as crude drugs and extracts. Several of the more potent and active substances are employed as isolated compounds, including many alkaloids such as morphine (pain killer), codeine (antitussive), papaverine (phosphodiesterase inhibitor), ephedrine (stimulant), ajmaline (antirhythmic), quinine (antimalarial), reserpine (antihypertensive), galanthamine (acetylcholine esterase inhi-

bitor), scopolamine (travel sickness), berberine (psoriasis), caffeine (stimulant), capsaicin (rheumatic pains), colchicines (gout), yohimbine (aphrodisiac), pilocarpine (glaucoma), and various types of cardiac glycosides (heart insufficiency) (Wink et al., 2005).

The capacity for plant cell, tissue, and organ cultures to produce and accumulate many of the same valuable chemical compounds as the parent plant in nature has been recognized almost since the inception of *in vitro* technology. The strong and growing demand in today's marketplace for natural, renewable products has refocused attention on *in vitro* plant materials as potential factories for secondary phytochemical products, and has paved the way for new research exploring secondary product expression *in vitro*. However, it is not only commercial significance that drives the research initiatives. The deliberate stimulation of defined chemical products within carefully regulated *in vitro* cultures provides an excellent forum for in-depth investigation of biochemical and metabolic pathways, under highly controlled microenvironmental regimes.

Plant-produced secondary compounds have been incorporated into a wide range of commercial and industrial applications, and fortuitously, in many cases, rigorously controlled plant *in vitro* cultures can generate the same valuable natural products. Plants and plant cell cultures have served as resources for flavors, aromas and fragrances, biobased fuels and plastics, enzymes, preservatives, cosmetics (cosmeceuticals), natural pigments, and bioactive compounds. There is a series of distinct advantages to producing a valuable secondary product in plant cell culture, rather than *in vivo* in the whole crop plant. These include: a) Production can be more reliable, simpler, and more predictable, b) Isolation of the phytochemical can be rapid and efficient, as compared to extraction from complex whole plants, c) Compounds produced *in vitro* can directly parallel compounds in the whole plant, d) Interfering compounds that occur in the field-grown plant can be avoided in cell cultures, e) Tissue and cell cultures can yield a source of defined standard phytochemicals in large volumes, f) Tissue and cell cultures are a potential model to test elicitation, g) Cell cultures can be radio labeled, such that the accumulated secondary products, when provided as feed to laboratory animals, can be traced metabolically.

Secondary products in plant cell culture can be generated on a continuous year-round basis; there are no seasonal constraints. Production is reliable, predictable, and independent of ambient weather. At least in some cases, the yield per gram fresh weight may exceed that which is found in nature. Disagreeable odours or flavors associated with the crop plant can be modified or eliminated *in vitro*. Plant cell culture eliminates potential political boundaries or geographic barriers to the production of a crop, such as the restriction of natural rubber production to the tropics or anthocyanin pigment production to climates with high light intensity. When a

valuable product is found in a wild or scarce plant species, intensive cell culture is a practical alternative to wild collection of fruits or other plant materials. Extraction from the *in vitro* tissues is much simpler than extraction from organized, complex tissues of a plant. Plant tissue culture techniques offer the rare opportunity to tailor the chemical profile of a phytochemical product, by manipulation of the chemical or physical microenvironment, to produce a compound of potentially more value for human use.

While research to date has succeeded in producing a wide range of valuable secondary phytochemicals in unorganized callus or suspension cultures, in other cases production requires more differentiated microplant or organ cultures (Dörnenberg and Knorr, 1997). This situation often occurs when the metabolite of interest is only produced in specialized plant tissues or glands in the parent plant. A prime example is ginseng (*Panax ginseng*). Since saponin and other valuable metabolites are specifically produced in ginseng roots, root culture is required *in vitro*. Similarly, herbal plants such as *Hypericum perforatum* (St. John's wort), which accumulates the anti-depressant hypericins and hyperforins in foliar glands, have not demonstrated the ability to accumulate phytochemicals in undifferentiated cells (Smith et al., 2002). As another example, biosynthesis of lysine to anabasine occurs in tobacco (*Nicotiana tabacum*) roots, followed by the conversion of anabasine to nicotine in leaves. Callus and shoot cultures of tobacco can produce only trace amounts of nicotine because they lack the organ-specific compound anabasine. In other cases, at least some degree of differentiation in a cell culture must occur before a product can be synthesized (e.g., vincristine or vinblastine from *Catharanthus roseus*). Reliance of a plant on a specialized structure for production of a secondary metabolite, in some cases, is a mechanism for keeping a potentially toxic compound sequestered. The three long-standing, classic examples of commercially viable production of a secondary metabolite *in vitro* — ginseng saponines, shikonin, and berberine — each feature products that have diversified uses, including medicinal applications. Ginseng is produced in large-scale root cultures, whereas the other two products are produced in highly colored cell cultures. A tremendous research and development effort has advanced a number of other *in vitro*-derived secondary products to semicommercial status, including vanillin and taxol production in cell cultures. In a myriad of other cases, the *in vitro* processes for secondary metabolite production have fallen far short of expectations and have never approached commercial status. Still, the arena of secondary product formation in cell cultures remains as an industrial pursuit. Engineers and biologists are currently joining forces on a global scale to develop new strategies for streamlining the critical bioprocesses.

Research efforts on a broad range of plant cell culture-derived extracts can be cited in each of these major pro-

duct categories: flavors (onion and garlic, peppermint and spearmint, fruit flavors, chocolate aroma, seaweed flavors, vanilla, celery, coffee, spice, sweeteners, and so on); edible colors for foods and medicines (mainly betalains and anthocyanins); non-food pigments for cosmetics and textiles (shikonin, berberine, and various other products); several examples of fragrances and essential oils; and bioactive natural insecticides and phytoalexins useful in current integrated pest management programs. Of course, intensive activity has centered on production of natural drugs or chemoprotective compounds from plant cell culture. Some of the most prominent pharmaceutical products in this latter category include ajmalicine (a drug for circulatory problems) from *C. roseus* and taxol (a phytochemical effective in treatment of ovarian cancer) from *Taxus* species. *In vitro* production of secondary metabolites from higher plants is surveyed and summarized in Table 1.

Organ cultures for secondary metabolite production

The technique of *in vitro* organ culture of *Fritillaria unibracteata* has been established and chemical composition of the metabolites has been confirmed. *F. unibracteata* can be rapidly propagated, directly from small cuttings of the bulb by the technique of organ culture. The cultured bulb can be harvested after a 50-day culture period in MS media supplemented with 4.44 M BA and 5.71 M IAA. The growth rate was about 30–50 times higher than that under natural wild growth conditions. The content of alkaloid and beneficial microelements in the cultured bulbs were higher than found in the wild bulb. It is therefore possible to establish a novel process to produce this natural plant drug in future studies and achieve scale-up for industrial production (Gao et al., 2004).

In vitro shoot multiplication of *Frangula alnus* was obtained on woody plant medium (WPM) with indole-3-acetic acid (IAA) and 6-benzylaminapurine (BAP), the highest metabolite production (1731 mg/100 g of total anthraquinone) was in the shoots grown on the MS medium with addition of 1-naphthylacetic acid (NAA) (0.1 mg l^{-1}) and thidiazuron (TDZ) (0.1 mg l^{-1}). Similarly, shoot multiplication of *Frangula rupestris* was obtained on MS medium supplemented with BAP (0.5 mg l^{-1}). The highest metabolite production of total anthraquinone was in the shoots grown on medium with 2,4-D (0.1 mg l^{-1}) and BAP (0.5 mg l^{-1}) (Kovacevic et al., 2005).

Shoot cultures of *Gentianella austriaca* (A. and J. Kerner) Dostal established from seedling epicotyls were maintained on MS medium supplemented with 2.22 M BA and 0.54 M NAA. Shoot cultures contained the same types of secondary metabolites as plants from nature. Xanthenes were the major constituents, with DMB (demethylbellidifolin), DGL (demethylbellidifolin-8-O-glucoside) and BGL (bellidifolin-8-O-glucoside) present at

roughly two times lower concentrations than in samples from nature. Secondary metabolite production was strongly affected by the presence of BA in the medium (Vinterhalter et al., 2008).

Precursor addition for improvement of secondary metabolite production

The accumulation of secondary metabolites in plants is part of the defense response against pathogenic attack, which is triggered and activated by elicitors, the signal compounds of plant defense responses (Zhao et al., 2005). Therefore, the treatment of plant cells with biotic and/or abiotic elicitors has been a useful strategy to enhance secondary metabolite production in cell cultures. The most frequently used elicitors in previous studies were fungal carbohydrates, yeast extract, MJ and chitosan. MJ, a proven signal compound, is the most effective elicitor of taxol production in *Taxus chinensis* Roxb. (Wu and Lin, 2003) and ginsenoside production in *Panax ginseng* C.A. Meyer (Yu et al., 2000; Yu et al., 2002; Kim et al., 2004; Thanh et al., 2005) cell/organ culture. In the present study, the effect of different concentrations of MJ on embryogenic cell growth and eleutheroside accumulation was tested and results revealed that addition of 200 M MJ was suitable for optimum accumulation of eleutheroside B, E, E1 and chlorogenic acid. However, addition of MJ at higher concentration (above 100 M) was detrimental for biomass accumulation. Similar to the present results, MJ inhibited the cell growth and promoted the secondary metabolite production with cell/adventitious root cultures of *Bupleurum falcatum* L. (Aoyagi et al., 2001), *Taxus* spp. (Yukimune et al., 1996; Ketchum et al., 1999) and *Panax ginseng* C.A. Meyer (Kim et al., 2004; Thanh et al., 2005). Differential accumulation of eleutherosides was observed during elicitation experiments.

In the bioreactor cultures, eleutheroside content increased significantly by elicitation of methyl jasmonate (MJ) when the *Eleutherococcus senticosus* embryo was cultured in Liquid MS with suspension. However, the fresh weight, dry weight and growth ratio of embryos was strongly inhibited by increasing MJ concentrations. The highest total eleutheroside (7.3 fold increment) and chlorogenic acid (3.9 fold increment) yield was obtained with 200 M methyl jasmonate treatment (Shohael et al., 2007).

The involvement of amino acids in the biosynthesis of hyperforin and adhyperforin was reported in *H. perforatum* shoot cultures. Valine and isoleucine, upon administration to the shoot cultures, were incorporated into acyl side chain of hyperforin and adhyperforin, respectively. Feeding the shoot cultures with unlabelled l-isoleucine at a concentration of 2 mM induced a 3-7-fold increase in the production of adhyperforin. The addition of 3 mM threonine, a precursor of isoleucine, stimulated a

Table 1. *In vitro* secondary metabolites from plant cell, tissue and organs cultures.

Plant name	Active ingredient	Culture medium and plant growth regulator(s)	Culture type	Reference (s)
<i>Aconitum heterophyllum</i>	Aconites	MS + 2,4-D + Kin	Hairy root	Giri et al., 1997
<i>Adhatoda vasica</i>	Vasine	MS + BAP + IAA	Shoot culture	Shalaka and Sandhya, 2009
<i>Agastache rugosa</i>	Rosmarinic acid	MS + 2,4-D + Kin + 3% sucrose	Hairy root	Lee et al., 2007
<i>Agave amaniensis</i>	Saponins	MS + Kinetin	Callus	Andrijany et al., 1999
<i>Ailanthus altissima</i>	Alkaloids	MS + 2,4-D + Kinetin	Suspension	Anderson et al., 1987
<i>Ajuga reptans</i>	Phytoecdysteroids		Hairy root	Matsumoto and Tanaka, 1991
<i>Allium sativum</i>	Allin	MS + IAA + Kinetin	Callus	Malpathak and David, 1986
<i>Aloe saponaria</i>	Glucosides	MS + 2,4-D + Kinetin	Suspension	Yagi et al., 1983
<i>Ambrosia tenuifolia</i>	Altamisine	MS + Kinetin	Callus	Goleniowski and Trippi, 1999
<i>Ammi majus</i>	Umbelliferone	MS + BAP	Shootlet	Krolicka et al., 2006
<i>Ammi visnaga</i>	Furanocoumarin	MS + IAA + GA3	Suspension	Kaul and Staba, 1967
<i>Amsonia elliptica</i>	Indole alkaloids		Hairy root	Sauerwein et al., 1991
<i>Anchusa officinalis</i>	Rosmarinic acid	B ₅ + 2,4-D	Suspension	De-Eknamkul and Ellis, 1985
<i>Angelica gigas</i>	Deoursin	MS (Liq.) + 2,4-D + GA3	Hairy root	Xu et al., 2008
<i>Anisodus luridus</i>	Tropane alkaloids	MS + 2,4-D + BA	Hairy root	Jobanovic et al., 1991
<i>Ammi majus</i>	Triterpenoid	MS + 2,4-D + BA	Suspension	Staniszewska et al., 2003
<i>Arachys hypogaea</i>	Resveratol	G5 + 2,4-D + Kin.	Hairy root	Kim et al., 2008
<i>Armoracia laphthifolia</i>	Fisicoccin	MS + IAA	Hairy root	Babakov et al., 1995
<i>Artemisia absinthum</i>	Essential oil	MS + NAA + BAP	Hairy root	Nin et al., 1997
<i>Artemisia annua</i>	Artemisinin	MS + IAA + Kinetin	Hairy root	Rao et al., 1998
<i>Artemisia annua</i>	Artemisinin	MS + NAA + Kinetin	Callus	Baldi and Dixit, 2008
<i>Aspidosperma ramiflorum</i>	Ramiflorin	MS + 2,4-D + BAP	Callus	Olivira et al., 2001
<i>Aspidosperma ramiflorum</i>	Ramiflorin alkaloid	MS + 2,4-D + BAP + 30 g/l Sucrose	Callus	Olivira et al., 2001
<i>Astragalus mongholicus</i>	Cycloartane saponin	MS + 2,4-D + Kin	Hairy root	Ionkova et al., 1997
<i>Astragalus mongholicus</i>	Cycloartane	MS + IAA + NAA	Hairy root	Ionkova et al., 1997
<i>Azadirachta indica</i>	Azadirachtin	MS + 2,4-D	Suspension	Sujanya et al., 2008
<i>Azadirachta indica</i>	Azadirachtin	MS + 2,4-D + Cyanobacterial elicitor	Suspension	Poornasri Devi et al., 2008
<i>Beeta vulgaris</i>	Betalain pigments	MS + IAA	Hairy root	Taya et al., 1992
<i>Brucea javanica</i>	Alkaloids	MS + 2,4-D + Kinetin	Suspension	Lie et al., 1990
<i>Brucea javanica</i>	Cathin	MS + IAA + GA3	Suspension	Wagiah et al., 2008
<i>Brugmansia candida</i>	Tropane	MS + 2,4-D + IAA	Hairy root	Marconi et al., 2008
<i>Brugmansia candida</i>	Tropane alkaloid	MS+BA+NAA	Hairy root	Giulietti et al., 1993
<i>Bupleurum falcatum</i>	Saikosaponins	B ₅ + IBA	Root	Kusakari et al., 2000
<i>Bupleurum falcatum</i>	Saikosaponins	LS + 2,4-D	Callus	Wang and Huang, 1982
<i>Calystegia sepium</i>	Cuscohygrine	MS + 2,4-D + BA	Hairy root	Jung and Tepfer, 1987
<i>Camellia chinensis</i>	Flavones	MS + 2,4-D + NAA	Callus	Nikolaeva et al., 2009
<i>Camellia sinensis</i>	Theamine	MS + IBA + Kinetin	Suspension	Orihara and Furuya, 1990
<i>Campanula medium</i>	Polyacetylenes	MS+IAA+BA	Hairy root	Tada et al., 1996
<i>Canavalia ensiformis</i>	Canavanine	LS + NAA + Picloram	Callus	Ramirez et al., 1992
<i>Capsicum annum</i>	Capsiacin	MS + 2,4-D+ GA3	Callus	Varindra et al., 2000
<i>Capsicum annum</i>	Capsiacin	MS + 2,4-D + Kin.	Callus	Umamaheswari and Lalitha, 2007
<i>Capsicum annum</i>	Capsaicin	MS + 2,4-D + Kinetin	Suspension	Johnson et al., 1990
<i>Cassia acutifolia</i>	Antraquinones	MS + 2,4-D + kinetin	Suspension	Nazif et al., 2000

Table 1. Contd.

<i>Cassia obtusifolia</i>	Anthraquinone	MS + TDZ + IAA	Hairy root	Ko et al., 1995
<i>Cassia senna</i>	Sennosides	MS + NAA + Kin	Callus	Shrivastava et al., 2006
<i>Catharanthus roseus</i>	Indole alkaloids	MS + IAA	Suspension	Moreno et al., 1993
		MS + NAA + Kinetin	Suspension	Zhao et al., 2001
<i>Catharanthus roseus</i>	Vincristine	MS + 2,4-D + GA3	Suspension	Lee-Parsone and Rogce, 2006
<i>Catharanthus roseus</i>	Indole alkaloid	MS + 2,4-D + GA3 + Vanadium	Suspension	Tallevi and Dicosmo, 1988
<i>Catharanthus roseus</i>	Catharathine	MS + 2,4-D + UV-B radiation	Suspension	Ramani and Jayabaskaran, 2008
<i>Catharanthus trichophyllus</i>	Indole alkaloids	MS + IAA + GA3	Hairy root	Davioud et al., 1989
<i>Cayratia trifoliata</i>	Stilbenes	MS + IAA + GA3	Suspension	Roat and Ramawat, 2009
<i>Centella asiatica</i>	Asiaticoside	MS + 2,4-D	Hairy root	Kim et al., 2007
<i>Centella asiatica</i>	Asiaticoside	MS + 2,4-D + Kin	Callus	Kiong et al., 2005
<i>Centella asiatica</i>	Asiaticoside	MS + BAP + IAA	Shoot	Kim et al., 2004
<i>Centella asiatica</i>	Asiaticoside	MS + 2,4-D	Hairy root	Paek et al., 1996
<i>Centranthes ruber</i>	Valepotriates	MS + IAA + Kin	Hairy root	Granicher et al., 1995
<i>Cephaelis ipecacuanha</i>	Alkaloids	MS + IAA	Root	Teshima et al., 1988
<i>Chaenatis douglasei</i>	Thiarbrins	MS + NAA	Hairy root	Constabel and Towers, 1988
<i>Chrysanthemum cinerariaefolium</i>	Pyrethrins	MS + 2,4-D + Kinetin	Callus	Rajasekaran et al., 1991
		MS + Kinetin	Suspension	Kuch et al., 1985
<i>Cinchona ledgeriana</i>	Quinine	MS + 2,4-D	Hairy root	Hamill et al., 1989
		B ₅ + 2,4-D	Suspension	Schripsema et al., 1999
		B ₅ + 2,4-D + Kinetin	Suspension	Wijnsma et al., 1985
<i>Cinchona succirubra</i>	Anthraquinone	MS + IAA + GA3	Suspension	Khoury et al., 1986
<i>Citrus</i> sp.	Limonin	MS + 2,4-D + Kinetin	Callus	Barthe et al., 1987
<i>Coffea arabica</i>	Caffeine	MS + 2,4-D + Kinetin	Callus	Waller et al., 1983
<i>Coleus forskohlii</i>	Forskolin	MS + IAA + Kin	Hairy root	Sasaki et al., 1998
<i>Corydalis ambigua</i>	Corydaline	MS + IAA + 3% sucrose	Embryo	Hiraoka et al., 2004
<i>Corydalis cava</i>	Corydaline	MS + IAA + GA3	Shoot	Rueffer et al., 1994
<i>Corydalis ophiocarpa</i>	Alkaloids	MS + 2,4-D + Kinetin	Callus	Iwasa and Takao, 1982
<i>Corydalis terminalis</i>	Corydalin	MS + 2,4-D + BAP	Callus	Taha et al., 2008
<i>Coscinium fenestratum</i>	Berberin	MS + 2,4-D + BAP	Callus	Khan et al., 2008
<i>Coscinium fenestratum</i>	Berberine	MS + IAA + BAP	Callus	Nair et al., 1992
<i>Coscinium fenestratum</i>	Berberine	MS + 2,4-D + GA3	Suspension	Narasimhan and Nair, 2004
<i>Crataegus sinaica</i>	Flavonoid	MS + 2,4-D + NAA + BAP	Callus	Maharik et al., 2009
<i>Croton sublyratus</i>	Plaunotol	MS+NAA+BA	Callus	Morimoto and Murai, 1989
<i>Cruciata glabra</i>	Anthraquinones	LS + NAA + Kinetin	Suspension	Dornenburg and Knorr, 1996
<i>Cryptolepis buchanani</i>	Cryptosin	B ₅ + 2,4-D + Kinetin	Callus	Venkateswara et al., 1987
<i>Cymbopogon citratus</i>	Essential oil	MS + IAA + GA3	Shoot	Quiala et al., 2006
<i>Datura stramonium</i>	Hyocyamine	MS + IAA	Hairy root	Hilton and Rhodes, 1993
<i>Digitalis purpurea</i>	Cardenolides	MS+BA	Suspension	Hagimori et al., 1982

Table 1. Contd.

<i>Digitalis purpurea</i>	Cardioactive glycosides	MS + 2,4-D + BA	Hairy root	Saito et al., 1990
<i>Diocorea doryophora</i>	Diogenin	MS + 2,4-D + BA	Suspension	Huang et al., 1993
<i>Dioscorea deltoidea</i>	Diosgenin	MS + 2,4-D	Suspension	Heble and Staba, 1980
<i>Drosera rotundifolia</i>	7-Methyljuglone	MS + BAP + NAA	Shoot culture	Hohtola et al., 2005
<i>Duboisia leichhardtii</i>	Alkaloids	LS+NAA+BA	Callus	Yamada and Endo, 1984
<i>Duboisia leichhardtii</i>	Scopalamine	MS + 2,4-D + BA	Hairy root	Muranaka et al., 1992
<i>Duboisia myoporoides</i>	Scopalamine	MS + IAA	Hairy root	Deno et al., 1987
<i>Echinacea purpurea</i>	Alkamides	MS + 2,4-D	Hairy root	Trysteen et al., 1991
<i>Eleutherococcus senticosus</i>	Eleuthrosides	MS + 2,4-D	Suspension	Shohael et al., 2007
<i>Ephedra</i> sp.	L-Ephedrine	MS + Kinetin + 2,4-D	Suspension	O'Dowd et al., 1993
<i>Eriobotrya japonica</i>	Triterpenes	LS+NAA+BA	Callus	Taniguchi et al., 2002
<i>Eucalyptus tereticornis</i>	Sterols and phenolic compounds	MS + 2,4-D	Callus	Venkateswara et al., 1986
<i>Fabiana imbricata</i>	Rutin	MS + NAA + 2,4-D	Callus and Suspension	Schmeda-Hirschmann et al., 2004
<i>Fagopyrum esculentum</i>	Flavonol	MS + IAA + GA3	Hairy root	Trotin et al., 1993
<i>Fagopyrum esculentum</i>	Rutin	MS + NAA	Hairy root	Lee et al., 2007
<i>Frangula alnus</i>	Anthraquinones	WPM + IAA + BAP	Callus	Kovacevic and Grabisic, 2005
<i>Fritillaria unibracteata</i>	Alkaloids	MS + 2,4-D + Kin	Multiple shoot	Gao et al., 2004
<i>Fumaria capreolata</i>	Alkaloids	LS + IAA	Suspension	Tanahashi and Zenk, 1985
<i>Gentiana macrophylla</i>	Glucoside	MS + IAA + Kin	Hairy root	Tiwari et al., 2007
<i>Gentiana</i> sp.	Glucosides	B ₅ + Kinetin	Callus	Skrzypczak et al., 1993
<i>Gentianella austriaca</i>	Xanthone	MS + BAP	Multiple shoot	Vinterhalter et al., 2008
<i>Geranium thunbergii</i>	Tannin	MS + 2,4-D + BAP	Hairy root	Ishimaru and Shimomura, 1991
<i>Ginkgo biloba</i>	Ginkoside-A	MS + NAA + Kinetin	Suspension	Carrier et al., 1991
<i>Glehnia littoralis</i>	Furanocoumarin	LS + 2,4-D + Kinetin	Suspension	Kitamura et al., 1998
<i>Glycyrrhiza echinata</i>	Flavonoids	MS + IAA + Kinetin	Callus	Ayabe et al., 1986
<i>Glycyrrhiza glabra</i>	Triterpenes	MS + IAA + Kinetin + 2,4-D	Callus	Ayabe et al., 1990
<i>Glycyrrhiza glabra</i>	Glycyrrhizin	MS + 2,4-D + GA3	Hairy root	Mehrotra et al., 2008
<i>Glycyrrhiza glabra</i>	Flavonoid	MS + IAA	Hairy root	Asada et al., 1998
<i>Gymnema sylvestre</i>	Gymnemic acid	MS + 2,4-D + IAA	Callus	Gopi and Vatsala, 2006
<i>Gymnema sylvestre</i>	Gymnemic acid	MS+IAA+BA	Callus	Devi et al., 2006
<i>Gynostemma pentaphyllum</i>	Saponin	MS + 2,4-D + BAP	Hairy root	Fei et al., 1993
<i>Gypsophila paniculata</i>	Saponin	MS + IAA + TDZ	Root suspension	Fulcheri et al., 1998
<i>Hemidesmus indicus</i>	Lupeol, Rutin	MS + BAP + NAA	Shoot culture	Misra et al., 2005
<i>Hyocyamus niger</i>	Tropane alkaloids	MS + 2,4-D + BA	Callus	Yamada and Hashimoto, 1982
<i>Hyocyamus niger</i>	Tropane alkaloid	MS + IAA + Kinetin	Hairy root	Jaziri et al., 1988
<i>Hyoscyamus albus</i>	Phytolexins	MS + NAA + GA3	Hairy root	Kuroyanagi et al., 1998
<i>Hyoscyamus muticus</i>	Hyoscyamine	MS + 2,4-D	Hairy root	Halperin and Flores, 1997
<i>Hypericum perforatum</i>	Hypericin	Liquid MS + NAA + GA3	Suspension	Hohtola et al., 2005
<i>Hypericum perforatum</i>	Hypericins	MS+BA+IAA	Multiple shoot	Kornfeld et al., 2007
<i>Hypericum perforatum</i>	Hypericin	MS+BA+TDZ	Multiple shoot	Santarem and Astarita, 2003
<i>Hypericum perforatum</i>	Hyperforin	MS + 2,4-D + Leusine	Multiple shoot	Karppinen et al., 2007
<i>Hyssopus officinalis</i>	Triterpenes	G5 + 2,4-D + IAA	Suspension	Skrzypek and Wysokinsku, 2003
<i>Hyssopus officinalis</i>	Sterols	MS + 2,4-D + NAA	Suspension	Skrzypek and Wysokinsku, 2003

Table 1. Contd.

<i>Ipomoea cairica</i>	Lignan	MS + IAA + Kin	Callus	Paska et al., 1999
<i>Isoplexis isabelliana</i>	Anthraquinone	MS + 2,4-D + Kinetin	Suspension	Arrebola et al., 1999
<i>Lactuca virosa</i>	Sesquiterpene lactones	MS + 2,4-D	Hairy root	Kisiel et al., 1995
<i>Leontopodium alpinum</i>	Essential oil	MS+IAA+BA	Hairy root	Hook, 1994
<i>Linum flavum</i>	5-Methoxyphyllotaxin	MS salts + B ₅ Vitamins	Suspension	Uden et al., 1990
<i>Linum flavum</i>	Lignan	MS + IAA + GA3	Hairy root	Oostdam et al., 1993
<i>Lithospermum erythrorhizon</i>	Shikonin derivatives	LS + IAA + Kinetin	Suspension	Fujita et al., 1981 ; Fukui et al., 1990
<i>Lithospermum erythrorhizon</i>	Shikonin	MS + 2,4-D + Kinetin	Hairy root	Fukui et al., 1998
<i>Lobelia cardinalis</i>	Polyacetylene glucosides	MS + 2,4-D	Hairy root	Yamanaka et al., 1996
<i>Lycium chinense</i>	Cerebroside	MS + 2,4-D, Kinetin	Suspension	Jang et al., 1998
<i>Mentha arvensis</i>	Terpenoid	MS+BA+NAA	Shoot	Phatak and Heble, 2002
<i>Momordica charantia</i>	Flavonoid	MS + BAP + NAA	Callus	Agarwal and Kamal, 2007
<i>Morinda citrifolia</i>	Anthraquinones	B ₅ + NAA	Suspension	Zenk et al., 1975
		B ₅ + NAA + Kinetin	Suspension	Assetti et al., 1995
<i>Mucuna pruriens</i>	L-Dopa	MS + IAA	Suspension	Wichres et al., 1993
		MS + 2,4-D		Brain, 1976
			Callus	
<i>Myristica fragrans</i>	Myristin	MS + NAA + TDZ	Shoot	Indira Iyer et al., 2009
<i>Nandina domestica</i>	Alkaloids	MS + 2,4-D + Kinetin	Callus	Ikuta and Itokawa, 1988
<i>Nicotiana glauca</i>	Anatabine	MS + IAA	Hairy root	Parr and Hamill, 1987
<i>Nicotiana glauca</i>	Alkaloids	LS + 2,4-D + Kinetin	Callus	Tabata ad Hiraoka, 1976
<i>Nicotiana glauca</i>	Nicotine	MS + NAA + Kinetin	Suspension	Mantell et al., 1983
<i>Ophiorrhiza rugosa</i> var. <i>decumbens</i>	Camptothecin	MS + BA + Kin.	Shoot	Vineesh et al., 2007
<i>Panax ginseng</i>	Saponin and spogenins	MS + 2,4-D	Callus	Furuya et al., 1973 ; Asaka et al., 1993
<i>Panax ginseng</i>	Glycoside	MS + NAA + Kin.	Hairy root	Jeong and Park, 2007
<i>Panax notoginseng</i>	Gensenosides	MS + 2,4-D + Kinetin	Suspension	Zhong and Zhu, 1995
<i>Papaver bracteatum</i>	Thebaine	MS + Kinetin + 2,4-D	Callus	Day et al., 1986
<i>Papaver somniferum</i>	Alkaloids	MS + Kinein	Callus	Furuya et al., 1972
	Morphine and codeine			Siah and Doran, 1991
		MS + 2,4-D + Kinetin	Suspension	
<i>Papaver somniferum</i>	Codeine	LS+BA+NAA	Hairy root	Williams and Ellis, 1992
<i>Peganum harmala</i>	Alkaloids	MS + 2,4-D	Suspension	Sasse et al., 1982
<i>Perezia cuernavacana</i>	Sesquiterpene quinone	MS+IAA+BA	Hairy root	Arellano et al., 1996
<i>Phiorrhiza pumila</i>	Alkaloids	LS + 2,4-D + NAA	Callus	Kitajima et al., 1998
<i>Phytolacca americana</i>	Betacyanin	MS + 2,4-D	Suspension	Sakuta et al., 1987
<i>Picrasma quassioides</i>	Quassin	B ₅ + 2,4-D + Kinetin	Suspension	Scragg and Allan, 1986
<i>Pimpinella anisum</i>	Essential oil	MS + IAA + BAP	Hairy root	Santos et al., 1998
<i>Piper solmsianum</i>	Piperine	MS + 2,4-D + BA	Suspension	Balbuena et al., 2009
<i>Plantago media</i>	Verbascoside	B ₅ + IAA + Kin.	Callus	Kunvari et al., 1999
<i>Platycodon grandiflorum</i>	Polyacetylene	MS + 2,4-D	Hairy root	Tada et al., 1995
<i>Pluchea lanceolata</i>	Quercetin	MS + NAA + BAP	Callus	Arya et al., 2008

Table 1. Contd.

<i>Plumbago rosea</i>	Plumbagin	MS + CaCl ₂	Callus	Komaraiah et al., 2003
<i>Plumbago zeylanica</i>	Plumbagin	MS + BAP + IBA	Hairy root	Verma et al., 2002
<i>Podophyllum hexandrum</i>	Podophyllotaxin	B ₅ + NAA	Suspension	Uden et al., 1989
<i>Podophyllum hexandrum</i>	Podophyllotaxin	MS + BAP + GA32	Shoot	Li et al., 2009
<i>Polygala amarella</i>	Saponin	MS + IAA	Callus	Desbene eet al., 1999
<i>Polygonum hydropiper</i>	Flavonoids	MS + 2,4-D + Kinetin	Suspension	Nakao et al., 1999
<i>Portulaca grandiflora</i>	Betacyanin	MS + 2,4-D + Kinetin	Callus	Schroder and Bohm, 1984
<i>Primula veris</i>	Saponins	MS + BAP + GA3	Shoot	Okrslar et al., 2007
<i>Psoralea cordifolia</i>	Isoflavones	MS + TDZ + BAP	Multiple shoot	Shinde et al., 2009
<i>Psoralea corylifolia</i>	Isoflavones	MS + TDZ + BAP	Multiple shoot	Shinde et al., 2009
<i>Ptelea trifoliata</i>	Alkaloids	MS + 2,4-D +Kinetin	Callus	Petit-Paly et al., 1987
<i>Rauvolfia sellowii</i>	Alkaloids	B ₅ + 2,4-D + Kinetin	Callus	Rech et al., 1998
<i>Rauvolfia serpentina</i>	Reserpine	LS + NAA +BA LS	Suspension Callus	Yamamoto and Yamada, 1996 Gerasimenko et al., 2001
<i>Rauvolfia serpentina</i>	Serpentine	MS + BAP + IAA	Callus	Salma et al., 2008
<i>Rauvolfia serpentina</i>	Reserpine	MS + IAA + Cu2+	Callus	Nurchgani et al., 2008
<i>Rauvolfia tetraphylla</i>	Reserpine	MS + 2,4-D + Tryptophan	Callus	Anitha and Kumari, 2006
<i>Rhamnus catharticus</i>	Anthraquinones	WPM + Kin + 2,4-D	Callus	Kovacevic and Grabisic, 2005
<i>Rheum ribes</i>	Catechin	MS+IBA+BA	Callus	Farzami and Ghorbant, 2005
<i>Rhodiola rosea</i>	Rosarin	MS + NAA + IAA	Callus culture	Hohtola et al., 2005
<i>Rhus javanica</i>	Gallotannins	LS + IAA +Kinetin	Root	Tanoguchi et al., 2000
<i>Rubia akane</i>	Anthraquinone	B ₅ + NAA + Kin	Hairy root	Park and Lee, 2009
<i>Rubia akane</i>	Anthraquinone	MS + 2,4-D + Chitosan	Suspension	Jin et al., 1999
<i>Rubia tinctorum</i>	Anthraquinone	MS + 2,4-D	Hairy root	Sato et al., 1991
<i>Ruta sp.</i>	Alkaloids and coumarins	MS + 2,4-D + Kinetin	Callus	Baumert et al., 1992
<i>Salvia miltiorrhiza</i>	Rosmarinic acid	MS + 2,4-D + BA MS + 2,4-D + Kinetin	Callus	Morimoto et al., 1994 Miyasaka et al., 1989
<i>Salvia officinalis</i>	Cryptotanshinone Flavonoid	LMS + IAA + BAP	Suspension Multiple shoot	Grzegorzcyk and Wysokinska, 2008
<i>Salvia officinalis</i>	Terpenoids	MS + 2,4-D + BA	Callus	Santos-Gome et al., 2002
<i>Saponaria officinalis</i>	Saponin	MS + IAA + TDZ	suspension	Fulcheri et al., 1998
<i>Saprosma fragrans</i>	Anthraquinone	MS + 2,4-D + NAA	Callus	Singh et al., 2006
<i>Scoparia dulcis</i>	Scopadulic acid	LMS + Kin + Phenyl urea	Callus	Hayashi et al., 1998
<i>Scopolia parviflora</i>	Alkaloids	LS + 2,4-D + IAA	Callus	Tabata et al., 1972
<i>Scutellaria baicalensis</i>	Flavonoids	MS + IAA	Hairy root	Zhou et al., 1997
<i>Scutellaria columnae</i>	Phenolics	MS + 2,4-D + Kinetin	Callus	Stojakwska and Kisiel, 1999
<i>Serratula tinctoria</i>	Ecdysteroid	MS + 2,4-D + BA	Hairy root	Delbeque et al., 1995
<i>Sesamum indicum</i>	Napthaquinone	MS + NAA + Kinetin	Hairy root	Ogasawara et al., 1993
<i>Silybium marianum</i>	Silymarin	MS + IAA + GA3	Hairy root	Rahnama et al., 2008
<i>Silybium mariyanm</i>	Flavonolignan	LS + TDZ	Root	Alikaridis et al., 2000
<i>Silybum marianum</i>	Silymarin	MS + IAA + Kin	Hairy root	Rahnama et al., 2008
<i>Silybum marianum</i>	Silymarin	MS+IAA+BA	Callus	Tumova et al., 2006
<i>Simmondsia chinensis</i>	Fixed oil	MS + TDZ + GA3	Callus	Aftab et al., 2008
<i>Simmondsia chinensis</i>	Fixed Oil	MS + IAA + 2iP	Callus	Aftab et al., 2008
<i>Solanum aculeatissi</i>	Steroidal saponin	MS + 2,4-D	Hairy root	Ikenaga et al., 1995
<i>Solanum chrysotrichum</i>	Saponin	MS + 2,4-D + Kinetin	Suspension	Villarreal et al., 1997
<i>Solanum laciniatum</i>	Solasodine	MS + 2,4-D + Kinetin	Suspension	Handler and Dodds, 1983

Table 1. Contd.

<i>Solanum paludosum</i>	Solamargine	MS + BA + Kinetin	Suspension	Badaouti et al., 1996
<i>Stevia rebaudiana</i>	Stevioside	MS+BA+NAA	Callus	Dheeranapattana et al., 2008
<i>Swertia japonica</i>	Amarogenetin	MS + IAA	Hairy root	Ishimaru et al., 1990
<i>Tabernaemontana divariacata</i>	Alkaloids	MS+NAA+BA	Suspension	Sierra et al., 1992
<i>Tagetes patula</i>	Thiophenes	MS + IAA + Kinetin	Hairy root	Arroo et al., 1995
<i>Tanacetum parthenium</i>	Sesquiterpene	MS + 2,4-D + Kinetin	Hairy root	Kisiel and Stojakowska, 1997
<i>Taxus baccata</i>	Taxol baccatin III	B ₅ + 2,4-D + Kinetin + GA ₃	Suspension	Cusido et al., 1999
<i>Taxus</i> spp.	Taxol	B ₅ + 2,4-D + BA	Suspension	Wu et al., 1992
<i>Thalictrum minus</i>	Berberin	LS + NAA + 2,4-D + BA	Suspension	Kobayashi et al., 1987
		LS+NAA+BA	Suspension	Nakagawa et al., 1986
<i>Tinospora cordifolia</i>	Berberin	MS + IAA + GA ₃	Suspension	Rama Rao et al., 2008
<i>Torreya nucifera</i>	Diterpenoids	MS + 2,5-D	Suspension	Orihara et al., 2002
<i>Trichosanthes kirilowii</i>	Protein	MS + IAA	Hairy root	Savary and Flores, 1994
<i>Trigonella foenu-graecum</i>	Saponins	MS + 2,4-D + Kinetin	Suspension	Brain and Williams, 1983
<i>Vaccinium myrtillus</i>	Flavonoids	MS + BAP + NAA	Callus culture	Hohtola et al., 2005
<i>Vinca major</i>	Vincamine	MS + BAP	Hairy root	Tanaka et al., 2004
<i>Vitis vinifera</i>	Anthocyanin	MS + BAP + NAA	Suspension	Qu et al., 2006
<i>Vitis vinifera</i>	Resveratrol	MS+IAA+GA ₃ +UV	Callus	Kin and Kunter, 2009
<i>Withania somnifera</i>	Withaferin A	MS+BA	Shoot	Ray and Jha, 2001
<i>Withania somnifera</i>	Withaferin	MS + IAA + Kintin	Hairy root	Banerjee et al., 1994
<i>Withania somnifera</i>	Withanoloid A	MS + IAA + Kin	Hairy root	Murthy et al., 2008
<i>Withania somnifera</i>	Steroidal lactone	MS + 2,4-D + BA	Callus	Mirjalili et al., 2009
<i>Zataria multiflora</i>	Rosmarinic acid	MS + IAA + Kin	Callus	Francoise et al., 2007

2-fold increase in the accumulation of adhyperforin (Karppinen et al., 2007). Production of triterpenes in leaf derived callus and cell suspension cultures of *Centella asiatica* was enhanced by the feeding of aminoacids. In the callus culture manifold increase of asiaticoside accumulation was reported with the addition of leucien (Kiong et al., 2005).

The effect of coniferyl alcohol as a precursor of flavonolignan biosynthesis on silymarin components production in *Silybum marianum* suspension culture was reported (Tumova et al., 2006). Coniferyl alcohol showed the changes in silymarin complex production. A significant increase of silydianin was observed only after 72 h of the application of 46 M coniferyl alcohol. The same precursor – coniferyl alcohol in the form of complex with -cyclodextrin was used as precursor for podophyllotoxin accumulation in *Podophyllum hexandrum* cell suspension cultures.

Elicitation of *in vitro* products

Pharmaceutically significant secondary metabolites or phytopharmaceuticals include alkaloids, glycosides, flavonoids, volatile oils, tannins, resins etc. Currently, most of these secondary metabolites are isolated from wild or cultivated plants because their chemical syn-

thesis is either extremely difficult or economically infeasible. Biotechnological production in plant cell cultures is an attractive alternative, but to date this has had only limited commercial success because of a lack of understanding of how these metabolites are synthesized. Plants and/or plant cells *in vitro*, show physiological and morphological responses to microbial, physical or chemical factors which are known as 'elicitors'. Elicitation is a process of inducing or enhancing synthesis of secondary metabolites by the plants to ensure their survival, persistence and competitiveness (Namdeo, 2007).

The production of secondary metabolites in callus, cell suspension and hairy roots of *Ammi majus* L. is by exposing them to elicitors: benzo(1,2,3)-thiadiazole-7-carbothionic acid S-methyl ester and autoclaved lysate of cell suspension of bacteria—*Enterobacter sakazaki* (Staniszewska et al., 2003). GC and GC–MS analysis of chloroform and methanol extracts indicated a higher accumulation of umbelliferone in the elicited tissues than in the control ones. Plants generally produce secondary metabolites in nature as a defense mechanism against pathogenic and insect attack. The study was applied in several abiotic elicitors to enhance growth and ginseng saponin biosynthesis in the hairy roots of *Panax ginseng* (Jeong and Park, 2007). Generally, elicitor treatments were found to inhibit the growth of the hairy roots, al-

though simultaneously enhancing ginseng saponin biosynthesis. Tannic acid profoundly inhibited the hairy root growth during growth period. Also, ginseng saponin content was not significantly different from that of the control. The addition of selenium at inoculum time did not significantly affect ginseng saponin biosynthesis. However, when 0.5 mM selenium was added as an elicitor after 21 d of culture, ginseng saponin content and productivity increased to about 1.31 and 1.33 times control levels, respectively. Also, the addition of 20 M NiSO₄ resulted in an increase in ginseng saponin content and productivity, to about 1.20 and 1.23 times control levels, respectively, and also did not inhibit the growth of the roots. Sodium chloride treatment inhibited hairy root growth, except at a concentration of 0.3% (w/v). Increases in the amounts of synthesized ginseng saponin were observed at all concentrations of added sodium chloride. At 0.1% (w/v) sodium chloride, ginseng saponin content and productivity were increased to approximately 1.15 and 1.13 times control values, respectively. These results suggest that processing time for the generation of ginseng saponin in a hairy root culture can be reduced via the application of an elicitor.

Chitosan was the biotic elicitor polysaccharide and it is eliciting the manifold increase of anthraquinone production in *Rubia akane* cell culture (Jin et al., 1999).

Metabolic engineering and production of secondary metabolites

Metabolic engineering involves the targeted and purposeful alteration of metabolic pathways found in an organism to achieve better understanding and use of cellular pathways for chemical transformation, energy transduction, and supramolecular assembly (Lessard, 1996). This technique applied to plants will permit endogenous biochemical pathways to be manipulated, resulting in the generation of transgenic crops in which the range, scope, or nature of a plant's existing natural products are modified to provide beneficial commercial, agronomic and/or post-harvest processing characteristics (Kinney, 1998). Over the last decades, plant cell cultures have been intensively investigated as a possible tool for the production of commercial plant secondary metabolites, including fine chemicals such as pharmaceuticals, agrochemicals, flavors, insecticides, fragrances and cosmetics (Whitmer et al., 2002). In spite of the efforts in the field of *in vitro* production of phytochemicals, few industrial processes have been developed, involving only a limited number of secondary products, such as shikonin, berberine, ginsenosides and paclitaxel (Ramachandra and Ravishankar, 2002). As in many cases production is too low for commercialization, metabolic engineering can provide various strategies to improve productivity, such as: increasing the number of producing cells, increasing the carbon flux through a

biosynthetic pathway by overexpression of genes codifying for rate-limiting enzymes or blocking the mechanism of feedback inhibition and competitive pathways and decreasing catabolism.

Many of the isolated pure compounds with biological activity are alkaloids, a diverse group of nitrogen-containing chemical ring structure compounds, with alkali-like chemical reactivity and pharmacological activity. Although the pharmacological effects of alkaloids have been studied, the biosynthetic pathways of these compounds are still obscure. Among the most famous are the tropane alkaloids, such as (-)-hyoscyamine, its racemate atropine, and scopolamine (hyoscyne), which have an 8-azabicyclo[3.2.1]octane esterified nucleus. These alkaloids are commonly found in plants of different families: Solanaceae, Erythroxylaceae, Convolvulaceae, Proteaceae, Euphorbiaceae, Rhizophoraceae and Cruciferae (Griffing and Lin, 2000). Related to the tropane alkaloids, a new group of nortropane alkaloids, the calystegines, was discovered only 15 years ago. Calystegines bear three to five hydroxyl groups in various positions, making them water-soluble, and they share metabolic steps and enzymes of the formation of tropane alkaloids.

Several genes in the biosynthetic pathways for scopolamine, nicotine, and berberine have been cloned, making the metabolic engineering of these alkaloids possible. Expression of two branching-point enzymes was engineered: putrescine *N*-methyltransferase (PMT) in transgenic plants of *Atropa belladonna* and *Nicotiana sylvestris* and (*S*)-scoulerine 9-*O*-methyltransferase (SMT) in cultured cells of *Coptis japonica* and *Eschscholzia californica*. Overexpression of PMT increased the nicotine content in *N. sylvestris*, whereas suppression of endogenous PMT activity severely decreased the nicotine content and induced abnormal morphologies. Ectopic expression of SMT caused the accumulation of benzyloquinoline alkaloids in *E. californica* (Sato et al., 2001).

Hairy root cultures as a source of secondary metabolites

The hairy root system based on inoculation with *Agrobacterium rhizogenes* has become popular in the two last decades as a method of producing secondary metabolites synthesized in plant roots (Palazon et al., 1997). Unorganized plant tissue cultures are frequently unable to produce secondary metabolites at the same levels as the intact plant. This is also the case of scopolamine production in undifferentiated *in vitro* cultures of Solanaceae, probably due to the specific location of some of the key enzymes involved in this biosynthetic pathway (Palazon et al., 2006). Suzuki et al. (1997) have demonstrated that the expression of the *pmt* gene was pericycle-specific, and it has also been shown

that H6H is localized in the root pericycle (Kanegae et al., 1994). In addition, Nakajima and Hashimoto (1999) have observed that TR proteins accumulate in the lateral roots of *Hyoscyamus niger*. Another possible reason for the low production of scopolamine in undifferentiated *in vitro* cultures could be that the auxin added to the callus and cell culture media for normal growth inhibit the activity of some of the key enzymes involved in scopolamine biosynthesis, such as PMT (Rothe et al., 2003).

The hairy root phenotype is characterized by fast hormone-independent growth, lack of geotropism, lateral branching and genetic stability. The secondary metabolites produced by hairy roots arising from the infection of plant material by *A. rhizogenes* are the same as those usually synthesized in intact parent roots, with similar or higher yields (Sevon and Oksman-Caldentey, 2002). This feature, together with genetic stability and generally rapid growth in simple media lacking phytohormones, makes them especially suitable for biochemical studies not easily undertaken with root cultures of an intact plant. During the infection process *A. rhizogenes* transfers a part of the DNA (transferred DNA, T-DNA) located in the root-inducing plasmid Ri to plant cells and the genes contained in this segment are expressed in the same way as the endogenous genes of the plant cells. Some *A. rhizogenes*, such as strain A4, have the T-DNA divided in two sections: the TR-DNA and TL-DNA, each of which can be incorporated separately into the plant genome. Two sets of pRi genes are involved in the root induction process: the *aux* genes located in the TR region of the pRi T-DNA and the *rol* (root loci) genes of the TL region (Jouanin, 1989). The *ags* genes responsible for opine biosynthesis in the transformed tissues are also located in the TR region (Binns and Tomashow, 1988). Opines are synthesized by plant transformed cells and are only used by *Agrobacterium* as a source of nitrogen and carbon. Due to the similarities of the *A. rhizogenes* and *A. tumefaciens* infection processes, and because both microorganisms are very closely related, it has been suggested that the most important *A. rhizogenes* oncogenes encode proteins involved in the regulation of plant hormone metabolism. *Aux* genes provide transformed cells with an additional source of auxin (Chriqui et al., 1996), but they do not seem essential for developing hairy root disease. However, *rol* genes have functions that are most likely other than that of producing mere alterations in plant hormone concentrations (Nilsson and Olsson, 1997). Several authors have investigated the effect of TR and TL regions of *A. rhizogenes* on growth and morphology of transformed roots and plants, but until now there have been few studies on the direct effects of oncogenes on secondary metabolism. As has been previously reported, a correlation exists between the expression of the *rolC* gene and tropane alkaloids (Fumanova and Syklovska, 2000), *Catharanthus roseus* alkaloids (Palazon et al., 1998), and ginsenoside production (Bulkagov et al., 1998). No correlation between

rolA and *rolB* expression and secondary metabolism was found in any of these studies. Moyano et al. (1999) showed that the inoculation of leaf sections of tobacco, *Duboisia* hybrid and *Datura metel* plants with the A4 strain of *A. rhizogenes* induced transformed roots with the capacity to produce putrescine-derived alkaloids such as nicotine, hyoscyamine and scopolamine. In general, the obtained hairy roots presented two morphologies: typical hairy roots with a high capacity to produce alkaloids, and callus-like roots with faster growth and lower alkaloid production. The *aux1* gene of *A. rhizogenes* located in the TR-DNA of *A. rhizogenes* was detected in all roots showing callus-like morphology. However, this gene was only detected in 25-60% of the established root cultures showing typical hairy morphology. These results demonstrate a significant role of *aux* genes in the morphology of transformed roots and the importance of typical hairy root morphology in the production of secondary metabolites. The studies with *Panax ginseng* hairy roots also support the effects of the genes located in the TR-DNA on root morphology and secondary metabolism (Mallol et al., 2002). The hairy roots are normally induced on aseptically wounded parts of plants by inoculating them with *A. rhizogenes*. In scopolamine-producing Solanaceae plants, roots usually emerge at the inoculation sites after 1-4 weeks, but in the case of other plant species such as *Taxus* it can be more than 4 months before the roots emerge. Root tips are cultured separately in a hormone-free medium, the most commonly used being MS (Murashige-Skoog, Gamborg's B5 or SH (Schenk and Hildebrandt, 1972)). The next step for establishing hairy root cultures is to select and characterize the root clones according to their capacity for growth and production of the desired compounds. Sometimes these productions were achieved after a laborious process to optimize the growth conditions, such as the selection of the more productive clones, and optimization of the production conditions by testing different ionic concentrations as well as the carbon source and pH of the medium.

Srivastava and Srivastava (2007) have recently summarized the attempts to adapt bioreactor design to hairy root cultures; stirred tank, airlift, bubble columns, connective flow, turbine blade, rotating drum, as well as different gas phase reactors have all been used successfully. In the case of tropane alkaloids, different types of bioreactors are used for scopolamine production. Wilson (1997) describes the only large droplet bioreactor system with a volume of 500 L designed for hairy root cultures of *Datura stramonium*. On a smaller scale, modified airlift and stirred tanks have been used for scopolamine production in hairy root cultures of *D. metel*, connective flow reactors for *H. muticus* (Carvalho and Curtis, 1998) and *Atropa belladonna* (Williams and Doran, 2000) and more recently a bubble column bioreactor has been employed for root cultures of *Scopolia parviflora* (Min et al., 2007). One such advance is the development of dis-

posable wave bioreactor systems, with working principle is based on wave-induced agitation, which significantly reduces stress levels. This type of bioreactor has been successfully used for *H. muticus* and *Panax ginseng* hairy root cultures (Eibl and Eibl, 2004).

Genetic manipulation in hairy root culture for secondary metabolite production

Transformed roots provide a promising alternative for the biotechnological exploitation of plant cells. *A. rhizogenes* mediated transformation of plants may be used in a manner analogous to the well-known procedure employing *A. tumefaciens*. *A. rhizogenes* mediated transformation has also been used to produce transgenic hairy root cultures and plantlets have been regenerated. None of the other T-DNA sequences are required for the transfer with the exception of the border sequences. The rest of the T-DNA can be replaced with the foreign DNA and introduced into cells from which whole plants can be regenerated. These foreign DNA sequences are stably inherited in a Mendelian manner (Zambryski et al., 1989). The *A. rhizogenes* mediated transformation has the advantage of being able to transfer any foreign gene of interest placed in binary vector to the transformed hairy root clone. It is also possible to selectively alter some plant secondary metabolites or to cause them to be secreted by introducing genes encoding enzymes that catalyze certain hydroxylation, methylation, and glycosylation reactions. An example of a gene of interest with regard to secondary metabolism that was introduced into hairy roots is the 6- β -hydroxylase gene of *Hyoscyamus muticus* which was introduced to hyocyanin rich *Atropa belladonna* by a binary vector system using *A. rhizogenes* (Hashimoto et al., 1993). Engineered roots showed an increased amount of enzyme activity and a five-fold higher concentration of scopolamine. Hairy root cultures of *Nicotiana rustica* with ornithin decarboxylase gene from yeast, and *Peganum harmala* with tryptophan decarboxylase gene from *Catharanthus roseus* (Berlin et al., 1993) have been shown to produce increased amounts of the secondary metabolites nicotine and serotonin when expressing transgenes from yeast. Downs et al. (1994) reported transgenic hairy roots in *Brassica napus* containing a glutamine synthase gene from soybean showing a three-fold increase in enzyme activity. This approach may be a reality for the commercial production of pharmaceutically important compounds using transgenic hairy root culture system.

Role of endophytes in *in vitro* production of secondary metabolites

There are three schools of thought on the origins of secondary metabolism in plants (Wink, 2008). There is

the argument that both plants and endophytic microbes co-evolved with pathways to produce these natural products. Another thought is that an ancient horizontal gene transfer took place between plants and microbes. The third suggests that either plants or endophytic fungi produce these secondary metabolites and transfer them to the other symbiont. Biosynthetic pathway studies using radio-labeled precursor amino acids reveal that plants and endophytic fungi have similar, but distinct metabolic pathways for production of secondary metabolites (Jennewein et al., 2001). Evidence to support the independent production of Taxol by endophytic fungi is the isolation of the gene 10-deacetylbaconin-III-10-O-acetyl transferase from the endophytic fungus *Cladosporium cladosporioides* MD213 isolated from *Taxus media* (yew species). This gene is involved in the biosynthetic pathway of Taxol and shares 99% identity with *T. media* (plant) and 97% identity with *T. wallichiana* var. *marirei* (plant). These data lead to the hypothesis that plants and endophytic fungi through mutualistic symbiosis produce similar secondary metabolites. Recently, it was reported that plants other than yew species also have endophytic fungi associated with them that make Taxol. This suggests that plants and fungi are independently capable of producing these important secondary metabolites. The question is whether bioactive phytochemicals of plants are produced by the plant itself or as a consequence of a mutualistic relationship with beneficial organisms in their tissue. The fact that a combination of inducing factors from both plants and endophytic fungi increased the accumulation of secondary metabolites in plants and fungi respectively (Zhang et al., 2009; Li et al., 2009) suggest that the fungal endophyte may play important roles in the biosynthesis of secondary metabolites. Therefore, the symbiotic association and effects of plants and endophytes on each other during the production of other important pharmacological bioactive natural products such as camptothecin, vinblastine, and podophyllotoxin need to be explored. This could provide the framework for future natural product production through genetic and metabolic engineering (Engels et al., 2008).

Bioreactors scaling up of production of secondary metabolites

This is the application of bioreactor system for large-scale cultivation of plant cells for the production of valuable bioactive compounds in an active field. Plant cells in liquid suspension offer a unique combination of physical and chemical environments that must be accommodated in large-scale bioreactor process. Some of the well known drawbacks of the cell suspension cultures include the instability of the productive cell lines, the slowness of the cell growth and limited knowledge about the metabolic pathway (Fulzele, 2005). There are indications

that sufficient oxygen supply and proper mixing in airlift bioreactors may not be suitable for high density plant cell suspension cultures. Well known problem shear sensitivity and rapid setting characteristics of plant cell aggregates and cell floating tendencies of the cell cultures have to be solved when bioreactors are designed.

The main constraint for commercial exploitation of *in vitro* cultures is the scaling up at industrial level. Hairy roots, callus and suspension cultures are complicated when it comes to scaling up and pose unique challenges. Mechanical agitation causes wounding of hairy roots and leads to callus formation. With a product of sufficiently high value it is feasible to use batch fermentation, harvest the roots, and extract the product. For less valuable products it may be desirable to establish a packed bed of roots to operate the reactor in a continuous process for extended periods collecting the product from the effluent stream. Scale up becomes difficult in providing nutrients from both liquid and gas phases simultaneously. Meristem dependent growth of root cultures in liquid medium results in a root ball with young growing roots on the periphery and a core of older tissue inside. Restriction of nutrient oxygen delivery to the central mass of tissue gives rise to a pocket of senescent tissue. Due to branching, the roots form an interlocked matrix that exhibits a hairy root culture as a source of bioactive chemicals depends on development of suitable bioreactor system where several physical and chemical parameters must be taken into consideration.

Immobilization scaling up of secondary metabolite accumulation

Advances in scale-up approaches and immobilization techniques contribute to a considerable increase in the number of applications of plant cell cultures for the production of compounds with a high added value. Plant-derived compounds with cancer chemotherapeutic or antioxidant properties use rosmarinic acid (RA) and taxol as representative examples. Critical issues are thoroughly discussed already, including the dependence of *in vitro*, compound-specific production on culture growth and differentiation, elicitation strategies, physiological effects of immobilization, and the current status of scale-up production systems (Ramawat and Merillon, 2008).

Cell cultures of *Plumbago rosea* were immobilized in calcium alginate and cultured in Murashige and Skoog's basal medium containing 10 mM CaCl₂ for the production of plumbagin, an important medicinal compound (Komaraiah et al., 2003). Studies were carried to find out the impact of immobilization on the increased accumulation of this secondary metabolite. Immobilization in calcium alginate enhanced the production of plumbagin by three, two and one folds compared to that of control, un-crosslinked alginate and CaCl₂ treated cells respec-

tively. Cells subjected to combined treatments of chitosan, immobilization and *in situ* extraction showed a synergistic effect and yielded 92.13 mg g⁻¹ of plumbagin which is 21, 5.7, 2.5 times higher than control, immobilized, immobilized and elicited cells, respectively.

Conclusion

Advances in biotechniques, particularly methods for culturing plant cell cultures should provide new means for the commercial processing of even rare plants and the chemicals they provide. The advantage of this method is that it can ultimately provide a continuous, reliable source of natural products. The major advantage of the cell cultures include synthesis of bioactive secondary metabolites, running in controlled environment, independently from climate and soil conditions. The use of *in vitro* plant cell culture for the production of chemicals and pharmaceuticals has made great strides building on advances in plant science. The increased use of genetic tools and an emerging picture of the structure and regulation of pathways for secondary metabolism will provide the basis for the production of commercially acceptable levels of product. The increased level of natural products for medicinal purposes coupled with the low product yields and supply concerns of plant harvest has renewed interest in large-scale plant cell culture technology. Knowledge of biosynthetic pathways of desired phytochemicals in plants as well as in cultures is often still in its infancy, and consequently strategies needed to develop an information based on a cellular and molecular level. These results show that *in vitro* plant cell cultures have potential for commercial production of secondary metabolites. The introduction of newer techniques of molecular biology, so as to produce transgenic cultures and to effect the expression and regulation of biosynthetic pathways, is also likely to be a significant step towards making cell cultures more generally applicable to the commercial production of secondary metabolites.

REFERENCES

- Aftab F, Akram S, Iqbal J (2008). Estimation of fixed oil from various explants and *in vitro* callus cultures of Jojoba (*Simmondsia chinensis*). Pak. J. Bot. 40: 1467-1471.
- Agarwal M, Kamal R (2007). Studies on flavonoid production using *in vitro* cultures of *Momordica charantia* L. Ind. J. Biotechnol. 6 : 277-279.
- Alikaridis F, Papadakis D, Pantelia K, Kephelas T (2000). Flavonolignan production from *Silybium marianum* transformed and untransformed root cultures. Fitoterapia 71: 379-384.
- Anderson LA, Roberts MF, Phillipson JD (1987). Studies on *Ailanthus altissima* cell suspension cultures. The effect of basal media on growth and alkaloid production. Plant Cell Rep. 6: 239-241.
- Andrijany VS, Indrayanto G, Soehono LD (1999). Simultaneous effect of calcium, magnesium, copper and cobalt on saponin steroids content in callus cultures of *Agave amaniensis*. Plant Cell Tiss. Org. Cult. 55: 103-108.

- Anitha S, Kumari BDR (2006). Stimulation of reserpine biosynthesis in the callus of *Rauvolfia tetraphylla* L. by precursor feeding. *Afr. J. Biotechnol.* 5: 659-661.
- Arellano J, Vazquez F, Villegas T, Hernandez G (1996). Establishment of transformed root cultures of *Perezia cuernavacana* producing the sesquiterpene quinone perezone. *Plant Cell Rep.* 15 : 455-458.
- Arrebola ML, Ringbom T, Verpoorte R (1999). Anthraquinones from *Isoplexis isabelliana* cell suspension cultures. *Phytochemistry* 52: 1283-1286.
- Arroo RRJ, Develi A, Meijers H, Van de Westerlo E, Kemp AK, Cores AF, Williams GJ (1995). Effect of exogenous auxin on root morphology and secondary metabolism in *Tagetes patula* hairy root cultures. *Physiol. Plant* 93 : 233-240.
- Arya D, Patn V, Kant U (2008). *In vitro* propagation and quercetin quantification in callus cultures of Rasna (*Pluchea lanceolata* Oliver & Hiern.). *Ind. J. Biotechnol.* 7 : 383-387.
- Asada Y, Li W, Yoshikawa T (1998). Isopyrenylated flavonoids from hairy root cultures of *Glycyrrhiza glabra*. *Phytochem.* 47 : 389-392.
- Asaka I, Li I, Hirokuni M, Asada Y, Furuya T (1993). Production of ginsenoside saponins by culturing ginseng (*Panax ginseng*) embryonic tissue in bioreactors. *Biotechnol. Lett.* 15: 1259-1264.
- Ayabe S, Iida K, Furuya T (1986). Induction of stress metabolites in immobilized *Glycyrrhiza echinata* cultured cells. *Plant Cell Rep.* 3: 186-189.
- Ayabe S, Takano H, Fujita T, Hirota H, Takahashi T (1990). Triterpenoid biosynthesis in tissue cultures of *Glycyrrhiza glabra* var. *glandulifera*. *Plant Cell Rep.* 9: 181-184.
- Babakov AV, Bartova LM, Dridze IL, Maisuryan AN, Margulis GU, Oganian RR, Voblikova VD, Muromtsev GS (1995). Cultures of transformed horseradish roots as source of fusicoccin-like lignans. *J. Plant Growth Reg.* 14 : 163-167.
- Badaoui H, Muguet B, Henry M (1996). Production of solamargine by *in vitro* cultures of *Solanum paludosum*. *Plant Cell Tiss. Org. Cult.* 45: 123-127.
- Balbuena TS, Santa-Catarina C, Silvera V, Kato MJ, Floh EIS (2009). *In vitro* morphogenesis and cell suspension culture establishment in *Piper solmsianum* DC. (Piperaceae). *Acta Bot. Bras.* 23: 229-236.
- Baldi A, Dixit VK (2008). Enhanced artemisinin production by cell cultures of *Artemisia annua*. *Curr. Trends in Biotechnol. Pharmacol.* 2 : 341-348.
- Banerjee S, Naqui AA, Mandal S, Ahuja PS (1994). Transformation of *Withania somnifera* (L.) Dunal by *Agrobacterium rhizogenes* : infectivity and phytochemical studies. *Phytochem Res.* 8 : 452-455.
- Barthe GA, Jourdan PS, McIntosh CA, Mansell RL (1987). Naringin and limonin production in callus culture and regenerated shoots from *Citrus* sp. *J. Plant Physiol.* 127: 55-65.
- Bassetti L, Hagendoorn M, Johannes T (1995). Surfactant-induced non-lethal release of anthraquinones from suspension cultures of *Morinda citrifolia*. *J. Biotechnol.* 39: 149-155.
- Baumert A, Groger D, Kuzovkina IN, Reisch J (1992). Secondary metabolites produced by callus cultures of various *Ruta* species. *Plant Cell Tiss. Org. Cult.* 28: 159-162.
- Berlin J, Ruegenhagen C, Dietze P, Fecker LF, Goddijn OJM, Hoge JHG (1993). Increased production of serotonin by suspension and root cultures of *Peganum harmala* transformed with a tryptophan decarboxylase cDNA clone from *Catharanthus roseus*. *Transgenic Res.* 2: 336-344.
- Binns AN, Tomashow JV (1988). Cell biology of *Agrobacterium* infection and transformation of plants. *Annu. Rev. Microbiol.* 42: 575-606.
- Bulgakov VP, Khodakovskaya MV, Labetskaya NV, Chermodev GK, Zhuravlev YN (1998). The impact of plant *rol C* oncogene on ginsenoside production by ginseng hairy root cultures. *Phytochemistry* 49: 1929-1934.
- Carvalho EB, Curtis WR (1998). Characterization of fluid-flow resistance in root cultures with a connective flow tubular bioreactor. *Biotechnol. Bioeng.* 60: 375-384.
- Chriqui D, Guivarch A, Dewitte W, Prinsen E, Van Onkelen H (1996). *Rol* genes and root initiation and development. *Plant Soil* 187: 47-55.
- Constabel CP, Towers GHN (1988). Thiarubrin accumulation in hairy root cultures of *Chaenatis douglasii*. *J. Plant Physiol.* 133 : 67-72.
- Davioud E, Kan C, Hamon J, Tempe J, Husson HP (1989). Production of indole alkaloids by *in vitro* root cultures from *Catharanthus trichophyllus*. *Phytochem.* 28 : 2675-2680.
- De-Eknamkul D, Ellis BE (1985). Effects of macronutrients of growth and rosmarine acid formation in cell suspension cultures of *Anchusa officinalis*. *Plant Cell Rep.* 4: 46-49.
- Delbeque JP, Beydon P, Chapuis L (1995). *In vitro* incorporation of radio labelled cholesterol and mevalonic acid into ecdysterone by hairy root cultures of a plant *Serrulata tinctoria*. *Eur. J. Entomol.* 92 : 301-307.
- Deno H, Yamagata T, Emoto T, Yoshioka T, Yamada Y, Fujita Y (1987). Scopalamine production by root cultures of *Duboisia myoporoides* II. Establishment of a hairy root culture by infection with *Agrobacterium rhizogenes*. *J. Plant Physiol.* 131 : 315-323.
- Devi CS, Murugesu S, Srinivasan VM (2006). Gymnemic acid production in suspension calli culture of *Gymnema sylvestris*. *J. appl. Sci.* 6 : 2263-2268.
- Dheeranapattana S, Wangprapa M, Jatisatienr A (2008). Effect of sodium acetate on stevioside production of *Stevia rebaudiana*. *Acta Hort. (ISHS).* 786 : 269-272.
- Downs CG, Christey MC, Davies KM, King GA, Seelye JF, Sinclair BK, Stevenson DG (1994). Hairy roots of *Brassica napus*: II glutamine synthase over expression alters ammonia assimilation and the response to phosphinothiric. *Plant Cell Rep.* 14: 41-46.
- Eibl R, Eibl D (2006). In *Plant Tissue Culture Engineering. Focus on Biotechnology*, vol 6. (Eds. Gupta, S. D. and Ibaraki) Springer: Berlin-Heidelberg-New York. pp. 203-227.
- Engels B, Dahm P, Jennewein S (2008). Metabolic engineering of taxadiene biosynthesis in yeast as a first step towards Taxol (Paclitaxel) production. *Metabolic Engineering* 10: 201-206.
- Farzami MS, Ghorbani M (2005). Formation of catechin in callus cultures and micropropagation of *Rheum ribes* L. *Pak. J. Biol. Sci.* 8 : 1346-1350.
- Fei HM, Mei KF, Shen X, Ye YM, Lin ZP, Peng LH (1993). Transformation of *Gynostemma pentaphyllum* by *Agrobacterium rhizogenes* saponin production in hairy root cultures. *Acta Bot. Sinica* 35 : 626-631.
- Fons F, Gargadanne A, Rapior S (2008). Culture of *Plantago* species as bioactive components resources: a 20 year review and recent applications. *Acta Bot. Gallica.* 155: 217-300.
- Francoise B, Hossein S, Halimeh H, Zahra NF (2007). Growth optimization of *Zataria multiflora* Boiss. Tissue cultures and rosmarinic acid production improvement. *Pak. J. Biol. Sci.* 10 : 3395-3399.
- Fukui H, Feroj HAFM, Ueoka T, Kyo M (1998). Formation and secretion of a new benzoquinone by hairy root cultures of *Lithospermum erythrorhizon*. *Phytochem.* 47 : 1037-1039.
- Fulcheri C, Morard P, Henry M (1998). Stimulation of the growth and the triterpenoid saponin accumulation of *Saponaria officinalis* cell and *Gypsophila paniculata* root suspension cultures by improvement of the mineral composition of the media. *J. Agric. Food Chem.* 46 : 2055-2061.
- Fulzele P (2005). Bioreactor for production of bioactive compounds. *IANCAS Bull.* 4: 35-42.
- Furmanova M, Syklovska BK (2000). Hairy root cultures of *Taxus x media* var. *Hicksii* Rehd. as a new source of paclitaxel and 10-deacetylpaclitaxin III. *Biotechnol. Lett.* 22: 606-616.
- Gao SL, Zhu DN, Cai ZH, Jiang Y, Xu DR (2004). Organ culture of a precious Chinese medicinal plant – *Fritillaria unibracteata*. *Plant Cell Tiss. Org. Cult.* 59: 197-201.
- Giri A, Banerjee S, Ahuja PS, Giri CC (1997). Production of hairy roots in *Aconitum heterophyllum* Wall. using *Agrobacterium rhizogenes*. *In Vitro Cell Dev. Biol. Plant* 33: 280-284.
- Giulietti AM, Parr AJ, Rhodes MJC (1993). Tropane alkaloid production in transformed root cultures of *Brugmansia candida*. *Panta Med.* 59 : 428-431.
- Goleniowski M, Tirppi VS (1999). Effect of growth medium composition on psilostachyinolides and altamisine production. *Plant Cell Tiss. Org. Cult.* 56: 215-218.
- Gopi C, Vatsala TM (2006). *In vitro* studies on effects of plant growth regulators on callus and suspension culture biomass yield from *Gymnema sylvestris* R.Br. *Afr. J. Biotechnol.* 5 : 1215-1219.
- Granicher F, Cristen P, Kaptanidis I (1995). Production of valepotriates by hairy root cultures of *Centranthes rubra* DC. *Plant Cell Rep.* 14:

- 294-298.
- Griffing WJ, Lin GD (2000). Chemotaxonomy and geographical distribution of tropane alkaloids. *Phytochemistry* 53: 623-637.
- Grzegorzczak I, Wysokinska H (2008). Liquid shoot culture of *Salvia officinalis* L. for micropropagation and production of antioxidant compounds : effect of triacontanol. *Acta Soc. Botanicorum Poloniae* 73 : 99-104.
- Halperin SJ, Flores HE (1997). Hyocyanine and proline accumulation in water stressed *Hyocyanus muticus* hairy root cultures. *In Vitro Cell Biol. Plant.* 33 : 240-244.
- Hamill JD, Robins RJ, Rhodes MJC (1989). Alkaloid production by transformed root cultures of *Cinchona ledgeriana*. *Planta Med.* 55 : 354-357.
- Hashimoto T, Yun DJ, Yamada Y (1993). Production of tropane alkaloids in genetically engineered root cultures. *Phytochem.* 32: 713-718.
- Hayashi T, Kasahara K, Sankawa U (1998). Efficient production of biologically active diterpenoid by leaf organ culture of *Scoparia dulcis*. *Cell Culture Biotechnol.* 46: 517-520.
- Hilton MG, Rhodes MJC (1993). Factors affecting the growth and hyocyanine production during batch culture of transformed roots of *Datura stramonium*. *Planta Med.* 59 : 340-344.
- Hiraoka, N., Bhatt, I.D., Sakurai, Y. and Chang, J.I. 2004. Alkaloid production by somatic embryo cultures of *Corydalis ambigua*. *Plant Biotechnol.* 21 : 361-366.
- Hohtola A, Jalonen J, Tolonen A, Jaakola L, Kamarainen T, Pakonen M, Karppinen K, Laine K, Neubauer P, Myllykoshi L, Gyorgy Z, Rautio A, Peltonen O (2005). Natural product formation by plants, enhancement, analysis, processing and testing. In : Sustainable use renewable natural resources – from principles to practices (Eds. Jalkanen, A. and Nygren, P). University of Helsinki Publication. pp. 34-69.
- Hook I (1994). Secondary metabolites in hairy root cultures of *Leontopodium alpinum* Cass (Edelweiss). *Plant Cell Tiss. Organ Cult.* 38 : 321-326.
- Ikenaga T, Oyama T, Muranaka T (1995). Growth and steroidal saponin production in hairy root cultures of *Solanum aculeatiss.* *Plant Cell Rep.* 14 : 413-417.
- Indira IR, Jayaraman G, Ramesh GA (2009). *In vitro* responses and production of phytochemicals of potential medicinal value in nutmeg, *Myristica fragrans* Houtt. *Indian J. Sci. Technol.* 2: 65-70.
- Ionkova I, Karting T, Alfermann W (1997). Cycloartane saponin production from hairy root cultures of *Astragalus mongholicus*. *Phytochem.* 45 : 1597-1600.
- Ionkova I, Karting T, Alfermann W (1997). Cycloartane saponin production in hairy root cultures of *Astragalus mongholicus*. *Phytochem.* 45 : 1597-15600.
- Ishimaru K Shimomura K (1991). Tannin production in hairy root cultures of *Geranium thunbergii*. *Phytochem.* 30 : 825-828.
- Ishimaru K, Sudo H, Salake M, Malsugama Y, Hasagawa Y, Takamoto S, Shimomura K Amaroggenin and amaroswertin and four xanthones from hairy root cultures of *Swertia japonica*. *Phytochem.* 29 : 1563-1565.
- Iwasa K, Takao N (1982). Formation of alkaloids in *Corydalis ophiocarpa* callus cultures. *Phytochem.* 21: 611-614.
- Jaziri M, Yoshimatsu K, Homes J, Vanhaelen M (1988). Tropane alkaloid production by hairy root cultures of *Datura stramonium* and *Hyocyanus niger*. *Phytochem.* 27 : 419-420.
- Jennewein S, Rithner CD, Williams RM, Croteau RB (2001). Taxol biosynthesis: Taxane 13 -hydroxylase is a cytochrome P450-dependent monooxygenase. *PNAS.* 98: 13595-13600.
- Jeong GA, Park DH (2007). Enhanced secondary metabolite biosynthesis by elicitation in transformed plant root system. *Appl. Biochem. Biotechnol.* 130 : 436-446.
- Jin JH, Shin JH, Kim JH, Chung IS, Lee HJ (1999). Effect of chitosan elicitation and media components on the production of anthraquinone colorants in madder (*Rubia akane* Nakai) cell culture. *Biotechnol. Bioprocess. Eng.* 4 : 300-304.
- Jobanovic V, Grubisic D, Giba Z, Menkovic N, Ristic M (1991). Alkaloids from hairy root cultures of *Anisodus luridus* (*Scolopia lurids* Dunal Solanaceae Tropane alkaloids). *Planta Med.* 2 : 102.
- Johnson T, Ravishankar GA, Venkataraman LV (1990). *In vitro* capsaicin production by immobilized cells and placental tissue of *Capsicum annuum* L. grown in liquid medium. *Plant Sci.* 70: 223-229.
- Jouanin L (1984). Restriction map of an agropine-type Ri-plasmid and its homologies with Ti-plasmids. *Plasmid* 12: 91-102.
- Jung G, Tepfer D (1987). Use of genetic transformation by the Ri T-DNA of *Agrobacterium rhizogenes* to stimulate biomass and tropane alkaloid production in *Atropa belladonna* and *Calystegia sepium* roots grown *in vitro*. *Plant Sci.* 50 : 145-151.
- Kanegae T, Kajiya H, Amano Y, Hashimoto T, Yamada Y (1994). Species -dependent expression of the hyoscyamine 6 -hydroxylase gene in the pericycle. *Plant Physiol.* 105: 483-490.
- Karppinen K, Hokkanen J, Tolonen A, Maltia S, Hohtola A (2007). Biosynthesis of hyperforin and adhyperforin from amino acid precursors in shoot cultures of *Hypericum perforatum*. *Phytochem.* 68 : 1038-1045.
- Kaul B, Staba EJ (1967). *Ammi visnaga* L. tissue cultures suspension on growth and examination for furanocoumarins. *Planta Med.* 15: 145-156.
- Khan T, Krupadanam D, Anwar Y (2008). The role of phytohormone on the production of berberine in the calli culture of an endangered medicinal plant, turmeric (*Coscinium fenestratum* L.) *Afr. J. Biotechnol.* 7 : 3244-3246.
- Khoury HE, Ibrahim RK, Rideau M (1986). Effect of nutritional and hormonal factors on growth and production of anthraquinone glucosides in cell-suspension culture of *Cinchona succirubra*. *Plant Cell Rep.* 5: 423-426.
- Kim JS, Lee SY, Park SU (2008). Resveratrol production in hairy root culture of peanut, *Arachis hypogaea* L. transformed with differet *Agrobacterium rhizogenes* strains. *Afr. J. Biotechnol.* 7 : 3788-3790.
- Kim OT, Bang KH, Shin YS, Lee MJ, Jang SJ, Hyun DY, Kim YC, Senong NS, Cha SW, Hwang B (2007) Enhanced production of asiaticoside from hairy root cultures of *Centella asiatica* (L.) Urban elicited by methyl jasmonate. *Plant Cell Rep.* 26: 1914-1949.
- Kim OT, Kim MY, Hong MH, Ahn JC, Huang B (2004). Stimulation of asiaticoside accumulation in the whole plant cultures of *Centella asiatica* (L.) Urban by elicitors. *Plant Cell Rep.* 23: 339-344.
- Kin N, Kunter B (2009). The effect of callus age, VU radiation and incubation time on trans-resveratrol production in grapevine callus culture. *Tarim Bilimleri Dergisi* 15: 9-13.
- Kinney AJ (1998). Manipulating flux through plant metabolic pathways. *Curr. Opin. Plant Biol.* 1: 173-178.
- Kiong AL, Mahmood M, Fodzillan NM, Daud SK (2005). Effects of precursor supplementation on the production of triterpenes by *Centella asiatica* callus culture. *Pak. J. Biol. Sci.* 8 : 1160-1169.
- Kisiel W, Stojakowska A (1997). A sesquiterpene coumarin ether from transformed roots of *Tanacetum parthenium*. *Phytochem.* 46 : 5515-5516.
- Kisiel W, Stojakowska A, Malaz J, Kohlmunzer S (1995). Sesquiterpene lactones in *Agrobacterium rhizogenes* transformed hairy root cultures of *Lactuca virosa*. *Phytochem.* 40 : 1139-1140.
- Ko KS, Ebizuka Y, Noguchi H, Sankawa U (1995). Production of polypeptide pigments in hairy root cultures of *Cassia* plants. *Chem. Pharm. Bull.* 43 : 274-278.
- Koblitz H, Koblitz D, Schmauder HP, Groger D (1983). Studies on tissue cultures of the genus *Cinchona* L. alkaloid production in cell suspension cultures. *Plant Cell Rep.* 2: 122-125.
- Komaraiah P, Ramakrishna SV, Reddanna P, Kavikishore PB (2003). Enhanced production of plumbagin in immobilized cells of *Plumbago rosea* by elicitation and *in situ* adsorption. *J. Biotechnol.* 10 : 181-187.
- Kornfeld A, Kaufman PB, Lu CR, Gibson DM, Bolling SF, Warber SL, Chang SC, Kirakosyan A (2007). The production of hypericins in two selected *Hypericum perforatum* shoot cultures is related to differences in black gland culture. *Plant Physiol. & Biochem.* 45 : 24-32.
- Krolicka A, Kartanowicz R, Wosinska S, Zpitter A, Kaminski M, Lojkowska E (2006). Induction of secondary metabolite production in transformed callus of *Ammi majus* L. grown after electromagnetic treatment of the culture medium. *Enzyme and Microbial Technology* 39: 1386 – 1389.
- Kuch JSH, Mac Kenzie IA, Pattenden G (1985). Production of chrysanthermic acid and pyrethrins by tissue cultures of *Chrysanthemum*

- cinerariaefolium*. Plant Cell Rep. 4: 118-119.
- Kunvari M, Paska C, Laszlo M, Gyurjan I (1999). Effect of different media parameters on the growth and verbascoside production on *Plantago media* L. tissue culture. In: 47th annual congress of the Society for medical plant Research, Amsterdam 26-30 July 1999.
- Kuroyanagi M, Arakava T, Mikami Y, Yoshida K, Kawahar N, Hayashi T, Ishimaru H (1998) Phytotoxins from hairy roots cultures of *Hyocymus albus* treated with methyl jasmonate. J. Nat. Prod. 61 : 1516-1519.
- Kusakari K, Yokoyama M, Inomata S (2000). Enhanced production of saikosaponins by root culture of *Bupleurum falcatum* L. using two step control of sugar concentration. Plant Cell Rep. 19: 1115-1120.
- Lee SY, Cho SJ, Park MH, Kim YK, Choi JI, Park SU (2007). Growth and rutin production in hairy root culture of buck weed (*Fagopyrum esculentum*) Prep. Biochem. Biotechnol. 37 : 239-246.
- Lee SY, Xu H, Kim YK, Park SU (2007). Rosmarinic acid production in hairy root cultures of *Agastache rugosa* Kuntze. World J. Microbiol. Biotechnol. 20 : 969-972.
- Lee-Parsons CWT, Rogce AJ (2006). Precursor limitations in methyl jasmonate-induced *Catharanthus roseus* cell cultures. Plant Cell Rep. 25: 607-612.
- Lessard P (1996). Metabolic engineering, the concept coalesces. Nat. Biotechnol. 14: 1654-1655.
- Li W, Li M, Yang DL, Xu R, Zhang Y (2009). Production of podophyllotoxin by root culture of *Podophyllum hexandrum* Royle. Electronic J. Biol. 5: 34-39.
- Li YC, Wen-Yi T (2009). Effects of paclitaxel-producing fungal endophytes on growth and paclitaxel formation of *Taxus cuspidate* cells. Plant Growth Regul. 58: 97-105.
- Li YC, Wen-Yi T (2009). Interactions of Taxol-producing endophytic fungus with its host (*Taxus spp.*) during Taxol accumulation. Cell Biol. Int. 33: 106-112.
- Liu KCS, Yang SH, Roberts MF, Phillipson JD (1990). Production of canthin-6-one alkaloids by cell suspension cultures of *Brucea javanica* (L.) Merr. Plant Cell Rep. 9: 261-263.
- Maharik N, Elgengaihi S, Taha H (2009). Anthocyanin production in callus cultures of *Crataegus sinaica* Bioss. Intrn. J. Academic Res. 1 : 30-34.
- Malpathak NP, David SB (1986). Flavor formation in tissue cultures of garlic (*Allium sativum* L.). Plant Cell Rep. 5: 446-447.
- Marconi PL, Selden LM, Cslcena EN, Alvarez MA, Pitta-Alvarez SI (2008). Changes in growth and tropane alkaloid production in long-term culture of hairy roots of *Brugmansia candida* Elect. J. Integrative Biosci. 3: 38-44.
- Matsumoto T, Tanaka N (1991). Production of phytoecdysteroids by hairy root cultures of *Ajuga reptans* var. *attropurpurea*. Agric. Biol. Chem. 55: 10-25.
- Mehrotra S, Kukreja AK, Khanuja SPS, Mishra BN (2008). Genetic transformation studies and scale up of hairy root culture of *Glycyrrhiza glabra* in bioreactor. 11: 717-728.
- Min JY, Jung HY, Kang SM, Kim YD, Kang YM, Park DJ, Prasad DT, Choi MS (2007). Production of tropane alkaloids by small-scale bubble column bioreactor cultures of *Scopolia parviflora* adventitious roots. Bioresour. Technol. 98: 1748-1753.
- Mirjalili MH, Moyano E, Bonfill M, Cusido RM, Palazon J (2009) Steroidal lactones from *Withania somnifera*, an antioxidant plant for novel medicine. Molecules 14 : 2373-2393.
- Misra N, Misra P, Datta SK, Mehrotra S (2005). In vitro biosynthesis of antioxidants from *Hemidesmus indicus* R.Br. cultures. In vitro Dev. Biol. Plant. 41 : 285-290.
- Moreno PRH, Heijden RVD, Verpoorte R (1993). Effect of terpenoid precursor feeding and elicitation on formation of indole alkaloids in cell suspension cultures of *Catharanthus roseus*. Plant Cell Rep. 12: 702-705.
- Morimoto H, Murai F (1989). The effect of gelling agents on planuotol accumulation in callus cultures of *Croton sublyratus* Kurz. Plant Cell Rep. 8: 210-213.
- Moyano E, Fornalé S, Palazón J, Cusidó RM, Bonfill M, Morales C, Piñol MT (1999). Effect of *Agrobacterium rhizogenes* T-DNA on alkaloid production in Solanaceae plants. Phytochem. 52: 1287-1292.
- Muranaka T, Ohakawa H, Yamada Y (1992). Scopolamine release into media by *Duboisia leichhardtii* hairy root clones. Appl. Microbiol. Biotechnol. 37 : 554-559.
- Murthy HN, Dijkstra C, Anthony P, White DA, Davey MR, Powers JB, Hahn EJ, Paek KY (2008). Establishment of *Withania somnifera* hairy root cultures for the production of Withanoloid A. J. Integ. Plant Biol. 50 : 915-981.
- Nair AJ, Sudhakaran PR, MAdhusudanan JR, Ramakrishna SU (1992). Berberine synthesis by callus and cells suspension cultures of *Cosciniun fenestratum*. Plant Cell Tiss. Org. Cult. 29 : 7-10.
- Nakajima K, Hashimoto T (1999). Two tropinone reductases that catalyze opposite stereospecific reductions in tropane alkaloid biosynthesis are localized in plants root with different cell-specific patterns. Plant Cell Physiol. 40: 1099-1107.
- Namdeo AG (2007). Plant cell elicitation for production of secondary metabolites: A review. Pharmacognosy Rev. 1: 69-79.
- Narasimhan S, Nair M (2004). Release of berberine and its crystallization in liquid medium of cell suspension cultures of *Cosciniun fenestratum* (Gaertn.) Colebr. Curr. Sci. 86 : 1369-1371.
- Nazif NM, Rady MR, Seif MM (2000). Stimulation of anthraquinone production in suspension cultures of *Cassia acutifolia* by salt stress. Fitoterapia 71: 34-40.
- Nikolaeva TN, Zagoskina NV, Zaprometov MN (2009). Production of phenolic compounds in callus cultures of tea plant under the effect of 2,4-D and NAA. Russ. J. Pl. Physiol. 56 : 45-49.
- Nilsson O, Olsson O (1997). Getting to the root: the role of the *Agrobacterium rhizogenes* rol genes in the formation of hairy roots. Physiol. Plant. 100: 463-473.
- Nin S, Bennici A, Roselli G, Mariotti D, Schiff S, Magherini R (1997). Agrobacterium mediated transformation of *Artemisia absinthum* L. (worm wood) and production of secondary metabolites. Plant Cell Rep. 16 : 725-730
- Nurchgani N, Solichatun S, Anggarwulan E (2008). The reserpine production and callus growth of Indian snake root (*Rauvolfia serpentina* (L.) Benth. ex Kurz.) cultured by addition of Cu²⁺. Biodiversitas 9: 177-179.
- Ogasawara T, Cheba K, Tada M (1993). Production in high-yield of a naphthaquinone by a hairy root culture of *Sesamum indicum*. Phytochem. 33 : 1095-1098.
- Okrslar V, Plaper I, Kovac M, Erjavec A, Obermajer T, Rebec A, Ravnikar M, Zel J (2007). Saponins in tissue culture of *Primula veris* L. In Vitro Cell Dev. Biol. Plant. 43: 644-651.
- Olivira AJB, Koika L, Reis FAM, Shepherd SL (2001). Callus culture of *Aspidosperma ramiflorum* Muell.-Arg. Growth and alkaloid production. Acta Scientia 23: 609-612.
- Olivira AJB, Koike L, Reis FAM, Shepherd SLK (2001). Callus culture of *Aspidosperma ramiflorum* Muell.-Arg.: growth and alkaloid production. Acta Scientia 23 : 609-612.
- Oostdam A, Mol JNM, Vanderplas LHW (1993). Establishment of hairy root cultures of *Linum flavum* producing the lignan 5-methoxy podophyllotoxin. Plant Cell Rep. 12 : 474-477.
- Orihara, Y. and Furuya, T. 1990. Production of theanine and other - glutamyl derivatives by *Camellia sinensis* cultured cells. Plant Cell Rep. 9: 65-68.
- Paek YW, Hwang SJ, Park DH, Hwang B (1996). Multiplication and transformation of medicinal plants for production of useful secondary metabolites: II establishment of hairy root cultures of *Centella asiatica*. J. Plant Biol. 39: 161-166.
- Palazon J, Cusido RM, Gonzalo J, Bonill M, Morales C, Pinol MT (1998). Relation between the amount of rol C gene product and indole alkaloid accumulation in *Catharanthus roseus* transformed root cultures. J. Plant Physiol. 153: 712-718.
- Palazon J, Moyano E, Bonfill M, Cusido RM, Pinol MT (2006). In Floriculture, Ornamental and Plant Biotechnology: Advances and Topical Issues. (Ed. Teixeira da Silva, J. A.) Global Science Books, Ltd: London, UK. pp. 209-221.
- Palazon J, Pinol MT, Cusido RM, Morales C, Bonfill M (1997). Application of transformed root technology to the production of bioactive metabolites. Recent Res. Dev. Plant Phys. 1: 125-143.
- Park SU, Lee SY (2009). Anthraquinone production by hairy root culture of *Rubia akane* Nakai : Influence of media and auxin treatment. Sci. Res. Essays 4 : 690-693.
- Parr AJ, Hamill JD (1987). Relationship between Agrobacterium rhizogenes transformed hairy roots and intact uninfected *Nicotiana*

- plants. *Phytochem.* 26 : 3241-3245.
- Paska C, Innocent G, Kunvari M, Laszlo M, Szilagyi L (1999). Lignan production by *Ipomea cairica* callus culture. *Phytochem.* 52 : 879-883.
- Philipson JD (1990). Plants as source of valuable products. In: B.V. Chalwood and M.J. Rhodes (Eds.), *Secondary products from plant tissue culture*, Oxford, Clarendon Press. pp.1-21.
- Poornasri DB, Vimala A, Sai I, Chandra S (2008). Effect of cyanobacterial elicitor on neem cell suspension cultures. *Ind. J. Sci. Technol.* 1: 1-5.
- Qu JG, Yu XJ, Zhang W, Jin MF (2006). Significant improved anthocyanins biosynthesis in suspension cultures of *Vitis vinifera* by process intensification. *Sheng Wu Gong Cheng Xue Bae.* 22: 299-305.
- Quiala E, Barbon R, Jimenez E, Feria MD, Chavez M, Capote A, Perez N (2006) Biomass production of *Cymbopogon citratus* (DC.) Stapf. A medicinal plant in temporary immersion systems. *In Vitro Cell Dev. Biol. Plant.* 42 : 298-300.
- Rahnama H, Hasanloo T, Shams MR, Sephehrifar R (2008). Silymarin production by hairy root culture of *Silybium marianum* (L.) Gaertn. *Iranian J. Biotechnol.* 6: 113-118.
- Rajasekaran T, Rajendran L, Ravishankar GA, Venkataraman LV (1991). Influence of nutrient stress on pyrethrin production by cultured cells of pyrethrum (*Chrysanthemum cinerariaefolium*). *Curr. Sci.* 60: 705-707.
- Rama Rao, B., Vijay kumar, D., Amrutha, R.N., Jalaja, N., Vaidyanath, K., Maruthi, A., Rao, S., Polararupu R, Kavi kishor PB (2008). Effect of growth regulators, carbon source and cell aggregate size on berberine production from cell cultures of *Tinospora cordifolia* Miers. *Curr. Trends in Biotechnol. Pharmacy* 2: 269-276.
- Ramachandra SR, Ravishankar GA (2002). Plant cell cultures: Chemical factories of secondary metabolites. *Biotechnol. Adv.* 20: 1001-153.
- Ramani S, Jayabaskaran C (2008). Enhanced catharine and vindoline production in suspension cultures of *Catharanthus roseus* by ultraviolet-B light. *J. Mol. Signal.* 3 : 9-14.
- Ramawat K, Merillon S (2008). Secondary metabolite production from plant cell cultures : the success stories of rosmarinic acid. *Plant Cell Tiss. Org. Cult.* 16 : 85-100.
- Ramirez M, Alpizer L, Quiroz J, Oropeza C (1992). Formation of L-canavanine in *in vitro* cultures of *Canavalia ensiformis* (L.) DC. *Plant Cell Tiss. Org. Cult.* 30: 231-235.
- Rao KV, Venkanna N, Lakshmi NM (1998). *Agrobacterium rhizogenes* mediated transformation of *Artemisia annua*. *J. Sci. Ind. Res.* 57 : 773-776.
- Rao RS, Ravishankar GA (2002). Plant tissue cultures; chemical factories of secondary metabolites. *Biotechnol. Adv.* 20: 101-153.
- Roat C, Ramawat KG (2009). Elicitor induced accumulation of stilbenes in cell suspension cultures of *Cayratia trifoliata* (L.) Domin. *Plant Biotechnol. Rep.* 3: 135-138.
- Rothe G, Hachiya A, Yamada Y, Hashimoto T, Dräger B (2003). Alkaloids in plants and roots cultures of *Atropa belladonna* overexpressing putrescine N-methyltransferase. *J. Exp. Bot.* 54: 2065-2070.
- Rueffer M, Bauer M, Zenk MH (1994). The formation of corydaline and related alkaloids in *Corydalis cava* *in vivo* and *in vitro*. *Can. J. Chem.* 72 : 170-175.
- Ruffer M, Baue M, Zenk MH (1994). The formation of corydaline and related alkaloids in *Corydalis cava* *in vivo* and *in vitro*. *Can. J. Chem.* 72: 170-175.
- Saito K, Yoshimatsu K, Murakoshi T (1990). Genetic transformation of foxglove (*Digitalis purpurea*) by chimeric foreign genes and production of cardioactive glycosides. *Plant Cell Rep.* 9 : 121-124.
- Salma U, Rahman MSM, Islam S, Haque N, Jubair TA, Haque AKMF, Mukti IJ (2008). The influence of different hormone concentration and combination on callus induction and regeneration of *Rauvolfia serpentina* (L.) Benth. *Pak. J. Biol. Sci.* 11: 1638-1641.
- Santarem ER, Astarita LV (2003). Multiple shoot formation in *Hypericum perforatum* L. and hypericin production. *Brazilian J. Plant Physiol.* 15 : 21-26.
- Santos PM, Figueredo AC, Olivera MM, Barroso JG, Pedro LG, Deans SG, Younus AKM (1998). Essential oils from hairy root cultures and from fruits and roots of *Pimpinella anisum*. *Phytochem.* 48 : 455-460.
- Santos-Gomes PC, Seabra RM, Andrade PB, Fernandes-Ferreira MM (2002). Phenolic antioxidant compounds produced by *in vitro* shoots of sage (*Salvia officinalis* L.). *Plant Sci.* 162: 981-987.
- Sasaki K, Udagava A, Ishimaru H, Hayashi T, Alfermann AW, Nakanishi F, Shimomura K (1998). High forskolin production in hairy roots of *Coleus forskohlii*. *Plant Cell Rep.* 17 : 457-459.
- Sato F, Hashimoto T, Hachiya A, Tamura KI, Choi KB, Morishige T, Fujimoto H, Yamada Y (2001). Metabolic engineering of plant alkaloid biosynthesis. *Proc. Natl. Acad. Sci. USA.* 2 : 367-372.
- Sato K, Yamazaki T, Okuyama E, Yoshihira K, Shimomura K (1991). Anthraquinone production by transformed root cultures of *Rubia tinctorum* : influence of phytohormones and sucrose concentration. *Phytochem.* 30 : 2977-2978.
- Sauerwein M, Yamazaki T, Shimomura K (1991). Hernandulcin in hairy root cultures of *Lippia dulcis*. *Plant Cell Rep.* 9: 579-581.
- Savary BJ, Flores HE (1994). Biosynthesis of defense related proteins in transformed root cultures of *Trichosanthes kirilowii* Maxim. var. *japonicum* (Kitam). *Plant Physiol.* 106 : 1195-1204.
- Schenk RV, Hildebrandt AC (1972). Medium techniques for induction and growth of monocotyledonous and dicotyledonous plant cell cultures. *Can. J. Bot.* 50: 199-204.
- Schmeda-Hirschmann G, Jordan M, Gertn A, Wilken D, Hormazabal E, Tapia AA (2004). Secondary metabolite content in *Fabiana imbricata* plants and *in vitro* cultures. *Z. Naturforsch.* 5: 48 -54.
- Schripsema J, Ramos-Valdivia A, Verpoorte R (1999). Robustaquinones, novel anthraquinones from an elicited *Cinchona robusta* suspension culture. *Phytochem.* 51: 55-60.
- Sevon N, Oksman-Caldentey KM (2002). *Agrobacterium rhizogenes*-mediates transformation: Root cultures as a source of alkaloids. *Planta Med.* 68: 859-868.
- Shalaka DK, Sandhya P (2009). Micropropagation and organogenesis in *Adhatoda vasica* for the estimation of vasine. *Pharmacognosy Magazine* 5 : 539-363.
- Shinde AN, Malpathak N, Fulzele DP (2009). Induced high frequency shoot regeneration and enhanced isoflavones production in *Psoralea corylifolia*. *Rec. Nat. Prod.* 3: 38-45.
- Shinde AN, Malpathak N, Fulzele DP (2009). Induced high frequency shoot regeneration and enhance a isoflavones production in *Psoralea corylifolia*. *Rec. Nat. Prod.* 3 : 38-45.
- Shohaeh AM, Murthy HN, Hahn EJ, Paek KY (2007). Methyl jasmonate induced overproduction of eleuthrosides in somatic embryos of *Eleutherococcus senticosus* cultured in bioreactors. *Elect. J. Biotechnol.* 10: 633-637.
- Shrivastava N, Patel T, Srivastava A (2006). Biosynthetic potential of *in vitro* grown callus cells of *Cassia senna* L. var. *senna*. *Curr. Sci.* 90 : 1472-1473.
- Singh DN, Verma N, Raghwanishi S, Shukla DK, Kulshreshtha DK (2006). Antifungal anthraquinone from *Saprosma fragrans*. *Bioorg. Med. Chem. Lett.* 16 : 4512-4514.
- Skrzypek Z, Wysokinski H (2003). Sterols and terpenes in cell cultures of *Hyssopus officinalis* L. *Ver Lag der Zeitschrift fur Naturforschung.* D. 312.
- Smetanska I (2008). Production of secondary metabolites using plant cell cultures. *Adv. Biochem. Eng. Biotechnol.* 111: 187 – 228.
- Staniszewska I, Krolicka A, Mali E, Ojkowska E, Szafranek J (2003). Elicitation of secondary metabolites in *in vitro* cultures of *Ammi majus* L. *Enzyme Microbiol. Technol.* 33 : 565-568.
- Sujanya S, Poornasri DB, Sai I (2008). *In vitro* production of azadirachtin from cell suspension cultures of *Azadirachta indica*. *J. Biosci.* 33: 113-120.
- Suzuki K, Yamada Y, Hashimoto T (1999). Expression of *Atropa belladonna* putrescine N-methyltransferase gene in root pericycle. *Plant Cell Physiol.* 40: 289-297.
- Tada H, Nakashima T, Kuntake H, Mori K, Tanaka M, Ishimaru K (1996). Polyacetylenes production by hairy root cultures of *Campanula medium* L. *J. Plant Physiol.* 147 : 617-619.
- Tada H, Shimomura K, Ishimaru K (1995). Polyacetylenes in *Platycodon grandiflorum* hairy roots and campanulaceous plants. *J. Plant Physiol.* 145 : 7-10.
- Taha HS, El-Rahman A, Fathalla M, Kareem AE, Aly NE (2008). Successful application for enhancement and production of anthocyanin

- pigment from calli cultures of some ornamental plants. *Aust. J. Basic and Appl. Sci.* 2: 1148-1156.
- Tallevi SG, Dicosmo F (1998). Stimulation of indole alkaloid content in vanadium treated *Catharanthus roseus* suspension cultures. *Planta Med.* 54: 149-152.
- Tanaka N, Takao M, Matsumoto T (2004). Vincamine production in multiple shoot culture derived from hairy roots of *Vinca major*. *Plant Cell Tiss. Org. Cult.* 41 : 61-64.
- Taya M, Mine K, Kinoka M, Tone S, Ichi T (1992). Production and release of pigments by cultures of transformed hairy roots of red beet. *J. Ferment Bioeng.* 73 : 31-36.
- Teshima D, Ikeda K, Satake M, Aoyama T, Shimomura K (1988). Production of emetic alkaloids by *in vitro* culture of *Cephaelis ipecacuanha* A. Richard. *Plant Cell Rep.* 7: 278-280.
- Tiwari KK, Trivedi M, Guang ZC, Guo GQ, Zheng GC (2007). Genetic transformation of *Gentiana macrophylla* with *Agrobacterium rhizogenes* : growth and production of secoiridoid glucoside gentiopicoside in transformed hairy root cultures. *Plant Cell Rep.* 26 : 199-210.
- Trotin F, Moumou Y, Vasseur J (1993). Flavonol production by *Fagopyrum esculentum* hairy roots and normal root cultures. *Phytochem.* 33 : 929-931.
- Trypsteen M, Van-Lijsekettens M, Van Severen R, Van Montagu M (1991). *Agrobacterium rhizogenes* mediated transformation of *Echinacea purpurea*. *Plant Cell Rep.* 10 : 85-89.
- Tumova L, Rimakova J, Tuma J, Duscck J (2006). *Silybum marianum in vitro* – flavolignan production. *Plant Cell Environ.* 52 : 454-458.
- Umamaheswari A, Lalitha V (2007). *In vitro* effect of various growth hormones in *Capsicum annum* L. on the callus induction and production of Capsaicin. *J. Plant Sci.* 2: 545-551.
- Varindra S, Saikia R, Sandhu S, Gosal SS (2000). Effect of nutrient limitation on capsaicin production in callus culture derived from pericarp and seedling explants of *Capsicum annum* L. varieties. *Pl. Tissue Cult.* 10: 9-16.
- Verma PC, Singh D, Rahman L, Gupta MM, Banerjee S (2002). *In vitro* studies in *Plumbago zeylanica* : rapid micropropagation and establishment of higher plumbagin yielding hairy root cultures. *J. Plant Physiol.* 159 : 547-552.
- Vineesh VR, Fijesh PV, Jelly LC, Jaimsha VK, Padikkala J (2007). *In vitro* production of camptothecin (an anticancer drug) through albino plants of *Ophiorrhiza rugosa* var. *decumbens*. *Curr. Sci.* 92: 1216-1219.
- Vinterhalter B, Jankovic T, Sovikin L, Nikolic R, Vinterhalter D (2008). Propagation and xanthone content of *Gentianella austriaca* shoot cultures. *Plant Cell Tiss. Org. Cult.* 94 : 329-335.
- Wagiah ME, Alam G, Wiryowidagdo S, Attia K (2008). Improved production of the indole alkaloid cathin-6-one from cell suspension cultures of *Brucea javanica* (L.) Merr. *Ind. J. Sci. Technol.* 1 : 1-6.
- Waller GR, Mac Vean CD, Suzuki T (1983). High production of caffeine and related enzyme activities in callus cultures of *Coffea arabica* L. *Plant Cell Rep.* 2: 109-112.
- Wang PJ, Huang CI (1982). Production of saikosaponins by callus and redifferentiated organs of *Bupleurum falcatum* L. In: *Plant Tissue Culture* (Ed. Fujiwara), Maruzen, Tokyo. pp. 71-72.
- Whitmer S, Van der Heijden R, Verpoorte R (2002). In *Plant Biotechnology and Transgenic Plants* Oksman-Caldentey, K. -M.; Barz W.H., Eds.; Marcel & Dekker: New York-Basel; pp. 373-405.
- Wijnsma R, Go TKA, Weerden IN, Harkes PAA, Verpoorte R, Svendsen AB (1985). Anthraquinones as phytotoxins in cell and tissue cultures of *Cinchona* sp. *Plant Cell Rep.* 4: 241-244.
- Williams GR, Doran PM (2000). Hairy root culture in a liquid-dispersed bioreactor: characterization of spatial heterogeneity. *Biotechnol. Progr.* 16: 391-401.
- Williams RD, Ellis BE (1992). Alkaloids from *Agrobacterium rhizogenes* transformed *Papaver somniferum* cultures. *Phytochem.* 32 : 719-723.
- Wilson, P.D.G. 1997. In *Hairy roots: Culture and applications*. Doran, P. M., Ed.; Harwood Academic Publishers: Amsterdam, pp. 179-190.
- Wink M, Alfermann AW, Franke R, Wetterauer B, Distl M, Windhovel J, Krohn O, Fuss E, Garden, H, Mohagheghzaden A, Wildi E, Ripplinger P (2005). Sustainable bioproduction of phytochemicals by plant *in vitro* cultures: anticancer agents. *Plant Genetic Resour.* 3: 90-100.
- Wyk BEV, Wink M (2004). Medicinal plants of the world. Pretoria, Briza.
- Xu H, Kim YK, Suh SY, Udin MR, Lee SY, Park SU (2008). Deourisin production from hairy root culture of *Angelica gigas*. *J. Korea Soc. Appl. Biol. Chem.* 51: 349-351.
- Yagi, A, Shoyama Y, Nishioka I (1983). Formation of tetrahydroantrone glucosides by callus tissue of *Aloe saponaria*. *Phytochem.* 22: 1483-1484.
- Yamanaka M, Ishibashi K, Shimomura K, Ishimaru K (1996). Polyacetylene glucosides in hairy root cultures of *Lobelia cardinalis*. *Phytochem.* 41 : 183-185.
- Zambryski P, Tempe J, Schell J (1989). Transfer and function of T-DNA genes from *Agrobacterium* Ti and Ri plasmids in plants. *Cell* 56: 193-201.
- Zhang P, Peng-Peng Z, Long-Jiang Y (2009). An Endophytic Taxol-Producing Fungus from *Taxus media*, *Cladosporium cladosporioides* MD2. *Curr. Microbiol.* 59: 227-232.
- Zhao J, Davis LC, Verpoorte R (2005). Elicitor signal transduction leading to production of plant secondary metabolites. *Biotechnol. Adv.* 23: 283-333.
- Zhao J, Zhu W, Hu Q (2001). Enhanced catharanthine production in *Catharanthus roseus* cell cultures by combined elicitor treatment in shake flasks and bioreactors. *Enzyme Microb. Technol.* 28: 673-681.
- Zhou Y, Hirotsani M, Yoshikawa T, Furuya T (1997). Flavonoids and phenylethanoids from hairy root cultures of *Scutellaria baicalensis*. *Phytochem.* 42 : 69-72.