

Full Length Research Paper

# Assessment of *Salmonella enterica* Contamination in Camel Milk Across Kenya

Matofari, J. W., Shitandi, A.\*, Shalo, P. L., Nanua, N. J., Younan, M.

Guildford Dairy Institute, Egerton University Njoro, P.O. Box 536 Egerton 20107 Kenya.

Accepted 13 April, 2024

This study was undertaken to conduct a baseline risk analysis of raw camel milk with special emphasis on *Salmonella enterica* serovars. Cross-sectional studies were designed to investigate the prevalence of *S. enterica* serovars in a major camel milk production zone of Kenya. A total of 196 samples were assessed for possible presence of *S. enterica*. The samples included composite milk from the individual camel udders, bulk milk from collection and market centres, faeces, soil and water samples. Of the 196 samples tested, 43% (84/196) were found to contain *Salmonella* species. Out of the 84, only 31% (26/84) was positively identified as *S. enterica*. *S. enterica* was found in all the sample categories that represented the camel milk production environment. The results suggest that raw camel milk contamination by *S. enterica* was influenced by post-harvest handling of the product rather than camel infection by the pathogen. It was concluded that a need exists to formulate better regulation strategies for the safe handling of camel milk on rural Kenyan farms.

**Key words:** Camel milk; *Salmonella enterica*, Milk safety, Kenya.

## INTRODUCTION

Camel milk constitutes 12% of 3 billion litres of the milk produced nationally in Kenya. The average camel milk production per annum is 0.366 million litres and 25% of the national population (30m), especially those living in the low lands depend on camel milk.

Since commercial exploitation of camel milk in Kenya has grown tremendously, there is a growing public health concern on its safety as it is informally marketed.

The handling of informally marketed milk has been reported to affect the safety and quality of milk with reports of food poisoning due to consumption of camel milk also being reported (El-Nawawi et al., 1982; Bachman, 1992).

*Salmonella* infection in camels has been reported in various countries, including Sudan (Curasson, 1998) Palestine (Olitziki and Ellenbogen, 1943), French North Africa (Donatien and Boue, 1994), USA (Bruner and Mu-ran, 1949) and, more recently, from Somalia (Cheyne et al., 1977), Ethiopia (Pegram and Tareke, 1981), Egypt (Refai et al., 1984; Yassein, 1985; Osman, 1995) and UAE (Wernery, 1992). Faye (1997) reported that *S. enterica* typhimurium and *S. enterica* enteritidis are more pre-valent in camels. Healthy camels can be carriers of *Sal-*

*monella* and organisms have been isolated from faeces and lymphnodes on slaughter of camels (Zaki, 1956; Hamada et al., 1983; El-Nawawi et al., 1982; Refai et al., 1984; Yassien, 1985; Selim, 1990).

Camels that are chronic carriers of *Salmonella* may present a human health hazard through consumption of camel products like milk. This study was undertaken to conduct a baseline risk analysis of raw camel milk with special emphasis on *S. enterica* serovars. The study covered the rural zones in Kenya.

## MATERIALS AND METHODS

Cross-sectional studies were designed to investigate the prevalence of *S. enterica* serovars. The samples for *S. enterica* analysis included the composite milk from the individual camel udders, bulk milk from collection and market centres, faecal, soil and water samples. For the composite and bulk milk, the containers were shaken to mix the samples well. A cup was used to take the sample after shaking the container. About 25 ml of the milk sample from the cup was poured into a sterile screw cap universal bottle and then capped. This was then put in a cool box maintained at 4°C. Faecal samples were taken using sterile cotton swabs wrapped on splint wood sticks. The cotton swab stick was pushed into the rectum in a screw manner of the lactating female camel whose milk sample had been taken. The swab was immediately transferred into sterile Stuart Transport Medium (Oxoid) in a screw cap Bijou bottle. The handle stick was broken and the swab remained in the transport me

\*Corresponding author. E-mail: anakalos@gmail.com.

**Table 1.** Prevalence of *Salmonella enterica* in sampled regions and markets.

Region	N	Udders	Bulk milk	Faeces	Water	Soil
AK	36	10	5	4	1	1
BL	35	11	5	2	1	0
CN	20	4	1	0	3	0
DG	30	5	0	0	3	1
Enh	19	2	2	0	1	0
FM	16	3	2	2	3	0
GI	14	4	4	2	2	0
HNm*	26	0	0	0	0	0
Total	196	39	19	10	14	2
Incidence (%)		20	10	5	7	1

\*Market isolates, A-G are isolates from different production areas and Districts of Northern Kenya. AK-Kalacha, BL-Logologo, CN- Ngurnit, DG- Gudas, Enh- North horr, FM- Moyale, GI- Isiolo, HNm- Nairobi market.

dium. The bottle was capped and put in the cool box. Water samples were also taken from the boreholes or wells that were being used as sources of drinking water for the camels. Borehole samples were taken by pumping the water out for 5 min and then taking the sample by slanting the mouth of a sterile glass bottle towards the nozzle of the water pipe. Samples (500 ml) were taken and the bottles capped. Well water was taken by lowering a bucket on a rope into the well, when the bucket reached the water level, it was swirled to stir the water and then lowered to scoop the water. The water from the bucket was then poured into a sterile 500 ml glass bottle, capped and then placed in the cool box. For soil samples, 200 g was scooped in the middle of the *boma* and wrapped in clean polythene papers, then transferred to the cool box. All samples were transported to the laboratory at KARI-Marsabit in a cool box within 12 h of sampling. A total of 196 samples were taken and analysed for the isolation of *S. enterica*.

#### Enumeration of *S. enterica* from the various sample categories

Twenty-five ml of milk and water and 25 g of faecal and soil samples were pipetted and weighed, respectively, and inoculated into 225 ml of Buffered Peptone Water (BPW) (Oxoid) as a pre-enrichment. The suspensions were incubated at 37°C for 24 h. After the incubation, the mixtures were shaken gently, and using a sterile pipette, 1 ml was pipetted and transferred into 10 ml Rappaport Vassiliadis medium (Oxoid). After incubation in a water bath at 42°C for 24 h, a loopful of growth was streaked onto both Xylose Lysine Desoxycolate (XLD) (Oxoid) agar and Brilliant Green agar (BGA) (Oxoid). The agar plates were incubated at 37°C for 24 to 48 h. Colonies that appeared dark on XLD and those that appeared pink on BGA were taken to be non-lactose fermenters and were purified on MacConkey agar (Oxoid). The purified colonies on Mac-Conkey agar were inoculated into the Triple Sugar Iron (TSI) (Oxoid) agar slants by stabbing the butt and streaking the slant. The colonies that appeared shiny, yellow and mucoid were taken to be lactose fermenters and were also purified on MacConkey agar.

Pure isolates (3-4 colonies) that were lactose negative on culture and purified on MacConkey agar were inoculated into 10 ml preparations of fermentable sugars (glucose, lactose, manitol, sorbitol, citrate, Triple Sugar Iron (TSI) and urea) with phenol red as the indicator and incubated at 37°C for 24 h.

#### Serotyping of *S. enterica* isolates

For serological identification, a slide agglutination test using O-grouping polyvalent sera and Vi serum was used. All the reagents

were left to reach room temperature. On a clean microscope slide, a drop of antiserum was placed at one end and a drop of sterile normal saline (0.85% NaCl) was placed at the opposite end of the same slide. Three to four colonies from the non-selective media were suspended in 0.3 ml sterile saline and a dense cell suspension was made. One loopful of the cell suspension was put onto each of the drops of serum and normal saline, and mixed well. The cell suspension and normal saline served as controls. The slide was gently shaken for 1 min. Agglutination within 1 min was regarded as positive for polyvalent O-group. Whenever any isolate agglutinated with one of the polyvalent O-sera, it was again tested against the corresponding monovalent antisera, using the same procedure as above. The reason for testing with monovalent O was to avoid cross-reaction with non-salmonella genera like *Escherichia*, *Shigella*, *Citrobacter* and *Proteus*. The name of the serum that produced agglutination was considered as the name of the O-antigen possessed by the tested *Salmonella* spp.

When there was no agglutination after 1 min from the above test, the same above procedure was carried out using Vi serum instead of the polyvalent sera. When a positive reaction was found, a dense suspension of the organism in sterile saline was made and autoclaved at 121°C for 15 min. After cooling to room temperature, the agglutination test was repeated with polyvalent serum and Vi serum using the heated cells. Live cells that were negative with polyvalent serum and positive with Vi serum before heating, and positive with polyvalent serum and negative with Vi serum after heating, were taken to be *S. enterica* Typhi.

For H-antigen, a tube agglutination test was used. H polyvalent and monovalent sera were left to reach room temperature. Pure cultures of suspected *Salmonella* spp. (grown for 8 h at 37°C in non-selective broth) were diluted with an equal volume of saline containing 1% formalin. An aliquot (0.4 - 0.5 ml) of the antigen suspension was added to 0.05 ml of each specific H serum in small test tubes. A control was prepared that only contained the antigen suspension. The tubes were shaken well for 2 min, allowed to stand in a water bath at 50 - 52°C for 1 h and then observed for agglutination. The name of the serum that produced agglutination corresponds to the name of the H-antigen possessed by the test organism.

## RESULTS

The prevalence of *S. enterica* in the sampled regions and the market outlets is illustrated in Table 1. Of the 196 samples tested, 84 were found to contain *Salmonella*

**Table 2.** Serological identification of *Salmonella enterica* isolates.

Serovar isolate	n	Poly O- Ag	Factor O-Ag	Subgroup	Serotype
Paratyphi	15	A-G	C-factor 6, 7	C <sub>1</sub>	Paratyphi C
Typhi	11	A-G	D-factor 9 (Vi)	D <sub>1</sub>	Typhi

**Table 3.** *Salmonella enterica* serovars present in 196 samples taken from camel milk production environment

Sample category	No. of positive isolates	Incidence%	<i>S. enterica</i> serovar (n)
Milk(udder/bulk)	15	57	Paratyphi C (7), Typhi (8)
Water	5	19	Paratyphi C (3), Typhi (2)
Faeces	5	19	Paratyphi C (4), Typhi (1)
Soil	1	4	Paratyphi C (0), Typhi (1)
Total	26		ParatyphiC (14),Typhi (12)

species. Out of the 84, 31% (26/84) were positively identified as *S. enterica*. *Salmonella* incidence was more prevalent at the production level than market level. At market level, there was no isolation of *Salmonella*. Milk was the main source of *Salmonella* (individual camel milk and pooled milk). However, it is not quite clear whether the origin of *Salmonella* was endogenous, that is, from the camel itself, or exogenous, that is., from the camel environment. At the udder (milking) level, the incidence of *S. enterica* was twice as high (incidence of 20%) than at the collecting centres (incidence 10%) (Table 1). Based on serology, Table 2 illustrates the two serovars that were identified based on poly O and Factor O agglutination tests. *S. enterica* was found in all the sample categories that represented the camel milk production environment, these being the milk, faeces, water and soil (Table 3). The serovar typhi was found in all four sample categories accounting for 46% (12/26) of the positively identified *S. enterica* while serovar Paratyphi C was found in three categories accounting for 54% (14/26) but missing in soil category (Table 3).

Individual camel udders and collecting centers (bulk) had a higher incidence (57%) of *S. enterica* contamination as compared to other categories. The serovars typhi and Paratyphi C had almost the same prevalence. Water and faeces accounted for 19% of *S. enterica* contamination (Table 3).

## DISCUSSION

*S. enterica* was found in all the sample categories that represented the camel milk production environment. Milk at the udder harvesting level and the bulk milk at collection points had the highest incidence (> 20%) of the pathogen (Table 3). The environment in which any food is produced is a key factor contributing to its quality. The environment in which camel milk is produced has been proven in this study to be a contributory factor to the

contamination of the camel milk by *S. enterica*. The fact that *Salmonella* were found in a wide spectrum of the categories sampled, including water, faeces, soil and milk is an indication that faecal contamination of camel milk production and market chain is common.

Two serovars of *S. enterica* were isolated from camel milk in this study; *S. enterica* Paratyphi C and *S. enterica* Typhi. The latter is a strict human-associated serovar that causes septicaemia. This host-adapted serovar is transmissible through faecal contamination of food (Kenneth, 2005). Camel milk is produced in the same environment where human faecal waste is deposited, as pastoralists normally do not construct pit latrines. Whenever rains come, the faecal waste could be carried as surface runoff to the nearest water body, commonly streams that are seasonal, dams, boreholes and shallow wells. These are, in turn, used as sources of drinking water for camels and humans. The milkers and milk-handlers of camel milk could be carriers of *S. enterica* Typhi.

The paratyphi group occurs in almost all domestic animals and is transmissible to man. They are known to cause food poisoning in humans (Pietzsch, 1981). Serovar Paratyphi C occurs sporadically and its mode of transmission is direct or indirect contact with faeces or contaminated food (PHAC, 2001). As a health hazard, serovar Paratyphi C causes bacterial enteric fever. It has also been reported by Kariuki et al. (1999) that Paratyphi C is resistant to most antimicrobials. In this study, raw camel milk contamination by *S. enterica* was more strongly influenced by post-harvest handling of the product rather than *S. enterica* infection of the camel (Table 1). In some studies, the presence of *S. enterica* in camels has been reported in disease assessment, especially camel calf diarrhoea (Salih et al., 1998a, 1998b; Nation et al., 1996; Malik et al., 1967; Ambwani and Jaktar, 1973; Wernery, 1992) and in lymph-nodes and intestines of slaughtered dromedaries in Egypt (Rafai et al., 1984; Yassien, 1985).

The lower prevalence of *Salmonella* in raw camel milk in this study does not mean food borne illness may not be caused, but in fact should be considered a potentially hazardous situation. The presence of the *Salmonella* in camel milk must be supported by several factors in the environment or chain of production and marketing. For *S. enterica* to contaminate the camel milk, it must have gone through the chain of infection. It must have had a source (host) and a mode of transmission to the milk. The milk has all the factors that support its growth. The sources of the pathogen constitute the risk factors that may be associated with the prevalence or incidence of the same pathogen in the environment. In this case, pastoralists, camels, milkers, milk-handlers, equipment used in milking and handling milk, water, soil, etc. are the likely sources of *Salmonella* in the environment of camel milk production. The pastoralists and the camels may be healthy carriers, and they may persistently shed the pathogen in the environment and through the milk. The pathogen finds its way into other transmissible avenues like water, soil, milk and equipment. This cycle forms a web of causation of the pathogen in the environment. In the pastoralist environment, there is no human or animal waste disposal system. The same water sources are used for domestic work and animal watering. This web of causation of the pathogen, based on the risk factors mentioned above, is maintained at production level. This may explain the reason why there is high incidence of *S. enterica* at production level.

There was no *Salmonella* isolated at the market level (Table 1). The time lapse from production to market centres seems to be a factor in the apparent absence of the *Salmonella* from the milk. The ambient temperatures at the production level (ASAL) are normally high, during the transportation of the milk; there is no temperature control hence microbial growth is not limited. The situation is made worse by the poor infrastructure that characterises these areas and the long distances to the lucrative markets in urban centres. Under these conditions, the milk undergoes physico-chemical changes due to the unlimited multiplication of the microbial population. This leads to acid development that may selectively inhibit *S. enterica* from multiplying (Foster and Spector, 1995; Abee et al., 1995; Nousiainen, 1993; Juven et al., 1991). Intrinsic factors of camel milk, such as cationic peptides that have antimicrobial properties may also inhibit *S. enterica* from multiplying. The survival tactics of *S. enterica* in these conditions include developing resistance to these cationic peptides (Christensen et al., 1988; Kagan et al., 1990; Cotter et al., 2000), entering viable but non-culturable (VBNC) state (Erikson et al., 2001; Anvian et al., 2001; Chmielewski et al., 1995) and escaping from the extracellular environment of the milk to the intracellular environment using the leukocytes in milk (Gallan et al., 1992; Rosenshine et al., 1993) hence macrophage survival (Dunlap et al., 1992) may explain the apparent absence of the *Salmonella* from the market milk.

## Conclusion

This study has shown that camel milk within the studied area is contaminated with *S. enterica* and there is a clear indication of faecal contamination of camel milk. The serovars involved were *S. enterica* serotype Typhi and *S. enterica* serotype Paratyphi C. *S. enterica* serotype Typhi is highly host-adapted to humans. This suggests that there is direct and indirect human faecal contamination of the camel milk and water through the chain of production and marketing.

## ACKNOWLEDGMENTS

The authors are grateful to the Division of Research and Extension, Egerton University for funding the research and the Department of Dairy Science for access to the laboratories.

## REFERENCES

- Abee TL, Krokell J, Hill C (1995). Bacteriocins: Modes of action and potentials in food preservation and control of food poisoning. *Int. J. Food Microbiol.* 28: 169-185.
- Ambwani H, Jaktar P (1973). *Salmonella* infections of camel in Bikaner. *Indian Vet. J.* 50 (1): 100-102.
- Anvian YA, Weiner R, Johnson J, de Rezende C, Joseph S (2001). *Salmonella enterica* serovar Typhimurium DT 104 displays a rugose phenotype. *Appl. Environ. Microbiol.* 67: 4048-4056.
- Bruner D, Moran (1949). *Salmonella* infections of domestic animals. *Conell Vet.* 39 (1): 53-63.
- Cheyne A, Pegram R, Cartright C. (1977). An outbreak of Salmonellosis in camels in North East of the Somali Democratic Republic. *Tropical Animal Health Production.* 9 (4): 238-240.
- Chmielewski RN, Frank J (1995). Formation of viable but non-culturable *Salmonella* during starvation in chemically defined solutions. *Lett. Appl. Microbiol.* 20: 380-384.
- Christensen B, Fink J, Merrifield RB, Mauzerall D (1988). Chemical-forming properties of cepropins and related model compounds incorporated into planar lipid membranes *Proc. National Academy Sci. USA.* 85: 5072-5076.
- Cotter P, DiRita VJ (2000). Bacterial virulence gene regulation. An evolutionary perspective. *Ann. Rev. Microbiol.* 54: 519-65.
- Curason G (1998). Une maladie du dromadaire analogue au farcin du boeuf. *Bull. Soc. Cent. Med. Vet. (Supplement to Rec. Med. Vet.)* 94 71: 491-496
- Dunlap NE, Benjamin WB, Bery AK, Eldridge JH, Brides DE (1992). A safe site for *Salmonella* Typhimurium is within splenic polymorphonuclear cells. *Microbiol. Pathog.* 13: 181-190.
- El-Nawawi FH, El-Dere H, Sayed A (1982). *Salmonella* in slaughter animals. *Arch. F. ubensmittel hygiene* 33: 33-36.
- Erickson E, Mallinson T, Gupte A, Joseph W (2001). *Salmonella* spp. indifferent levels of water activity in closed microcosms. *J. Ind. Microbiol. Biotechnol.* 26: 222-225.
- Faye B (1997). Guide de l'élevage du dromadaire. Sanofi Sante Nutrition Animale, La Ballastiere-BP126, 33501 Libourne, Cedex, France. pp. 115-116.
- Foster JW, Spector MP (1995). How *Salmonella* survive against the odds. *Ann. Rev. Microbiol.* 49: 145-74.
- Gallan J, Pace J, Hayman J (1992). Involvement of the epidermal growth factor receptor in the invasion of cultured mammalian cells by *Salmonella* Typhimurium *Nature.* 357: 588-589.
- Hamada S, El-Sawah H, Sherif I, Joussef M, Hidik M (1983). *Salmonella* of the mesenteric lymphnodes of slaughtered cattle, buffaloes and camels. *J. Arab. Vet. Med. Ass.* 23(4): 272-277.

- Juven B, Meinersmann RJ, Stern NJ (1991). Antagonistic effects of Lactobacilli and Pediococci to control intestinal colonization of human enteropathogens in life poultry. *J. Appl. Bacteriol.* 70: 95-103.
- Kagan BL, Selsted ME, Ganz T, Albert J, Lehrer RI (1990). Antimicrobial defensin peptides from voltage dependent ion permeable channels in planar lipid bilayer membranes. *Proc. Natl. Acad. Sci. USA* 87: 210-214.
- Kariuki S, Cheesbrough J, Mavridis AK, Hart CA (1999). Typing of *Salmonella enterica* serotype Paratyphi C isolates from various countries by plasmid profiles and pulsed-field gel electrophoresis. *J. Clin. Microbiol.* 1: 2058-2062.
- Kenneth T (2005). Salmonella and Salmonellosis In: *Todar textbook of bacteriology*, University of Wisconsin-Madison, Department of Bacteriology.
- Malik PD, Datta SK, Singh IP, Kalra DS (1967). Salmonella serotypes from camel in India. *J. Res. Punjab Agric. University. Ludhiana.* 4: 123-126.
- Nation G, Moore JE, Tinson AH, MacAlmont M, Murphy PG, Harron D (1996). Treatment of Salmonella and Campylobacter-associated diarrhoea in camels (*Camelus dromedarius*) with enrofloxacin. *European Congress of Chemotherapy, Glasgow, Scotland, May 14-17.*
- Olitzki L, Ellenbogen V (1943). A Salmonella strain isolated from camels in Palestine. *J. Comp. Physiol. Thei.* 53 (1): 75-79.
- Osman AR (1995). *Path. Study on intestinal infections in camels.* Thesis. Faculty of Veterinary Medicine Cairo University.
- Pegram RG, Tareke F. (1981). Observation on the health of Afar livestock. *Ethopian Vet. J.* 5: 11-15.
- PHAC (Public Health Agency of Canada) (2001). *Salmonella Paratyphi C. Material safety data sheets: Health Canada.*
- Pietzsch O (1981). Salmonella In: *Blobel H, Schliesser T (Hrsg): Handbuch der bakteriellen infektionen bei Tieren. Bd. Jena, VEB Gustar Fischer.* 3: 344-452.
- Refai M, EL-Said W, Osman K, Lofti Z, Safwat EE, Elias S (1984). Salmonella in slaughtered camels in Egypt. *ZagaZig Vet. J.* 9: 266-267.
- Rosenshine J, Finlay BB (1993). Exploitation of host-signal transduction pathways and cytoskeletal functions by invasive bacterial. *Bio. Essays.* 15: 17-24.
- Salih O, Mohamed O, Shigidi MT (1998a). The epidemiological factors associated with camel calf diarrhoea in Eastern Sudan. *Proc. Intl. Meeting on camel production and future perspectives. May 2-3. Fac. of Agric. Sci. Al Ain, UAE.*
- Shigidi MTO, Mohamed HO, Mcdough A, Chang Y (1998b). The bacterial causes of camel calf diarrhoea in Eastern Sudan. *Proc. Intl. Meeting on camel production and future perspectives. May 2-3. Fac. of Agric. Sci. Al Ain, UAE.*
- Selim AM (1990). Salmonellosis in camels in Egypt. In: *Wardeh MF, Wilson RT, Saied AA, Proc. Inter. Conf. On camel prod. and improve. Dec, Tobruk, Libya.* pp. 10-13
- Wernery U (1992). The prevalence of Salmonella infections in camels (*Camelus dromedarius*) in the UAE. *Br. Vet. J.* 148 (5): 445-450.
- Yassiem NA (1985). *Salmonella in slaughtered camels.* MVSci. Thesis. Faculty of Veterinary Medicine Cairo University.
- Zaki OA (1956). The incidence of Salmonella infections in camels. *J. Egypt Publicat on Health Association.* 3 (2): 75-79.