

Full Length Research Paper

Effects of morphogenetic and diurnal variability on the hypericin content in St. John's Wort (*Hypericum perforatum* L.)

Oya Kaçar*, Erdinç Göksu and Nedime Azkan

Uludag University, Faculty of Agriculture, Department of Field Crops, Bursa, Turkey.

Accepted 12 June, 2020

This research was conducted to investigate effects of morphogenetic and diurnal variability on the hypericin content of St. John's Wort (*Hypericum perforatum* L.) populations originating from Turkey for maximum utilization of the active substance in plants. During 2002 and 2003, field trials were conducted at Uludag University, Faculty of Agriculture, Department of Field Crops, Bursa, Turkey. Samples of bud, flower and capsule of *H. perforatum* L. populations were collected in the second year of cultivation (2003). Factorial arrangements of three *H. perforatum* L. populations (Bursa, Edirne, Izmir), three part of plant (bud, flower, capsule) and six collection h (8:00, 10:00 a.m; 12:00 (noon), 2:00, 4:00, 6:00 p.m) were evaluated in a completely randomized block design with three replications. Hypericin content (%) in parts of *H. perforatum* L. populations was determined according to DAC (1986). Consequently, the content of hypericin in the examined populations varied from 0.260% in Bursa to 0.283% in Izmir. Evaluating plant parts revealed that the hypericin content both in floral parts (0.309%) and buds (0.308%) were higher than capsule tissues (0.208%) for all populations. When collection hours were examined for the hypericin content of plants, the highest content was recorded at 10:00 a.m (0.279%) and the lowest value at 4:00 p.m (0.272%) and 6:00 p.m (0.272%). As a result, this study showed that the highest hypericin ratio was determined in flowers and buds generally collected between 8:00 and 10:00 a.m within a day for examined *Hypericum* populations.

Key words: St. John's Wort, *Hypericum perforatum* L., morphogenetic variability, diurnal variability, hypericin content.

INTRODUCTION

Hypericum perforatum L, also known as St. John's Wort, is a herbaceous perennial plant belonging to the *Clusiaceae* family (Upton et al., 1997). *H. perforatum* L. (St. John's Wort) is widespread in Mediterranean region, Europe, Asia, North and South Africa, Western and Eastern North America (Jensen et al., 1995). Although it was known from ancient Greek and Roman times, St. John's Wort has recently become a widely popular herbal remedy due to its antidepressive effects (Linde et al., 1996; DeSmet and Mohen, 1996).

Hypericum contains numerous compounds with docu-

mented biological activity groups that have stimulated the most interesting compounds including the naphthodianthrones hypericin and pseudohypericin, a broad range of flavonoids quercetin, quercitrin, amentoflavone and hyperin, the phloroglucinols hyperforin and adhyperforin, the essential oil and xanthenes (Upton et al., 1997). From a pharmacological point of view, the hypericins are the most interesting compounds of *H. perforatum* L. (Patocka, 2003). Because of their use for mild depression treatment, clinical trials suggest that hypericin is the primary biologically active compound (Lavie et al., 1995; Linde et al., 1996). The naphthodianthrones, hypericin and pseudohypericin are known to contribute to the antidepressant action of this species, and most of *Hypericum* phytomedicines are currently standardized according to their hypericin content (Briskin, 2000). Hypericin also

*Corresponding author. E-mail: okacar@uludag.edu.tr. Tel: +90 224 2941523. Fax: +90 224 442 88 36.

Table 1. Climatic data for the experimental field during the growing period of *H. perforatum* L.

Month	Mean temperature (°C)			Rainfall (mm)			Relative humidity (%)		
	Long term	2002	2003	Long term	2002	2003	Long term	2002	2003
1	5.5	3.2	8.6	88.8	62.3	65.3	74.8	65.4	68.3
2	6.2	8.9	2.7	77.5	42.7	106.2	72.5	67.2	71.2
3	8.3	10.3	4.4	69.8	87.9	33.1	71.7	71.3	64.0
4	12.9	11.5	9.9	62.9	126.5	112.1	69.9	76.0	70.4
5	17.8	17.3	18.8	50.0	50.5	45.7	69.2	67.9	67.7
6	22.1	23.0	23.8	30.4	25.2	2.4	61.1	62.1	54.8
7	24.5	26.7	25.3	24.0	49.9	-	58.8	64.4	64.4
8	24.1	24.6	25.6	18.9	31.1	-	60.4	65.2	65.2
9	20.1	20.8	19.2	40.1	67.2	66.9	65.8	70.3	70.3
10	15.6	15.8	16.6	60.4	119.3	125.1	72.1	75.2	75.2
11	12.3	10.7	10.1	76.3	67.9	64.5	74.4	72.6	77.3
12	7.6	5.0	6.2	99.9	28.8	91.0	74.2	67.0	71.8
Total	-	-	-	699.0	759.3	712.3	-	-	-
Mean	14.8	14.8	14.3	-	-	-	68.7	68.7	68.4

shows a significant antiviral and antiretroviral activity (Meruelo et al., 1988; Vlietinck et al., 1998).

Hypericin and pseudohypericin may function as defensive allelochemicals in St. John's Wort (Fields et al., 1990). Hypericin is found in dark glandular structures visible located on the epidermal surface of leaves, flowers and to some extent the stems of plants (Fields et al., 1990; Curtis and Lerstern, 1991; Fornasiero et al., 1998; Crompton et al., 1988; Sirvent et al., 2002)

The formation of secondary metabolites is typical of the plants depend on both genetic and environmental factors. Besides climatic conditions of the growing site, particularly edaphic factors also play an important role in variation of these secondary metabolites (Franz, 1983; Palevitch, 1987). Variations of hypericin that derivate naphthodianthrone depend on not only genetic characteristics but also environmental factors, plant development stage, plant parts, harvest time, processing and storage methods (Bombardelli and Morazzoni, 1995; Büter et al., 1998; Jensen et al., 1995; Palevitch, 1991; Upton et al., 1997; Büter and Büter, 2002). Sirvent et al. (2002) stated that further studies are needed to investigate the relationships between the hypericin content and these factors.

The objective of this study was to assess the effects of plant parts (bud, flower and capsule) and collecting hour on hypericin content of three *H. perforatum* L. populations obtained from different regions of Turkey.

MATERIALS AND METHODS

During 2002 and 2003, three populations of *H. perforatum* L. (Bursa, Edirne, Izmir) were cultivated on the experimental field of Uludag University, Faculty of Agriculture, Department of Field Crops, Bursa (40° 11' N, 29° 04' E). The soil was clay loam, slightly alkaline and rich in phosphorus and potassium containing 1.8%

organic matter. Long-term average of annual total precipitation is 699 mm, mean temperature for the whole year is 14.3°C and relative humidity is 68.7%. In 2002 total precipitation 759.3 mm year⁻¹, mean temperature for the whole year was 14.8°C and relative humidity was 68.7%; in 2003 total precipitation 712.3 mm year⁻¹, mean temperature for the whole year was 14.3°C and relative humidity was 68.4%. The weather conditions during the experimental period are presented in Table 1.

The populations were propagated by seeds and thus they were regarded as populations. *Hypericum* seedlings were pre-grown in the green house during February to May, 2002 and transferred to the field on May 20/21, 2002. Seedlings were planted in 40 x 20 cm distances. Total 60 kg/ha nitrogen and 60 kg/ha P₂O₅ were applied in spring and autumn of each year. During the vegetation period, plots were irrigated as needed.

Samples of bud, flower and capsule of *H. perforatum* L. populations were collected at different collection h (8:00, 10:00 a.m; 12:00 (noon), 2:00, 4:00, 6:00 p.m) within a day in the second year (2003) of cultivation. The *Hypericum* plants could not reach their final size and morphology before the second year of cultivation. Similar to our observations, Büter et al. (1998) also stated that in the first year plants were smaller with less main shoots and less flowers resulting in considerably lower total plant yields. Moreover Osinska and Weglarz (2001) reported that the optimal time for harvest appeared to be the full blooming stage, especially in the second year of vegetation as far as the mass of herb and the content of investigated compounds are concerned. Similarly, Kordana and Zalecki (1997) reported that in the second year of cultivation yields of raw material, increased by 100% and hypericin content increased by 50%, compared with the first year. The relatively high hypericin content in the second year was possibly due to the higher proportion of flowers, leaves on the top drug herbage analysed. Therefore, in the present study, the observation and measurements were obtained from the plants of second year (2003).

Each replicate comprised a bulk sample of five individual plants. From each plant, 25 buds at onset of flowering (closed buds with yellow petals already visible), flower at full flowering (fully opened flowers) and capsule at seed formation (brown capsules) were removed and combined in order to obtain one extract per population, plant part and collection hour. Plant samples of *H. perforatum* L. were collected in 30.05.2003 for buds, 06.06.2003 for flowers and 27.06.2003 for capsules. Finally, harvested parts of plant were

Table 2. Effect of different populations, different parts of plant and their interactions on hypericin content (%) in *H. perforatum* L.

Parts of plant	Populations		
	Bursa	Edirne	Izmir
Bud	0.308 b*	0.309 b	0.308 b
Flower	0.306 b	0.312 a	0.308 b
Capsule	0.166 e	0.226 d	0.233 c
Mean	0.260 b	0.282 a	0.283 a

*Means with the same letter are not significantly different at $p=0.05$.

LSD 0.05 for population means: 0.001144.

LSD 0.05 for population x plant parts: 0.001982.

air-dried in a closed room at 25 - 30°C and stored at room temperature at dark conditions.

Extraction of hypericin

Hypericin content (%) in parts of *H. perforatum* L. populations was determined according to DAC (1986). The parts of harvested five individual plants were well homogenized in order to eliminate the variability between plants. Air dried plant material was powdered by a coffee mill. A 1 g powder was Soxhlett-extracted with chloroform. The extract was evaporated to dryness and the dried powder was then extracted with methanol. The hypericin content in methanol extracts was determined by a spectrophotometer at 590 nm.

Statistical analyses

Factorial arrangements of three *H. perforatum* L. populations (Bursa, Edirne, Izmir), three part of plant (bud, flower, capsule) and six collection hours (8:00, 10:00 a.m.; 12:00 (noon), 2:00, 4:00, 6:00 p.m) were evaluated in a completely randomized block design with three replications. All data were subjected to analysis of variance for each character using MSTAT-C (version 2.1, Michigan State University, 1991) and MINITAB (University of Texas at Austin) software. For mean separation, the F-protected least significant difference (LSD) was calculated at the 0.05 probability level according to Steel and Torrie (1980).

RESULTS AND DISCUSSION

Hypericin contents of *H. perforatum* L. populations, different parts of plant, different collection hours and their interactions were presented in Tables 2, 3 and 4. Significant differences were detected in the mean concentrations of hypericin of investigated factors and their interactions.

The content of hypericin in the examined populations varied from 0.260% in Bursa to 0.283% in Izmir. The average hypericin content of Izmir population (0.283%) was found to be similar to those of Edirne (0.282%) (Table 2).

Evaluating parts of plant, it was also found that hypericin content in both floral parts (0.309%) and buds

Table 3. Effect of different parts of plant, collection hour with in a day and their interactions on hypericin content in *H. perforatum* L.

Collection hour	Parts of plant		
	Bud	Flower	Capsule
8:00 a.m	0.309 b*	0.310 a	0.207 e
10:00 a.m	0.306 b	0.310 a	0.222 c
12:00 p.m (noon)	0.308 b	0.308 b	0.213 d
2:00 p.m	0.309 b	0.307 b	0.206 e
4:00 p.m	0.310 a	0.309 b	0.198 f
6:00 p.m	0.307 b	0.307 b	0.203 e
Mean of Plant Parts	0.308 a	0.309 a	0.208 b

*Means with the same letter are not significantly different at $p=0.05$.

LSD 0.05 for Parts of Plant: 0.001144.

LSD 0.05 for Parts of Plant x Collection Hour: 0.002803.

(0.308%) were higher than that of capsule tissues (0.208%) for all populations (Table 3). Regarding interactions between population and parts of plant, the distribution of the average amounts of hypericin in buds were similar to all populations. Significantly more hypericin (0.312%) were found in the flowers of Edirne population (Table 2). Also, hypericin content of capsules was found lower in Bursa population (0.166%) than those in other populations. Thus, the lowest hypericin content was determined in Bursa population (0.260%) (Table 2). Earlier investigations of *H. perforatum* L. in different countries indicated a higher hypericin concentration in the flowers compared to the other parts of plant (Cellarova et al., 1994; Schütt, 1996; Constantine and Karchesy, 1998; Tekelova et al., 2000; Southwell and Bourke, 2001; Sirvent et al., 2002).

Previous studies showed that hypericin contents varied among different parts of plant, ranging from 0.003 - 0.609% in flowers (Constantine and Karchesy, 1998; Melikian et al., 1998; Campbell et al., 1992; Büter et al., 1998; Sirvent et al., 2002; Poutaroud and Girardin, 2004), 0.215% in flower and bud (Southwell and Campbell, 1991), 0.03% in leaves+stems+flowers (Constantine and Karchesy, 1998), 0.038-0.045% in top leaves; 0.029-0.035% in bottom leaves; 0.012-0.0115% in side stem; 0.004% in main stem; 0.073-0.077% in capsules (Southwell and Campbell, 1991; Campbell et al., 1992). It has been reported that dried flowers can yield as 1.8% hypericin (Upton et al., 1997). As a result of these studies, hypericin concentrations generally declined towards from flowers to capsules. Repcak and Martonfi (1997) found that hypericin and pseudohypericin were located mainly in the petals and in the stamina of *Hypericum*. This finding may explain the progressive decrease of dianthrone content in older flowers, as their stamina has lost anthers and developing fruits have only rests of the corona. Similarly, the petals of small buds represent a smaller portion of their weight. The content of hypericin increased from the first bud phases to flowers just opened, then decreased from over the blooming to unripe

Table 4. Effect of different populations, collection hour with in a day and their interactions on hypericin content in *H. perforatum* L.

Collection hour	Populations			Mean of collection hour
	Bursa	Edirne	Izmir	
8:00 a.m	0.266 d*	0.285 b	0.275 c	0.276 b
10:00 a.m	0.261 d	0.282 b	0.295 a	0.279 a
12:00 p.m (noon)	0.255 e	0.284 b	0.290 a	0.277 b
2:00 p.m	0.267 d	0.283 b	0.272 c	0.274 b
4:00 p.m	0.260 d	0.278 c	0.279 c	0.272 c
6:00 p.m	0.250 e	0.282 b	0.285 b	0.272 c

* Means with the same letter are not significantly different at $p=0.05$.

LSD 0.05 for Collection Hour: 0.001618.

LSD 0.05 for Population x Collection Hour: 0.002803.

fruits (Tekelova et al., 2000). Kitanov (2000) reported that the level of hypericins was very low before budding period, but increased rapidly and reached its maximum at the stage of budding and blossoming. Hypericin content gradually decreased towards ripening of the fruits and reached the same level as before budding.

When collection hours were examined for the hypericin content of plants, the highest content was recorded at 10:00 a.m (0.279%) and the lowest value at 4:00 p.m (0.272%) and 6:00 p.m (0.272%) (Table 4). Generally the higher hypericin contents were obtained from samples collected at 8:00 and 10:00 a.m compared to 12:00 (noon) and 6:00 p.m within a day (Table 4).

The accumulation level of hypericin may show variability (Pluhar et al., 2001). Differences between the studies can be explained by the different methods used, geographical and ecological factors, population variability, using herbarium or fresh plant materials and phases of plant collection and processing of the harvested plant material (Cellarova et al., 1994; Büter et al., 1998; Denke et al., 1999; Kitanov, 2000; Pluhar et al., 2001; Ayan et al., 2004). The size, number and chemical content of such glandular structures in plants can be influenced by a variety factors including nutrient availability (Mutikaninen and Walls, 1995; Guillet et al., 1997), light quality (Büter et al., 1998) and light intensity (Yamamaura et al., 1989; Upadhyaya and Furness, 1994; Guillet et al., 1997).

Martonfi and Repcak (1994) and Repcak and Martonfi (1997) pointed out that flower and its parts in medicinal and aromatic plants are the main organs that produced active principles in *H. perforatum* L. Therefore each treatment that changed the ratio of flower/herb affected active substances properties of St. John's Wort. Furthermore the extremely high hypericin content was possibly based on the higher proportion of flowers, leaves and fine stems (Pluhar et al., 2000). Only the 'flowering segment', is commonly harvested and processed for pharmaceutical use, since this segment contains the major part of the putative active compounds (Bomme, 1997; Southwell and Bourke, 2001; Pluhar et al., 2001).

The correct developmental stage, i.e., keeping the appropriate harvesting time, was imperative for the production of a *Hyperici* herba drug with a satisfying hypericin or hyperforin content (Büter and Büter, 2002). In earlier studies, different stages have been reported for the maximum hypericin concentrations. These stages were: the beginning of the blooming period (Kitanov, 1995), the budding and blossoming period (Kitanov, 2000), the massive flower bud formation period (Kireeva et al., 1999), the late budding stage (Franke et al., 1999), the period that flowers just opened (Tekelova et al., 2000), the period that flowers were at anthesis (Walker et al., 2001), the flowering period or shortly before (British, 1976; ESCOP, 1996), the full flowering period (Kartnig et al., 1997; Pluhar and Zelnik, 1994; Seidler et al., 1999), the stage that flowering had almost finished, when fruiting capsules were forming (Southwell and Bourke, 2001), the stage at which equal ratios of buds and capsules are present and when primary flower turns into a green capsule (Braunewell, 1991).

Our results demonstrate that in general, higher concentrations of hypericin were found in samples collected in Izmir and Edirne populations as compared to Bursa population. When these results are considered, it is possible to say that the highest hypericin ratio was determined in flowers and buds collected between 8:00 and 10:00 a.m within a day for examined *Hypericum* populations.

REFERENCES

- Ayan AK, Cırak C, Kevseroglu K, Özen T (2004). Hypericin in some *Hypericum* species from Turkey, Asian J. Plant Sci. 3(2): 200-202.
- Bombardelli E, Morazzoni P (1995). *Hypericum perforatum*. Fitoterapia. 66: 43-68.
- Bomme U (1997). Produktion stechnologie von Johanniskraut (*Hypericum perforatum* L.). Z. Arzn. Gew. pfl. 2: 127-134.
- Braunewell H (1991). Ökologische, Ontogenetische Einflüsse auf Ertrag und Inhaltsstoffgehalt von *Hypericum* ssp. Diss. Giessen 1991 Justus-Leibig-University.
- Briskin DP (2000). Medicinal plants and phytomedicines. Linking plant biochemistry and physiology to human health. Plant Physiol. 124: 507-514.

- British Herbal Pharmacopoeia (1976). *Hypericum*. Part One. pp. 177. Büter B, Orlicchio C, Soldati A, Berger K (1998). Significance of genetic and environmental aspects in the field cultivation of *Hypericum perforatum*. *Planta Med.* 64: 431-437.
- Büter KB, Büter B (2002). Ontogenetic variation regarding hypericin and hyperforin levels in four accessions of *H. perforatum* L. *J. Herbs, Spices Med. Plant.* 9(2/3): 95-100.
- Campbell MH, May CE, Southwell IA, Tomlinson JD, Michael PW (1992). Variation and varietal determination in *Hypericum perforatum* L. (St. John's wort) in New South Wales. *Plant Prot. Quart.* 7(2):43-45.
- Cellarova E, Daxnerova Z, Kimakova K, Haluskova J (1994). The variability of the hypericin contents in the regenerants of *Hypericum perforatum*. *Act. Biotechnol.* 14:267-274.
- Constantine GH, Karchesy J (1998). Variations in hypericin concentrations in *Hypericum perforatum* L. and commercial products. *Pharm. Biol.* 36(5): 365-367.
- Crompton CW, Hall IW, Jensen KIN, Hildebrand PD (1988). The biology of Canadian weeds. 83. *Hypericum perforatum* L. *Can. J. Plant Sci.* 68: 149-162.
- Curtis JD, Lersten NR (1991). Internal secretory structures in *Hypericum* (Clusiaceae): *H. perforatum* L. and *H. balearicum* L. *New Phytol.* 114: 571-580.
- DAC (1986). *Deutscher Arzneimittel-Codex 3. Ergänzung* (1991) *Johanniskraut- Hyperici Herba.* J- 010. Frankfurt am Main: Govi Verlag.
- Denke A, Schempp H, Mann E, Schneider W, Elstner EF (1999). Biochemical activities of extracts from *Hypericum perforatum* L. 4 th communication: Influence of different cultivation methods. *Arzneimittel-Forsch.* 49: 120-125.
- DeSmet PA, Mohen WA (1996). St. John's Wort as an antidepressant. *Brit. Med. J.* 313: 241-242.
- ESCOP (1996). *Monograph St. John's Wort.* European Scientific Cooperative on Phytotherapy Monographs on The Medical Uses of Plant Drugs. *Hyperici herba*, pp. 1-10.
- Fields P, Arnason GJT, Fulcher RG (1990). The spectral properties of *Hypericum perforatum* leaves: The implications for photoactivated defenses. *Can. J. Bot.* 68: 1166-1170.
- Fornasiero RB, Bianchi A, Pinetti A (1998). Anatomical and ultrastructural observations in *Hypericum perforatum* L. leaves. *J. Herbs Spices Med. Plants.* 5: 21-33.
- Franke R, Schenk R, Bauermann U (1999). Variability in *Hypericum perforatum* L. breeding lines. *Acta Hort.* 502: 167-173.
- Franz CH (1983). Nutrient and water management for medicinal and aromatic plants. *Acta Hort.* 132: 203-215.
- Guillet G, Lorenzetti F, Belanger A, Arnason JT, Bernays E (1997). Production of glands in leaves of *porophyllum* spp. (Asteraceae): ecological and genetic determinants and implications for insect herbivores. *J. Ecol.*, 85: 647-655.
- Jensen KIN, Gaul SO, Specht EG, Doohan DJ (1995). Hypericin content of Nova Scotia biotypes of *H. perforatum* L. *Can. J. Plant. Sci.*, 75: 923-926.
- Kartnig T, Heydel B, Lasser L (1997). St. John's Wort cultivated in Switzerland. *Agrarforschung.* 4(7): 299-302.
- Kireeva TB, Sharanov UL, Letchamo W (1999). Biochemical and eco-physiological studies on *Hypericum* spp., J. JANICK (Editor), *Perspectives on New Crops and New Uses.* ASHS Press, Alexandria, VA: pp. 467-468.
- Kitanov G (2000). The dynamics and content of hypericins in *Hypericum perforatum* L. and *Hypericum maculatum* Crantz growing in Bulgaria. *Acta Pharmaceutica-Zagreb.* 50(1): 65-68.
- Kitanov GM (1995). Hypericins in *Hypericum* species. *Marmara Üniv.Eczacılık Derg.* 11 (1-2): 343-350.
- Kordana S, Zalecki R (1997). Research on the cultivation of *Hypericum perforatum* L. *Hortic. Abstr.* 67(9): 801.
- Lavie G, Mazur Y, Lavie D, Meruelo D (1995). The chemical and biological properties of hypericin-acompound with a broad spectrum of biological activities. *Med. Res. Rev.*, 15: 111-119.
- Linde K, Ramirez G, Mulrow CD, Pauls A, Weiden Hammer W, Melchart D (1996). St. John's Wort for depression-an overview and meta-analysis of randomised clinical trials. *Brit. Med. J.* 313: 253-258.
- Martonfi P, Repcak M (1994). Secondary metabolites during flower ontogenesis of *Hypericum perforatum* L. *Zahradnictvi.* 21(1):37-44.
- Melikian E, Boroyan R, Karaguezian A, Charchoghlian A, Gabrielian E, Panossian A (1998). Hypericin content in St. John's Wort (*Hypericum perforatum* L.) growing in Armenia. *Pharm. Pharmacol. Lett.* 8(3): 01-102.
- Meruelo D, Lavie G, Lavie D (1988). Therapeutic agents with dramatic antiretroviral activity and little toxicity at effective doses: aromatic polycyclic diones hypericin and pseudohypericin. *Proc. Natl. Acad. Sci. U. S. A.* 85: 5230-5234.
- Mutikaninen P, Walls M (1995). Growth, reproduction and defense in nettles: Responses to herbivory modified by competition and fertilization. *Oecologia*, 104: 487-495.
- Osinska E, Weglarz Z (2001). Comparative study on three *Hypericum* species growing wild in Poland. *Proceedings of the International Conference of Medicinal and Aromatic Plants Possibilities and Limitations of Medicinal and Aromatic Plant Production in the 21st Century*, 8-11 July 2001, Budapest, Hungary, pp. 41-43.
- Palevitch D (1987). Recent advances in the cultivation of medicinal and aromatic plants. *Acta Hort.* 208: 29-35.
- Palevitch D (1991). *Agronomy Applied to Medicinal Plant Conservation*, In *The Conservation of Medicinal Plants.* Edited by Akerele O, Heywood V, Syngue H. Cambridge University Pres, Cambridge, U.K. pp. 167-178.
- Patocka J (2003). The chemistry, pharmacology and toxicology of the biologically active constituents of the herb *Hypericum perforatum* L. *J. Appl. Biomed.* 1: 61-70.
- Pluhar ZS, Bernath J, Neumayer E (2001). Morphological, production, biological and chemical diversity of St. John's Wort (*Hypericum perforatum* L.). *Proceedings of The International Conference on Medical and Aromatic Plants Possibilities and Limitations of Medicinal and Aromatic Plant Production in The 21st Century*, 8-11 July, Buda-pest, Hungary., pp. 33-40.
- Pluhar ZS, Rehak O, Nemeth E (2000). Comparative investigation on *Hypericum perforatum* L. populations of different origin. *Inter. J. Horticultural Sci.*, 6(1):56-60.
- Pluhar ZS, Zelnik K (1994). Introduction of *Hypericum perforatum* cv. Topas. *Proceedings of International Conference on Medicinal and Aromatic Plants, Trento, Italy, June 2-3*, pp. 628-630.
- Poutaraud A, Girardin P (2004). Agronomic and chemical characterization of 39 *Hypericum perforatum* accessions between 1998 and 2000. *Plant Breed.* 123(5): 480.
- Repcak M, Martonfi P (1997). The localization of secondary substances in *Hypericum perforatum* flower. *Biol. (Bratislava), Section Bot.* 52(1): 91-94.
- Schütt H (1996). *Morphologische, phytochemische und botanische Untersuchungen zur Selektion hypericin, pseudohypericin und flavonoidreicher Hypericum perforatum L. Stamme.* (Morphological, phytochemical and botanical studies on the selection of hypericin, pseudohypericin and flavonoid-rich *Hypericum perforatum* L. strains) E. Schweizerbart'sche Verlagsbuchhandlung, Berlin. *Diss. Bot. Band*, p. 263.
- Seidler LK, Dabrowska J, Zygmunt B (1999). Content of active substances in herb of St. John's Wort (*H. perforatum* L.) cv. Topas in different growth phases. *Herba-Polonica.* 45(3): 169-172.
- Sirvent TM, Walker L, Vance N, Gibson DM (2002). Variation in hypericins from wild populations of *H. perforatum* L. in the Pacific Northwest of the U.S.A. *Econ. Bot.* 56(1): 41-48.
- Southwell IA, Bourke CA (2001). Seasonal variation in hypericin content of *Hypericum perforatum* L. (St. John's Wort). *Phytochemistry*, 56: 437-441.
- Southwell IA, Campbell MH (1991). Hypericin content variation in *Hypericum perforatum* in Australia. *Phytochemistry*, 30(2): 475-478.
- Steel RGD, Torrie JH (1980). *Principles and procedures of statistics: A biometrical approach.* (McGraw-Hill Co. New York).
- Tekelova D, Repcak M, Zemkova E, Toth J (2000). Quantitative changes of dianthrones, hyperforin and flavonoids content in the flower ontogenesis of *Hypericum perforatum*. *Planta Med.* 66: 778-780.
- Upadhyaya MK, Furness NH (1994). Influence of light intensity and water stress on leaf surface characteristics of *cynoglossum officinale*, *Centaurea* spp. and *Tragopogon* spp. *Can. J. Bot.*, 72: 1379-1386.
- Upton R, Graff A, Williamson E, Bunting D, Gatherum DM, Walker EB,

- Butterweck V, Liefänder U, Nahrstedt A, Winterhoff H, Cott J (1997). St. John's Wort Monograph in: American Herbal Pharmacopoeia and Therapeutic Compendium. HerbalGram. 40: 1-32.
- Vlietinck AJ, De Bruyne T, Apers S, Pieters L (1988). Plant-derived leading compounds for chemotherapy of human immunodeficiency virus (HIV) infection. *Planta Med.* 64: 97-109.
- Walker L, Sirvent T, Gibson D, Vance N (2001). Regional differences in hypericin and pseudohypericin concentrations and five morphological traits among *Hypericum perforatum* plants in the Northwestern United States. *Can. J. Bot.* 79(10): 1248-1255.
- Yamamura T, Tanaka S, Tabata M (1989). Light-dependent formation of glandular trichomes and monoterpenes in thyme seedlings. *Phytochemistry.* 28: 741-744.