

Full Length Research Paper

Ability of drought selection indices to identify the best drought tolerant and high yielding genotypes in Mediterranean environment

Saad Rahab Eddie^{1, 2*}, Ahmed L. Hadje², Adel G. Batma³ and Mohammed Taarabt⁴

¹National Institute of Agricultural Research, BP 578, Meknès, Morocco.

²Hassan II Veterinary and Agronomy Institute, BP 6202, Rabat, Morocco.

³National Institute of Agricultural Research, BP 589, Settat, Morocco.

⁴University Mohammed VI Polytechnique, UM6P, Benguerir, Morocco.

Accepted 22 August, 2016

Drought is the most severe production constraint for wheat worldwide. Evaluating performance of bread wheat lines and predicting drought tolerance is an essential part of the breeding process. The objective of this study is to investigate the efficiency of several indices in identifying wheat genotypes combining drought tolerance and high yield potential. Twenty-four indices, which were most frequently used in plant breeding, were compared based on grain yield of 40 bread wheat genotypes grown under two contrasting environments (stressed and non-stressed) during 2 cropping seasons 2014 and 2015. The trials were laid out as completely randomized block design of 3 replicates. Experienced stress was moderate because it caused less than 50% reduction in yield in both seasons. Analysis of variance of grain yield showed significant differences among genotypes, years, sites and genotype × site interaction. All drought indices revealed significant differences among genotypes in both seasons, except GM, SNPI and ATI. Based on correlations and principal component analysis, repeatable strong positive correlations were found between the indices (MP, MRP, REI, GMP, STI, MSTIk1, MSTIk2, HM and RDY) and grain yield under both moisture conditions during the two seasons. These indices can be considered as suitable criteria for selection of drought tolerant and high yielding genotypes under moderate stress Mediterranean environment. Moreover, these indices were able to select the highest mean yields under 20% of selection pressure with low variation across environments; especially STI, GMP and MP. The genotypes “Gladius” (9) and “AUS30355” (11) were consistently selected in both environments during two cropping seasons.

Key words: *Triticum aestivum*, drought stress, tolerance indices, grain yield.

INTRODUCTION

In the Mediterranean region, climate change is associated with more frequent and intense periods of

drought as overall rainfall levels decline. The negative effect of drought stress on agriculture sector has been

*Corresponding author. E-mail: eddie.saad@yahoo.com

qualified as a major problem in many parts of the world (Nouraein et al., 2013; Passioura, 2007), limiting the expression of crops yield potential and stability, especially in dryland areas (40% of world surface) (Karamanos et al., 2012).

Wheat is one of the most important crops for food security worldwide (Bishaw et al., 2011; Travlos, 2012). In Morocco, bread wheat is a staple food grown under various environments and agro-ecosystems. It occupies 70% of cereal cultivated area (2 Million hectare) with an average production of 2.8 Million tons. It is usually cultivated as a rain-fed crop in regions characterized by irregular annual precipitations and/or unequal distribution of rainfalls within a season (Jlibene, 2009).

Drought is a major constraint decreasing yield and potential production. Plant growth and productivity are adversely affected by water stress leading to heavy yield losses. Besides the water scarcity status, the exploration of new ways for an efficient use of water input is primordial for food security and sustainable environment. Breeding is one of the most efficient options to overcome this complex stress through the development of new varieties adapted to drought and climate instability. However, the lack of accurate reproducible screening techniques limits the success of the breeding programs (Ramirez and Kelly, 1998; Farshadfar and Elyasi, 2012; Farshadfar et al., 2012a).

Despite the lack of understanding of the drought tolerance mechanisms, the grain yield remain the basis of genotypes selection for improving drought tolerance (Talebi et al., 2009; Shirinzadeh et al., 2010; Geravandi et al., 2011; Farshadfar et al., 2012a). Some researchers believe in selection based on only favorable conditions where the low magnitude genotype \times environment interaction permits to express the genetic potential yield (Richards, 1996; Rajaram and Van Ginkle, 2001; Betran et al., 2003); or only under stress conditions (Gavuzzi et al., 1997). However, high potential yield under non-stress conditions does not necessarily result in improved yield under stress conditions and genotypes with high yield may not be stress tolerant to drought and the reverse is true (Blum, 1996; Sio-Se Mardeh et al., 2006). Currently, many authors have chosen a mid-point and believe that selection considering yield under both non-stress and stress conditions is more efficient especially under unpredictable rain-fed conditions with various yearly drought scenarios (Mitra, 2001; Farshadfar et al., 2001; Moosavi et al., 2008; Mohammadi et al., 2010; Farshadfar et al., 2012a, b, 2014).

Thus, many drought indices have been proposed for screening drought tolerant genotypes based on yield under stressed and non-stressed environments (Mitra, 2001; Talebi et al., 2009; Pireivatlou et al., 2010; Mohammadi et al., 2010; Nouri et al., 2011) aiming at assisting the identification of stable, high yielding, drought tolerant genotypes: Stress susceptibility index (SSI) (Fischer and Maurer, 1978), drought response index

(DRI) (Bidinger et al., 1987), relative drought index (RDI) (Fischer and Wood, 1979), mean productivity (MP), tolerance index (TOL) (Rosielle and Hamblin, 1981), drought tolerance efficiency (DTE) (Fischer and Wood, 1981), yield stability index (YSI) (Bousslama and Schapaugh, 1984), superiority index (Pi) (Lin et al., 1986), geometric mean productivity (GMP), stress tolerance index (STI) (Fernandez, 1992), drought resistance index (DI) (Lan, 1998), mean relative performance (MRP), relative efficiency index (REI) (Hossain et al., 1999), relative adaptability to drought (bN) (Karamanos and Papatheohari, 1999), modified stress tolerance indices 1 and 2 (MSTIk) (Farshadfar and Sutka, 2002), abiotic tolerance index (ATI), stress susceptibility percentage index (SSPI), stress/non stress production index (SNPI) (Moosavi et al., 2008), harmonic mean of yield (HM) (Dadbakhch et al., 2011), sensitivity drought index (SDI) (Farshadfar and Javadinia, 2011), golden mean (GM) (Moradi et al., 2012) and relative decrease in yield (RDY) (Farshadfar and Elyasi, 2012). The best indices are those which have high correlation with grain yield in both conditions and would be able to identify potential upper yielding and drought tolerant genotypes (Fernandez, 1992; Mitra, 2001; Farshadfar et al., 2001; Boussen et al., 2010).

In this perspective, the objectives of the study were to (i) investigate the repeatable ability and efficiency of 24 drought selection indices to identify the best drought tolerant and high yielding genotypes adapted to both stressed and non-stressed conditions in a Mediterranean environment, (ii) study the inter-relationships among them and (iii) identify the genotypes adapted to stressed environments.

MATERIALS AND METHODS

Plant materials and experimental design

Forty bread wheat genotypes, originating from different breeding programs, ICARDA, CIMMYT, Australia and Morocco, were chosen for evaluation based on their presumed differences for yield performance under different moisture conditions (Table 1). Those genotypes were evaluated for grain yield, in two contrasting sites, representing stressed and non-stressed conditions, during two cropping seasons. The yield data were then used to derive 24 selection indices.

Each experiment was laid out in a completely randomized block design (RCBD) with three replications. Each plot is composed of 6 rows of 5 m; with row to row distance of 0.25 m. The sowing was done in late November and harvesting in mid- May for stressed and mid-July in non-stressed experimental sites. Seeding rate was 300 grains/m². Fertilizer application (18-46-00) was 1.5 quintal/ha at planting and 1 quintal/ha (Ammonitrate 33.5%) at tillering stage. The plants were protected against foliar diseases by fungicides, and weeds were controlled manually and by herbicides when needed. Yield (t/ha) was obtained based on 9 m² of harvested plot.

Experimental sites

Two experimental stations of the National Institute of Agricultural

Table 1. List of the 40 bread wheat genotypes.

Entry code	Name	Origine	Entry code	Name	Origin
1	NEJMAH-11	ICARDA	21	SB062	CIMMYT
2	NEJMAH-14	ICARDA	22	SB109	CIMMYT
3	SHIHAB-12	ICARDA	23	SB169	CIMMYT
4	AL-ZEHRAA-2	ICARDA	24	SsrT02	CIMMYT
5	BAASHA-21	ICARDA	25	SsrT09	CIMMYT
6	AMIR-2	ICARDA	26	SsrT14	CIMMYT
7	ATTILA	CIMMYT	27	SsrT16	CIMMYT
8	SOKOLL	CIMMYT	28	SsrT17	CIMMYT
9	GLADIUS	AUSTRALIA	29	SsrW35	CIMMYT
10	AUS30354	CIMMYT	30	SsrW47	CIMMYT
11	AUS30355	CIMMYT	31	ARREHANE	Morocco
12	AUS30518	CIMMYT	32	ACHTAR	Morocco
13	AUS30523	CIMMYT	33	MARCHOUCH	Morocco
14	QG-170-4.1	CIMMYT	34	KANZ	Morocco
15	QG-58-5.1	CIMMYT	35	AMAL	Morocco
16	HARTOG	AUSTRALIA	36	MASSIRA	Morocco
17	DRYSDALE	AUSTRALIA	37	AGUILAL	Morocco
18	SB003	CIMMYT	38	BT05A104	Morocco
19	SB165	CIMMYT	39	BT05A106	Morocco
20	SB069	CIMMYT	40	RAJAE	Morocco

Research of Morocco, namely Taoujdate and Sidi El Aidi, were used as sites for experimentation, for two cropping seasons 2013-2014 and 2014-2015. Taoujdate site (Fes province) represented the non-stressed or favorable environment (Altitude: 550 m, Latitude: 33°N, Longitude: 5°; long term average rainfall 500 mm; deep clayey soil (Tirs)); while Sidi El Aidi site (Settat province) represented the stressed semi-arid environment (Altitude 240 m, latitude 33°07'16", longitude 7°37'48"W; long term average rainfall 300 mm; limestone-clay texture soil). During the rest of the document, whenever indicated, an environment will be referred to as a combination of site by year.

Calculation of indices

Drought tolerance indices per cultivar "i" were calculated based on grain yield per plot for stress (Y_{si}), non-stress (Y_{pi}) environments and mean of grain yield under stressed (Y_s) and non-stress conditions (Y_p) as indicated in Table 2. In statistical basis, the efficiency of the drought indices will be evaluated based on their ability of discrimination between genotypes, correlation with grain yields of both environments and their efficiency to target the best high yielding and stable genotypes.

Statistical analysis

Data were subjected to analysis of variance for grain yield and drought indices using one-way ANOVA for data of each particular trial, two-way ANOVA for combined data across year, three-way ANOVA for combined data across site and year. For grain yield, the combined three way ANOVA was performed considering the effect of year, experimental site and genotype according to the model $Y = \text{year} + \text{site} + \text{bloc}(\text{site}) + \text{genotype} + \text{genotype by year} + \text{genotype by site} + \text{genotype by year by site} + \text{error}$. For drought indices, the combined two-way ANOVA was performed considering the effect

of year and genotype according to the model $Y = \text{year} + \text{genotype} + \text{bloc} + \text{genotype by year} + \text{error}$, while the one way ANOVA was used for each trial separately to detect the genotypic effect per year using the model $Y = \text{genotype} + \text{bloc} + \text{error}$. For each combined ANOVA, the magnitude of variation attributable to each factor was estimated as percentage of variance explained (VE %) of total sum of squares.

Ranks were assigned to genotypes for each index and simple correlation analysis using Spearman's coefficient was performed to elucidate the relationships among the selection indices for each cropping season, and their association with grain yield. Based on indices formula, the genotype with the highest value for Y_s , Y_p , MP, MRP, REI, GMP, STI, MSTI_{k1}, MSTI_{k2}, HM, YI, RDI, DI, GM, SNPI, DTE and DRI and the lowest value for SSI, TOL, Pi, SDI, SSPI, ATI, RDY, b and bN received a rank 1.

Principal component (PC) analysis method based on rank correlation matrix data was used to elucidate graphically the relationships among drought indices at once. The ANOVA was performed using GENSTAT (Discovery edition 3, VSN International, UK). The correlations and PC analysis were carried out using XLSTAT (Free trial version 2015, Addinsoft, Inc., Brooklyn, NY, USA).

RESULTS AND DISCUSSION

Pattern of the cropping seasons

For the stressed site (Sidi El Aidi), the rainfall amount was about 181 and 237 mm respectively in 2014 and 2015 cropping seasons. In non-stressed conditions (Taoujdate), the rainfall amount was about 278 and 412 mm during the respective seasons 2014 and 2015. Additional irrigation (about 100 mm) was applied during

Table 2. List of the 24 drought tolerance indices and formula.

Index	Abbr.	Formula	References
Mean productivity	MP	$(Y_{pi} + Y_{si}) / 2$	Rosielle and Hamblin, 1981
Mean relative performance	MRP	$(Y_{si} / Y_s) + (Y_{pi} / Y_p)$	Hossain et al., 1999
Stress susceptibility index	SSI	$1 - (Y_{si} / Y_{pi}) / SI$ Where Stress intensity (SI) = $1 - (Y_s / Y_p)$	Fischer and Maurer, 1978
Stress tolerance index	TOL	$Y_{pi} - Y_{si}$	Rosielle and Hamblin, 1981
Geometric mean productivity	GMP	$\sqrt{Y_{pi} \times Y_{si}}$	Fernandez, 1992
Relative efficiency index	REI	$(Y_{si} / Y_s) * (Y_{pi} / Y_p)$	Hossain et al., 1999
Stress tolerance index	STI	$(Y_{si} \times Y_{pi}) / (Y_p)^2$	Fernandez, 1992
Modified stress tolerance index 1	MSTik1	$(Y_{pi})^2 / (Y_p)^2 \times STI$	Farshadfar and Sutka, 2002
Modified stress tolerance index 2	MSTik2	$(Y_{si})^2 / (Y_s)^2 \times STI$	Farshadfar and Sutka, 2002
Harmonic mean of yield	HM	$2 \times (Y_{pi} \times Y_{si}) / (Y_{pi} + Y_{si})$	Dadbakhsh et al., 2011
Coefficient of regression	b	$\Sigma Y_{ij} Y_j / \Sigma Y^2$ where i refers to genotypes and j to environment; Y is the overall mean of all genotypes in both environments.	Bansal and Sinha, 1991
Relative adaptability to drought	bN	b / a ; where b = Slope of regression model; a = intercept of regression model	Karamanos and Papatheohari, 1999
Yield Index	YI	Y_{si} / Y_s	Gavuzzi et al., 1997; Lin et al., 1986
Superiority Index	Pi	$\Sigma_{j=1}^n (X_{ij} - M_j)^2 / 4$; where X_{ij} = Grain yield of the ith genotype in the jth environment, M = Yield of the highest yielding genotype in the environment j	Clarke et al., 1992; Lin et al., 1986
Sensitivity drought index	SDI	$(Y_{pi} - Y_{si}) / Y_{pi}$	Farshadfar and Javadinia, 2011
Relative drought index	RDI	$(Y_{si} / Y_{pi}) / (Y_s / Y_p)$	Fischer and Wood, 1979
Drought resistance index	DI	$Y_{si} \times (Y_{si} / Y_{pi}) / (Y_s)$	Lan, 1998
Golden mean	GM	$(Y_{pi} + Y_{si}) / (Y_{pi} - Y_{si})$	Moradi et al., 2012
Abiotic tolerance index	ATI	$((Y_{pi} - Y_{si}) / (Y_p / Y_s)) * (\sqrt{Y_{pi}} * Y_{si})$	Moosavi et al., 2008
Stress Susceptibility percentage index	SSPI	$((Y_{pi} - Y_{si}) / (2 * Y_p)) * 100$	Moosavi et al., 2008
Stress/non-stress production index	SNPI	$(\sqrt[3]{(Y_{pi} + Y_{si})} / (Y_{pi} - Y_{si})) * \sqrt[3]{Y_{pi}} * Y_{si} * Y_{si}$	Moosavi et al., 2008
Drought response index	DRI	$(Y_A - Y_{ES}) / S_{ES}$; where Y_A = Yield estimate by regression in stress conditions; Y_{ES} = Real yield in stress conditions; S_{ES} = Standard error of estimated grain yield of all genotypes	Bidinger et al., 1987
Relative decrease in yield	RDY	$100 - ((Y_{si} / 100) * Y_{pi})$	Farshadfar and Elyasi, 2012
Drought tolerance efficiency	DTE	$(Y_{si} / Y_{pi}) * 100$	Fischer and Wood, 1981

Y_{si} : Yield under stress for genotype "i"; Y_{pi} : Yield under non-stress for genotype "i"; Y_s : Mean of grain yield under stressed; Y_p : Mean of grain yield under non-stress conditions.

critical growing stages. The drought stress occurred essentially at mid-cycle during the reproductive stage (pre-flowering and flowering) (Figure 1).

In non-stressed environment, the mean grain yield was higher during 2015 (4.49 t/ha) compared to 2014 (3.35 t/ha). However, the mean yield was 1.93 t/ha in 2014 compared to 3.05 t/ha during 2015 under stress conditions. During both seasons, the grain yield of genotypes showed greater variation under non stress compared to stress conditions. This variation can be explained by the differences in genotypes response to

different moisture conditions (Mohammadi et al., 2010). Stress intensity in the first and second cropping season was respectively 0.43 (43% of yield reduction) and 0.32 (32% of reduction). Thus, the drought intensity was moderate for both seasons (below 50%). However, this index evaluates only drought stress intensity of the whole experiment and not for different genotypes.

Analysis of variance

Based on combined ANOVA, statistically significant




Figure 1. Rainfall amount (mm) in non-stressed (NS) and stressed (S) sites during the two cropping seasons 2013-14 and 2014-15. The oval forms refer to the timing and extension of the flowering stage.

Table 3. Combined analysis of variance for grain yield of 40 genotypes.

Source of variation	df	M.S	Percentage of variation explained (%)
Block	2	10.82	2.59
Year	1	146.68**	35.14
Site	1	255.73**	61.26
Genotype	39	1.63**	0.39
Year.Genotype	39	0.60	0.14
Year. Site	1	0.12	0.03
Site.Genotype	39	0.80*	0.19
Year. Site.Genotype	39	0.60	0.14
Residual	296(22)	0.48	0.11

*, ** Significant at 5 and 1% levels, respectively; † df: degree of freedom; M.S: mean square

differences for grain yield were found among years, sites, genotypes and for genotype × site interaction. However, the other interactions year × genotype, year × site and year × site × genotype were not significant (Table 3). The magnitude of variation attributable to the years, sites and genotypes was respectively 35, 61 and 0.4% (Table 3). These results indicated that the genotypes represented a broad range of response to drought stress based on its intensity influenced by the environmental variations (Mohammadi et al., 2011; Farshadfar et al., 2012a).

The results of combined analysis of variance of selection indices are presented in Table 4. Significant differences were observed between years for ATI, DTE, GMP, HM, MP, Pi, RDY, SDI, SNPI, SSPI, STI and bN. Those indices were influenced mainly by year effect as confirmed by the percentage of explained variance per factor (Table 4); whereas, the indices DRI, MRP, MSTIk1, MSTIk2, RDI, REI, SSI, TOL and YI showed an important genotypic variation compared to the year effect (Table 4). The interaction genotype × year was significant only for SSI, Pi, ATI and RDY. Thus, those indices ranked

differently the genotypes depending on the variation of stress intensity between years (Table 4).

All drought indices showed significant differences among genotypes except SNPI, b and bN during both seasons, GM which discriminated between genotypes only in 2015 and ATI which discriminated between genotypes only in 2014 season. Those results demonstrated that almost all indices revealed an important genetic diversity and were able to discriminate between the genotypes. However, the efficient indices should be also able to select the genotypes combining high yield and drought tolerance (Mitra, 2001; Farshadfar et al., 2001).

Correlation analysis

Correlation analysis revealed a positive but non-significant association between grain yield of stressed and non-stressed conditions of 0.16 and 0.3 during 2014 and 2015 seasons respectively. Correlation seemed to

Table 4. Mean Square of analysis of variance of drought tolerance indices for the 40 genotypes over and within each season 2013-14 and 2014-15.

Source of variation	Two way ANOVA					ANOVA 2014		ANOVA 2015
	Year (Y)	VE (%)	Genotype (G)	VE (%)	Y x G	VE (%)	Genotype	Genotype
MP	70.87**	94.43	0.82**	0.10	0.36	0.04	0.36**	0.83**
MRP	0.05	1.87	0.38**	14.44	0.15	5.60	0.26**	0.27**
SSI	0.23	7.77	1.10**	36.17	0.76*	24.98	7.83**	12.41*
TOL	0.007	0.02	1.63**	36.07	1.12	24.66	0.96**	1.52*
GMP	74.33**	93.55	1.03**	1.30	0.47	0.06	0.49**	1.01**
REI	0.06	0.24	0.39**	16.25	0.14	0.57	0.26**	0.27**
STI	0.47**	37.54	0.15**	12.23	0.06	0.45	0.086**	0.13**
MSTik1	0.43	15.59	0.45**	16.30	0.16	5.89	0.27**	0.36**
MSTik2	0.0003	0.00	1.24**	18.79	0.45	6.88	0.68*	1.01**
HM	76.16**	92.77	1.25**	1.52	0.59	0.07	0.62**	1.19**
YI	0.02	0.14	0.24**	16.33	0.11	0.70	0.17**	0.16**
Pi	25.57**	75.57	1.73**	0.51	1.16*	0.34	0.59**	2.31**
ATI	72.32**	71.70	5.23**	5.20	4.69*	4.60	2.46**	6.95
DI	0.49	22.10	0.37**	16.90	0.21	9.60	0.19*	0.32**
DRI	0.16	2.90	2.03**	27.00	0.88	11.70	1.24*	1.38**
DTE	8815**	66.70	1352.5**	10.23	879.8	6.65	652.9**	1311.5*
GM	39369	39.64	18920	19.05	19123	19.26	37596	419.5*
RDI	0.021	0.15	0.34**	24.70	0.21	15.71	0.199**	0.28*
RDY	0.29**	95.04	0.004**	0.14	0.002*	0.07	652.9**	1311.5*
SDI	0.88**	66.69	0.14**	10.23	0.09	0.66	0.066**	0.13*
SNPI	499.8**	77.02	29.58	0.45	34.25	0.53	30.54	32.54
SSPI	1656**	64.95	263.1**	10.23	177.0	0.69	212.6**	188.4*
b	0.01	-	1.49	-	-	-	-	-
bN	1.2*	-	0.23	-	-	-	-	-

*, ** Significant at 5 and 1% levels, respectively; VE (%): Percentage of variation explained.

have improved from the dry season of 2014 to the wet season of 2015 where the stressed site was 41 mm wetter than the non-stressed site of 2014. Similar findings were reported by Fernandez (1992), Clarke et al. (1992), Sio-Se Mardeh et al. (2006), Mohammadi et al. (2010), Boussen et al. (2010), Nouri et al. (2011) Dadbakhsh et al. (2011) and Farshadfar et al. (2013) suggesting that high yield under non stress condition will not result necessarily in improved yield under stress conditions (and the opposite is true) because the genes controlling yield and drought resistance/tolerance are different (Rosielle and Hamblin, 1981; Golabadi et al., 2006; Anwar et al., 2011). Thus, under such conditions, the indices correlated with both moisture conditions are the most suitable to select stable genotypes with good yield performances (Mitra, 2001; Farshadfar et al., 2001; Farshadfar et al., 2012a, b; 2013; 2014).

Correlation between grain yield and drought tolerance indices

To determine the most desirable drought tolerance

criteria, the Spearman coefficient of correlation (based on ranks) between grain yield in both moisture conditions and each of the drought indices were calculated for 2013-14 (Table 5) and 2014-15 seasons (Table 6).

During 2014 (under 43% of stress intensity) (Table 5), yield under stress condition (Ys) was highly significantly and positively correlated with the indices MP, MRP, REI, SSI, TOL, GMP, STI, MSTik1, MSTik2, HM, YI, SDI, RDI, DI, SSPI, ATI, RDY, DTE and DRI; and moderately correlated with the coefficient of regression b. When the stress became less severe (32%) during 2015 (Table 6), the latter correlations remained the same but became stronger, except for ATI ($r=0.56$ in 2015 besides 0.69 in 2014). Moreover, other relationships appeared between Ys and (GM, SNPI and bN) respectively. Based on these results over both cropping seasons, significant positive repeatable correlations were found between yield under stress conditions (YS) and the drought indices (MP, MRP, REI, SSI, TOL, GMP, STI, MSTik2, HM, YI, SDI, RDI, DI, SSPI, RDY, DTE, DRI; ATI, MSTik1 and b). These relationships were influenced by the drought intensity (difference between Ys and Yp) and indicated that genotypes selected based on these indices are

Table 5. Spearman's rank correlation between grain yield and drought indices in 2014.

	YS	YP	MP	MRP	REI	SSI	TOL	GMP	STI	MSTik1	MSTik2	HM	YI	
YS	1													
YP	0.16	1												
MP	0.76	0.74	1											
MRP	0.85	0.62	0.98	1										
REI	0.86	0.60	0.97	0.99	1									
SSI	0.78	-0.43	0.23	0.39	0.40	1								
TOL	0.74	0.4	0.23	0.36	0.37	0.94	1							
GMP	0.81	0.64	0.97	0.97	0.97	0.33	0.38	1						
STI	0.81	0.64	0.97	0.97	0.97	0.33	0.38	1.00	1					
MSTik1	0.59	0.87	0.97	0.91	0.90	0.02	0.04	0.92	0.92	1				
MSTik2	0.91	0.39	0.87	0.91	0.91	0.56	0.63	0.94	0.94	0.76	1			
HM	0.82	0.61	0.96	0.96	0.96	0.35	0.41	0.99	0.99	0.90	0.95	1		
YI	0.94	0.23	0.78	0.84	0.84	0.68	0.75	0.87	0.87	0.64	0.98	0.88	1	
Pi	0.15	0.30	0.27	0.26	0.24	-0.08	-0.06	0.22	0.22	0.26	0.18	0.22	0.13	
SDI	0.79	-0.33	0.30	0.43	0.44	0.94	0.99	0.44	0.44	0.11	0.68	0.47	0.80	
RDI	0.79	-0.33	0.30	0.43	0.44	0.94	0.99	0.44	0.44	0.11	0.68	0.47	0.80	
DI	0.90	-0.05	0.56	0.66	0.66	0.85	0.91	0.68	0.68	0.39	0.87	0.70	0.94	
SSPI	0.74	-0.41	0.23	0.36	0.37	0.94	1.00	0.38	0.38	0.04	0.62	0.40	0.75	
GM	-0.05	0.16	0.08	0.11	0.13	-0.15	-0.14	0.09	0.09	0.12	-0.01	0.11	-0.07	
ATI	0.69	-0.48	0.15	0.29	0.30	0.93	0.99	0.30	0.30	-0.05	0.56	0.33	0.70	
SNPI	0.01	0.35	0.25	0.24	0.27	-0.22	-0.21	0.23	0.23	0.30	0.09	0.24	0.01	
RDY	0.86	0.60	0.97	0.99	1.00	0.40	0.37	0.97	0.97	0.90	0.91	0.96	0.84	
DTE	0.79	-0.33	0.30	0.43	0.44	0.94	0.99	0.44	0.44	0.11	0.68	0.47	0.80	
b	0.50	-0.41	0.06	0.19	0.20	0.71	0.65	0.11	0.11	-0.09	0.28	0.12	0.39	
bN	-0.24	0.11	-0.07	-0.13	-0.11	-0.29	-0.30	-0.09	-0.09	-0.02	-0.16	-0.09	-0.21	
DRI	0.94	0.22	0.76	0.83	0.83	0.70	0.76	0.86	0.86	0.63	0.97	0.87	0.99	
	Pi	SDI	RDI	DI	SSPI	GM	ATI	SNPI	RDY	DTE	b	S ²	bN	DRI
Pi	1													
SDI	-0.05	1												
RDI	-0.05	1.00	1											
DI	0.03	0.94	0.94	1										
SSPI	-0.06	0.99	0.99	0.91	1									
GM	0.17	-0.13	-0.13	-0.17	-0.13	1								
ATI	-0.08	0.97	0.97	0.86	0.99	-0.15	1							
SNPI	0.20	-0.17	-0.17	-0.14	-0.21	0.95	-0.25	1						
RDY	0.24	0.44	0.44	0.66	0.37	0.13	0.30	0.26	1					
DTE	-0.05	1.00	1.00	0.94	0.99	-0.13	0.97	-0.17	0.44	1				
b	-0.11	0.64	0.64	0.55	0.65	-0.12	0.64	-0.15	0.20	0.64	1			
bN	-0.02	-0.31	-0.31	-0.26	-0.30	0.01	-0.31	0.02	-0.11	-0.31	-0.53	0.26	1	
DRI	0.12	0.81	0.81	0.95	0.76	-0.07	0.71	0.00	0.83	0.81	0.39	0.33	-0.22	1

characterized by drought tolerance criteria and will improve yield under stress conditions.

These observed relationships are in consistence with numerous studies. Many studies reported positive relationships between Ys and the most popular and widely used indices MP, GMP, STI, SSI, TOL (Khalili et al., 2004; Golabadi et al., 2006; Gholinezadeh et al., 2010, Mohammadi et al., 2010, Farshadfar et al., 2012a,

Mevlut et Sait, 2011; Nouri et al., 2011; Mevlut and Sait, 2011; İlker et al., 2011; Reza Eivazi et al., 2013; Rahmani et al., 2013). Jafari et al. (2009); Gholinezadeh et al. (2010), Farshadfar and Eiyasi (2012) and Farshadfar et al. (2012b, 2013, 2014) noticed also positive significant correlation between YS and HM, YI, DI, MSTik1, MSTik2 and DRI. The coefficient of regression (b) expressed significant positive correlation with yield under stress to

Table 6. Spearman's rank correlation coefficients between yields and drought tolerance indices 2015.

	YS	YP	MP	MRP	REI	SSI	TOL	GMP	STI	MSTik1	MSTik2	HM	YI	
YS	1													
YP	0.31	1												
MP	0.86	0.71	1											
MRP	0.93	0.59	0.98	1										
REI	0.93	0.58	0.98	0.99	1									
SSI	0.91	-0.07	0.62	0.74	0.74	1								
TOL	0.80	-0.27	0.44	0.57	0.58	0.97	1							
GMP	0.93	0.58	0.98	0.99	1.00	0.74	0.58	1						
STI	0.93	0.58	0.98	0.99	1.00	0.74	0.58	1.00	1					
MSTik1	0.72	0.85	0.96	0.91	0.91	0.42	0.23	0.91	0.91	1				
MSTik2	0.99	0.40	0.91	0.97	0.97	0.87	0.74	0.97	0.97	0.79	1			
HM	0.96	0.49	0.95	0.99	0.99	0.81	0.67	0.99	0.99	0.85	0.99	1		
YI	1.00	0.31	0.86	0.93	0.93	0.91	0.80	0.93	0.93	0.72	0.99	0.96	1	
Pi	-0.04	0.10	0.03	0.04	0.03	-0.12	-0.17	0.03	0.03	0.09	-0.04	-0.02	-0.04	
SDI	0.91	-0.07	0.62	0.74	0.74	1.00	0.97	0.74	0.74	0.42	0.87	0.81	0.91	
RDI	0.91	-0.07	0.62	0.74	0.74	1.00	0.97	0.74	0.74	0.42	0.87	0.81	0.91	
DI	0.97	0.12	0.75	0.85	0.85	0.97	0.90	0.85	0.85	0.57	0.94	0.90	0.98	
SSPI	0.80	-0.27	0.44	0.57	0.58	0.97	1.00	0.58	0.58	0.23	0.74	0.67	0.80	
GM	0.80	0.08	0.66	0.72	0.75	0.85	0.82	0.75	0.75	0.53	0.79	0.80	0.80	
ATI	0.56	-0.55	0.13	0.27	0.28	0.83	0.93	0.28	0.28	-0.09	0.47	0.39	0.56	
SNPI	0.87	0.31	0.81	0.85	0.87	0.81	0.73	0.87	0.87	0.71	0.88	0.90	0.87	
RDY	0.93	0.58	0.98	0.99	1.00	0.74	0.58	1.00	1.00	0.91	0.97	0.99	0.93	
DTE	0.91	-0.07	0.62	0.74	0.74	1.00	0.97	0.74	0.74	0.42	0.87	0.81	0.91	
b	0.79	-0.26	0.43	0.56	0.57	0.96	0.99	0.57	0.57	0.22	0.72	0.65	0.79	
bN	0.91	-0.05	0.63	0.74	0.74	0.99	0.96	0.74	0.74	0.43	0.86	0.80	0.91	
DRI	0.95	0.04	0.69	0.79	0.80	0.97	0.92	0.80	0.80	0.50	0.91	0.86	0.95	
	Pi	SDI	RDI	DI	SSPI	GM	ATI	SNPI	RDY	DTE	b	S ²	bN	DRI
Pi	1													
SDI	-0.12	1												
RDI	-0.12	1.00	1											
DI	-0.09	0.97	0.97	1										
SSPI	-0.17	0.97	0.97	0.90	1									
GM	-0.17	0.85	0.85	0.83	0.82	1								
ATI	-0.21	0.83	0.83	0.71	0.93	0.69	1							
SNPI	-0.12	0.81	0.81	0.85	0.73	0.96	0.52	1						
RDY	0.03	0.74	0.74	0.85	0.58	0.75	0.28	0.87	1					
DTE	-0.12	1.00	1.00	0.97	0.97	0.85	0.83	0.81	0.74	1				
b	-0.22	0.96	0.96	0.89	0.99	0.81	0.93	0.72	0.57	0.96	1			
bN	-0.17	0.99	0.99	0.96	0.96	0.84	0.82	0.81	0.74	0.99	0.96	0.18	1	
DRI	-0.15	0.97	0.97	0.98	0.92	0.83	0.75	0.82	0.80	0.97	0.90	0.11	0.96	1

Bold values are significant at 5% level of probability.

identify the drought tolerant genotypes in Guttieri et al. (2001), Clarke et al. (1992), Ahmadi et al. (2004), Moghaddam and Hadizadeh (2002), Mevlut and Sait (2011) and Khadarahmpour et al. (2011). Gholinezadeh et al. (2010) and Mohammadi et al. (2012) reported also significant positive correlation between RDI and Ys.

Significant relationships between YS and REI, MRP and between Ys and DTE were also reported by Singh et al. (2011) and Kumar et al. (2014). Naghavi et al. (2013) observed significant differences between Ys and SSPI. This finding was in agreement with Naghavi et al. (2013) and in contradiction with Farshadfar et al. (2012a, 2014);

Moosavi et al., 2008).

The correlation between YS and SNPI disappeared when the stress reached 43% of intensity. However, many studies confirmed this significant positive correlation (Farshadfar et al., 2012a; 2012b; 2014; Moosavi et al., 2008). The same pattern was observed between Ys and GM. Mohammadi et al. (2011, 2012) found a significant positive correlation between YS and GM at 22.6 and 26.4% stress intensity. On the other hand, the relationships between YS and SDI, RDY disagreed with the findings of Farshadfar et al. (2012b). The positive correlation between Ys and ATI was in disagreement with the results of Farshadfar et al. (2012a, 2012b); Moosavi et al. (2008) which attested the absence of relationship between those two indices.

Under non-stressed environment, yield (Yp) during 2014 season (43% of stress intensity) was highly and positively correlated with MP, MRP, GMP, STI, MSTI1, HM, RDY and REI; moderately correlated with MSTI2 (0.39) (Table 5). During 2015 (32% of stress intensity), the same correlations were obtained with more moderate association (<0.4), except for MSTI2 and RDY (which remained almost the same) and ATI (stronger correlation with YF) (Table 6). Moreover, the moderate positive correlation with TOL, SNPI and the negative ones with SSI, SDI, RDI, SSPI, DTE and b were lost (Table 6). Based on the two cropping season results, the yield under favorable conditions (Yp) had strong positive repeatable correlation with MP, MRP, REI, GMP, STI, MSTI1, HM, RDY; moderate correlation with MSTI2 and significant negative correlation with ATI. Those indices permit to select genotypes with high yield potential (Yp). They are also influenced by the variation between yield under stressed and non-stressed conditions except RDY and MSTI2.

The positive correlations observed between Yp and (MSTI1, MP, GMP, STI, MSTI2, HM and RDY) are in agreement with the results obtained by Moosavi et al. (2008), Gholinezhad et al. (2014); Farshadfar et al. (2012b; 2013); Mevlut and Sait (2011), Farshadfar and Elyasi (2012) and Naghavi et al. (2013). However, the absence or the negative correlation between Yp and SSI, TOL, SSPI, SNPI, YI and DI are in disagreement with the same authors. Non-significant correlations between SSI and Yp were found in Ahmadi et al. (2004), Golabadi et al. (2006), Moosavi et al. (2008), Khodarahmpour et al. (2011), Drikvand et al. (2012), Mohammadi et al. (2012), Farshadfar et al. (2014) and Mohammadi et al. (2011). The presence of negative correlation between TOL and Yp under moderately severe conditions (60%) in comparison with moderate conditions (34%) was also found by Mohammadi et al. (2011) and the absence of correlation was reported in other studies (Moosavi et al., 2008; Khodarahmpour et al., 2011). Moosavi et al. (2008), Farshadfar et al. (2012b) and Naghavi et al. (2013) obtained positive correlation between Yp, SSPI and SNPI. However, the absence of correlation between

SSPI and Yp found in our study was in agreement with Farshadfar and Elyasi (2012). Moreover, the non-significant correlation between Yp and DI was in agreement with the results obtained by Farshadfar et al. (2012b). Similar to our findings, REI and MRP were useful in identifying genotypes with high yield potential in Singh et al. (2011). Furthermore, no correlation were found between (Pi, b, bN, YSI) and Yp as supported by Mohammadi et al. (2011).

In this study, no significant associations were found between Pi and the yield under both conditions. The same finding was observed in Mevlut and Sait (2011), however, this is not in agreement with other studies (Mohammadi et al., 2010; Mevlut and Sait, 2011). Moreover, our findings were in disagreement with the results obtained by Mohammadi et al. (2012) concerning the correlation between Yp and GM, RDI, YSI and DRI. In addition, the negative correlation between ATI and Yp is in disagreement with Moosavi et al. (2008) and Farshadfar et al. (2012, 2012a); Rahmani et al. (2013) where there was a positive association.

Overall, under moderate stress, the drought indices MP, MRP, REI, GMP, STI, MSTI1, MSTI2, HM and RDY were correlated with both moisture conditions (non-stressed and stressed) during the two cropping seasons. Thus, they can be considered as repeatable suitable criteria for selection for drought tolerant and high yielding genotypes. These results can be supported by numerous studies (Golabadi et al., 2006; Boussem et al., 2010; Nouraein et al., 2013; Farshadfar and Sutka, 2002; Ilker et al., 2011; Jafari et al., 2009; Mohammadi et al., 2003; 2010; 2011; Khodarahmpour et al., 2011; Farshadfar and Elyasi, 2012; Farshadfar et al., 2012a, b; 2013; Drikvand et al., 2012; Naghavi et al., 2013).

Relationships between drought tolerance indices

The relationships between the different drought indices will allow us to suggest one as alternative for the others that belong to the same group based on their strong correlation for the evaluation of the drought tolerant and high yielding genotypes. In the presence of a large number of indices, the principal component analysis (PCA) was used to assess in a simple graphic the relationships, similarities and dissimilarities between all attributes at once, based on the rank correlation.

In 2014 cropping season, the first and second components explained 80% of the total variation (55.5 and 24.2% respectively) (Figure 2). The PCA1 and PCA2 mainly distinguish the indices in different groups. The yield under stress (YS) and the indices MP, MRP, REI, SSI, TOL, GMP, STI, MSTI2, HM, YI, SDI, RDI, DI, SSPI, ATI, RDY, DTE and DRI were correlated with the first component. This component can be called "stress tolerance component". The cosine of the angle between the vectors of two indices approximates the correlation




Figure 2. Biplot of drought indices based on principal component analysis for 2013-14 season.

between them. The angle between MSTIK1, MSTIK2, MP, MRP, RDY, HM, STI, GMP, REI, DRI, YI and DI was well below 90° (acute angle) showing high correlations and similarities in ranking the genotypes ($0.64 < \text{coefficient of correlation } (r) < 0.99$); except for DI which showed moderate correlations. Inside this group, an overlapping of vectors (zero angle) was found between RDY, HM, STI, GMP, MRP, and REI ($0.96 < r < 1$) and between YI and DRI ($r = 0.999$) showing same ranking genotypes basis. Similar relationships were observed between SSI, ATI, DTE, TOL, SDI, SSPI and RDI ($0.93 < r < 1$) indicating that these indices are identical in genotype rankings. The second component PCA2 was highly positively correlated with the potential yield (Yp) and MSTIK1 ($r = 0.87$) and moderately correlated to the coefficient of regression (b) ($r = 0.41$). This component can be called “yield potential component”. The index MSTIK1 had positive and strong correlation to both components but with more emphasis on the stress tolerance component. The indices GM and SNPI were correlated to PCA3 whereas bN was correlated to PCA4; those two components explained only 6.5 and 5% respectively of variation between the indices. Finally, Pi was correlated to PCA5 which explained 3.45% of variation. Thus, the indices (GM, SNPI, bN and Pi) had low contribution to the variation between genotypes.

In 2015 cropping season, the first and second components explained 90.3% of the variation between all indices in 2015 (Figure 3). YS, MP, MRP, REI, SSI, TOL, GMP, STI, MSTIK2, HM, YI, SDI, RDI, DI, SSPI, GM, SNPI, RDY, DTE, DRI, bN were positively correlated with PCA1 (73% of variation), whilst b had negative correlation. The angle between MSTIK1, MSTIK2, MP, MRP, RDY, STI, GMP, REI, HM, YI and SNPI was well below 90 degrees showing similarities in ranking the genotypes ($0.83 < r < 1$). An acute angle was observed between MRP, GMP, RDY, STI and REI ($0.997 < r < 1$) displaying that these indices are identical in genotype rankings. An acute angle was also found between YS, SNPI and YI ($0.87 < r < 1$). Similar relationships were observed between TOL, SSPI, bN, DTE, RDI, SDI, SSI, GM, DI and DRI with an angle below 90° ($0.81 < r < 1$). A zero angle was found between SSI, SDI, bN, DTE and RDI ($0.99 < r < 1$). A zero angle was found between TOL and SSPI ($r = 1$), indicating that they ranked similarly the genotypes, as indicated by the zero angle between their vectors. Similar relationships were observed GM, DI and DRI ($0.82 < r < 0.98$). The PCA2 (17% of variation) is correlated positively with Yp, MSTIK1 and negatively with ATI. The grain yield (Yp) was highly correlated to MSTIK1 ($r = 0.85$) and moderately negatively correlated to ATI ($r = -0.55$). Similarly to the first year, the index MSTIK1 had




Figure 3. Biplot of drought indices based on principal component analysis for 2014-15 season.

positive and strong correlation to both components but with more emphasis on the first one. Finally, Pi had significant moderate correlation with PCA3 which explained only 4% of total variability.

Based on the results obtained in two cropping seasons, no relationship was found between the grain yield in non-stressed and stressed environments as indicated by their correlation to different components and by the right angle between their vectors. Strong repeatable significant correlations were found between MRP, RDY, STI, GMP, REI, HM, MP, YI, MSTik1 and MSTik2. This indicated that one of these indices could be used interchangeably as an alternative for the others in genotypes selection, especially the first 5 ones. The observed relationships are in concordance with those observed by Normand et al. (2001), Golabadi et al. (2006), Melvut and Sait (2011), Mohammed et al. (2011, 2012), Drikvand et al. (2012), Moradi et al. (2012), Farshadfar et al. (2012a, b), Reza Eivazi et al. (2013), Rahmani et al. (2013), Naghavi et al. (2013) and Farshadfar et al. (2014). Strong repeatable relationships were also observed between SSI, TOL, SDI, RDI, DTE and SSPI. Similar findings were reported by Normand et al. (2001), Golabadi et al. (2006), Bousset et

al. (2010), Mohammadi et al. (2010, 2012), Dadbakhsh et al. (2011), Farshadfar et al. (2012a, b), Rahmani et al. (2013) and Farshadfar et al. (2014). No correlation was found between the superiority index (Pi) and the other drought indices. Those findings were not in agreement with the findings of Mohammadi et al. (2010) and Melvut and Sait (2011).

Screening genotypes

Tables 7 and 8 represent the average mean yield, variance between genotypes and mean variance of the top 20% genotypes selection based on each index ranking during the two cropping seasons. Those parameters will be able to define the drought indices providing the best accurate genotypes selection at 20% selection pressure.

During the first cropping season 2014 (Table 7), mean values of indices showed similarities in top ranking genotypes for MP, MRP, REI, GMP, MSTik1, HM, RDY and STI. The top similar genotypes for this group were 2, 6, 9, 10 and 11. The indices DI, SSI, TOL, SDI, RDI,

Table 7. Descriptive statistics of the top 20% genotype selection for drought indices during 2013-2014.

Indices	Mean yield of 20% top genotypes	Variance inter 20% genotypes	Mean variance of the top genotypes	Top 20% genotypes selected
ATI	2.70	0.13	0.53	30, 2, 21, 8, 37, 11, 39,36
b	2.68	0.12	0.38	24, 40, 5, 26, 35, 18, 17, 23
bN	2.78	0.07	0.14	7, 3, 9, 25, 15, 14, 31,6
DI	2.77	0.11	0.14	2, 30, 21, 8, 11, 23, 36,37
DRI	2.99	0.03	0.46	2, 11, 21, 8, 23, 9, 1, 17
DTE	2.70	0.13	0.53	30, 21, 37, 8, 2, 39, 11,36
GM	2.70	0.12	0.60	1, 20, 4, 32, 9, 31, 12, 27
GMP	3.08	0.02	0.14	2, 9, 11, 6, 17, 10, 23,4
HM	3.08	0.02	0.59	2, 9, 11, 6, 17, 10, 23, 4
MP	3.10	0.01	0.28	2, 6, 9, 10, 11, 17, 19, 31
MRP	3.10	0.01	0.63	2, 11, 9, 6, 10, 34, 4, 31
MSTik1	3.10	0.01	0.95	6, 2, 9, 10, 17, 11, 19, 31
MSTik2	3.03	0.03	0.59	2, 11, 9, 23, 8, 21, 17,6
Pi	2.73	0.09	0.14	18, 31, 22, 11, 26, 9, 15, 3
RDI	2.70	0.13	0.14	30, 21, 37, 8, 2, 39, 11,36
RDY	3.10	0.015	0.53	2, 9, 11, 6, 10, 34, 4, 31
REI	3.10	0.015	0.53	2, 9, 11, 6, 10, 34, 4, 31
SDI	2.70	0.13	0.42	30, 21, 37, 8, 2, 39, 11,36
SNPI	2.94	0.02	0.81	1, 20, 4, 9, 31, 17, 19, 34
SSI	2.70	0.13	0.14	30, 21, 37, 39, 8, 2, 11,36
SSPI	2.70	0.13	0.34	30, 21, 8, 2, 37, 39, 11,36
STI	3.08	0.02	0.14	2, 9, 11, 6, 17, 10, 23,4
TOL	2.70	0.13	0.16	30, 21, 8, 2, 37, 39, 11,36
YI	2.99	0.03	0.63	2, 11, 8, 21, 23, 9, 1, 17

SSPI, ATI and DTE selected the genotypes 30, 21, 37, 39, 8, 2 and 11 as best performances. The genotypes 2, 11, 9, 23, 8, 21 and 17 were the best selections for MSTik2 and YI. According to GM and SNPI, the genotypes 1, 20, 4, 32, 9 and 31 exhibited the best rankings. The superiority index (Pi) selected the genotypes 18, 31, 22, 11 and 26. The coefficient of regression (b) identified the genotypes 24, 40, 5, 26 and 35 as best performances. The least values of the index bN were obtained by the genotypes 7, 3, 9, 25, 14 and 15. Finally, the highest values of DRI were obtained by the genotypes 2, 11, 21, 8, 23 and 9.

During the second season 2015 (Table 8), the indices MP, MRP, REI, GMP, STI, MSTik1, MSTik2, HM, YI and RDY selected the genotypes 11, 8, 9, 34, 10, 28 and 6 as top ranking. Similarly, SSI, TOL, SDI, RDI, DI, SSPI, DTE, bN, DRI, GM and SNPI selected the genotypes 2, 14, 9, 8 and 10 as best performances; except for the genotype 2 which was not selected by GM and SNPI. The least values of Pi were exhibited by the genotypes 16, 22, 33, 31, 15 and 18. The genotypes 5, 21, 16, 18, 29 and 37 expressed the lowest values of b. Finally, the genotypes 2, 14, 38, 9, 1, 20 and 40 were the top rankings for the index ATI.

Based on those results, the similarities between

drought indices in genotypes selection were in concordance with the previous correlation and biplot results. These findings showed that the indices MP, MRP, REI, GMP, MSTik1 and HM ranked similarly the genotypes. Their values are based on relative performance under various moisture conditions with little emphasis on yield stability. They have higher power in the separation of group A from the other Fernandez groups (Rosielle and Hamblin, 1981; Fernandez, 1992; Ramirez and Kelly, 1998; Golabadi et al., 2006; Talebi et al., 2009; Singh et al., 2011). The drought index RDY belonged also to the same grouping in the present study; even if this index emphasize on the selection of genotypes which have the minimum reduction in grain yield due to the moisture stress (Deshmukh et al., 2004); so it is more related to yield stability. For MSTik2 and YI, they were affiliated to this grouping only during 2015 when the stress intensity was 32%. Once the stress became harder during 2015 (43% of yield reduction), they constituted a separate common group because their formulation (equation) is based mainly on yield under stress conditions (Gavuzzi et al., 1997; Singh et al., 2011; Rahmani et al., 2013) in comparison to the other indices. In contrast, DI, SSI, TOL, SDI, RDI, DI, SSPI, DTE, bN and DRI emphasized more on yield stability and low

Table 8. Descriptive statistics of the top 20% genotype selection for drought indices during 2014-2015 cropping season.

Indices	Mean yield of 20% top genotypes	Variance inter 20% genotypes	Mean variance top 20% genotypes	Top 20% Genotypes selected
ATI	3.80	0.26	0.13	2, 14, 38, 9, 1, 20, 40, 8
b	3.96	0.23	0.12	5, 21, 16, 18, 29, 37, 23, 26
bN	4.14	0.16	0.13	2, 14, 9, 8, 10, 34, 1, 32
DI	4.29	0.20	0.14	2, 8, 9, 14, 11, 10, 34, 32
DRI	4.31	0.20	0.16	2, 8, 14, 9, 11, 6, 10, 34
DTE	4.14	0.16	0.13	2, 14, 9, 8, 10, 34, 1, 32
GM	4.16	0.14	0.15	14, 9, 8, 10, 34, 1, 32, 20
GMP	4.46	0.06	0.26	11, 8, 9, 34, 10, 28, 6, 32
HM	4.46	0.06	0.26	11, 8, 9, 34, 10, 28, 6, 32
MP	4.46	0.06	0.30	11, 8, 9, 34, 10, 28, 6
MRP	4.46	0.06	0.26	11, 8, 9, 34, 10, 28, 6
MSTIk1	4.46	0.06	0.33	11, 8, 9, 34, 10, 28, 6, 3
MSTIk2	4.46	0.06	0.26	11, 8, 9, 34, 10, 28, 6, 32
Pi	3.56	0.25	0.85	16, 22, 33, 31, 15, 18, 9, 27
RDI	4.14	0.16	0.13	2, 14, 9, 8, 10, 34, 1, 32
RDY	4.46	0.06	0.26	11, 8, 9, 34, 10, 28, 6, 32
REI	4.46	0.06	0.26	11, 8, 9, 34, 10, 28, 6
SDI	4.14	0.16	0.13	2, 14, 9, 8, 10, 34, 1, 32
SNPI	4.38	0.13	0.19	14, 8, 9, 11, 10, 34, 32, 28
SSI	4.14	0.16	0.13	2, 14, 9, 8, 10, 34, 1, 32
SSPI	3.96	0.23	0.12	2, 14, 9, 8, 1, 20, 38, 10
STI	4.46	0.06	0.26	11, 8, 9, 34, 10, 28, 6, 32
TOL	3.96	0.23	0.12	2, 14, 9, 8, 1, 20, 38, 10
YI	4.40	0.14	0.22	11, 8, 9, 10, 34, 2, 28, 6

changes between potential and actual yields under moisture conditions. In this case, top ranking genotypes are not necessarily high yielding (Fischer and Maurer, 1978; Fernandez, 1992; Clarke et al., 1992; Ramirez and Kelly, 1998; Guttieri et al., 2001; Sio- Se Mardeh et al., 2006; Golabadi et al., 2006). GM and SNPI had the same top ranking genotypes as SSI, TOL, SDI, RDI, DI, SSPI, DTE, bN and DRI during 2015. However, when the gap between the potential yield and yield under stress became larger (43% of reduction during 2014), those two indices were grouped together in a separate cluster because they put more emphasis on yield stability and high yield under stressed conditions (Farshadfar et al., 2002; Moradi et al., 2012) compared to the other indices.

The best indices should be able to select the highest and stable performances. Based on the mean yield and mean variance of the 20% top genotypes selection (Tables 7 and 8), the indices MP, MRP, REI, RDY, GMP, STI, HM, MSTIk1 and MSTIk2 identified the highest mean yields during 2014 (3.09 t/ha) and 2015 (4.47 t/ha) cropping seasons. In contrast, the mean variance of the top 20% genotypes varied from 0.14 to 0.95 in 2014 and from 0.26 to 0.33 during 2015. Thus, the mean variance between yield under stress and non-stress environments became higher at 43% of drought intensity compared to

32%. However, the indices GMP, MP and STI were able to exhibit the best combination of high mean yield and low mean variance.

Conclusion

The indices MP, MRP, REI, GMP, STI, MSTIk1, MSTIk2, HM and RDY showed high discrimination between genotypes, exhibited the best correlation with both yields under stressed and non-stressed environments and were able to identify the highest mean yielding genotypes with low variance across environments, especially STI, GMP and MP. These indices can be considered as suitable criteria for selection of drought tolerant and high yielding genotypes under moderate stress Mediterranean environment. The indices MP, MRP, REI, GMP, STI and RDY can be used interchangeably. The genotypes "AUS30355" and "Gladius" were recognized as best stable performances in the different moisture conditions. Our conclusions may be limited in terms of drought scenarios (duration, timing). The stress severity of our stressed environments was moderate consisting of drought at mid-stage (pre-flowering and flowering). More stress severities and drought scenarios may need to be

studied before confirming the general suitability of the different indices. Practically, these indices can be used immediately for semi arid environments of moderate drought severity, like areas located north of the latitude 33°N. Moreover, one particular disadvantage of these indices is their limitation on two contrasting environments at a time only, while the breeding for large adaptation usually uses a network of a wide range of environments. A development of data processing software will be useful in this case.

Abbreviations: **ATI**, abiotic tolerance index; **GM**, golden mean; **GMP**, geometric mean productivity; **HM**, harmony mean; **MP**, mean productivity; **MRP**, mean relative performance; **MSTik1**, modified stress tolerance index 1; **MSTik2**, modified stress tolerance index 2; **RDY**, relative decrease in yield; **REI**, relative efficiency index; **STI**, stress tolerance index; **SNPI**, stress/non-stress production index.

Conflict of Interests

The authors have not declared any conflict of interests.

REFERENCES

- Ahmadi A, Mohammadi V, Siosemardeh A, Poustinik (2004). Evaluation of wheat yield and drought resistance indices across water regimes. University of Tehran, Karaj, Iran. College of Agriculture, Kurdistan University, Sanandaj, Iran.
- Anwar J, Subhani GM, Hussain M, Ahmad J, Hussain M, Munir M (2011). Drought tolerance indices and their correlation with yield in exotic wheat genotypes. Pak. J. Bot. 43(3):1527-1530.
- Bansal KC, Sinha SK (1991). Assessment of drought resistance in 20 accessions of *Triticum aestivum* L. and related species, total dry matter and grain yield stability. Euphytica 56:7-14.
- Betran FJ, Beck D, Banziger M, Edmeades GO (2003). Genetic analysis of inbred and hybrid grain yield under stress and non-stress environments in tropical maize. Crop Sci. 43:807-817.
- Bidinger FR, Mahalakshmi V, Rao GDP (1987). Assessment of drought resistance in pearl millet (*Pennisetum americanum* (L) Leeke). I Factors affecting yields under stress. Aust. J. Agric. Res. 38:37-48.
- Bishaw Z, Struik PC, Van Gestel AJG (2011). Wheat and barley seed system in Syria: farmers' varietal perceptions, seed sources and seed management. Int. J. Plant Prod. 5(4):323-347.
- Blum A (1996). Crop responses to drought and the interpretation of adaptation. Plant Growth Regul. 20:135-148.
- Bousslama M, Schapaugh WT (1984). Stress tolerance in soybean. Part 1: Evaluation of three screening techniques for heat and drought tolerance. Crop Sci. 24:933-937.
- Boussen H, Ben Salem M, Slama A, Mallek-Maalej E, Rezgui S (2010). Evaluation of drought tolerance indices in durum wheat recombinant inbred lines. In: Lopez-Francos A. (Comp.), Lopez-Francos A. (Collab.). Economics of drought and drought preparedness in a climate change context. Zaragoza: CIHEAM/FAO/ICARDA/GDAR/CEIGRAM/MARM. Options Méditerranéennes Série A, Séminaires Méditerranéennes : N° 95.
- Clarke JM, De Pauw RM, Townley-Smith TM (1992). Evaluation of methods for quantification of drought tolerance in wheat. Crop Sci. 32:732-728.
- Dadbakhsh A, Yazdansepar A, Ahmadizadeh M (2011). Study drought stress on yield of wheat (*Triticum aestivum* L.) genotypes by drought tolerance indices. Adv. Envid. Biol. 5(7):1804-1810.
- Deshmukh DV, Mhase LB, Jamadagni BM (2004). Evaluation of chickpea genotypes for drought tolerance. Ind. J. Pulses Res. 17:47-49.
- Drikvand R, Doosty B, Hosseinpour T (2012). Response of rainfed wheat genotypes to drought stress using drought tolerance indices. J. Agric. Sci. 4(7):126.
- Eivazi AR, Mohammadi S, Razaee M, Ashori S, Pour FH (2013). Effective selection criteria for assessing drought tolerant indices in barley (*Hordeum vulgare* L.) accessions. Int. J. Agron. Plant Prod. 4(4):813-821.
- Farshadfar E, Ghanadha M, Zahravi M, Sutka J (2001). Generation mean analysis of drought tolerance in wheat. Acta. Agron. Hung. 49:59-66.
- Farshadfar E, Sutka J (2002). Multivariate analysis of drought tolerance in wheat substitution lines. Cereal Res. Commun. 31:33-39.
- Farshadfar E, Mohammadi R, Sutka J (2002). Association between field and laboratory predictors of drought tolerance in wheat disomic lines. Acta. Agron. Hung. 50(3):377-381.
- Farshadfar E, Javadinia J (2011). Evaluation of chickpea (*Cicer arietinum* L.) genotypes for drought tolerance. Seed and Plant Improv. J. 27(4):517-537.
- Farshadfar E, Farshadfar M, Dabiri S (2012a). Comparison between effective selection criteria of drought tolerance in bread wheat landraces of Iran. Ann. Biol. Res. 3(7):3381-3389.
- Farshadfar E, Elyasi P (2012). Screening quantitative indicators of drought tolerance in bread wheat (*T. aestivum*) landraces. Pelagia Research Library. Eur. J. Exp. Biol. 2(3):577-584.
- Farshadfar E, Pour SMM, Pour AAR (2012b). Repeatability of drought tolerance indices in bread wheat genotypes. Int. J. Agric. Crop Sci. 4(13):891-903.
- Farshadfar E, Mohammadi R, Farshadfar M, Dabiri S (2013). Relationships and repeatability of drought tolerance indices in wheat-rye disomic addition lines. Aust. J. Crop Sci. 7(1):130-138.
- Farshadfar E, Sheibanirad A, Soltanian M (2014). Screening landraces of bread wheat genotypes for drought tolerance in the field and laboratory. Int. J. Farm. Allied Sci. 3(3):304-311.
- Fernandez GCJ (1992). Effective selection criteria for assessing plant stress tolerance. In: Kuo CG, ed. Adaptation of Food Crops to Temperature and Water Stress. Shanhu: Asian Vegetable Research and Development Center. Taiwan Publication 93(410):257-270.
- Fischer KS, Wood G (1981). Breeding and selection for drought tolerance in tropical maize. In the Proceeding of the Symposium on Principles and Methods in Crop Improvement for Drought Resistance with Emphasis on Rice, IRRRI, Philippines.
- Fischer RA, Maurer R (1978). Drought resistance in spring wheat cultivars. Aust. J. Agric. Res. 29:897-912.
- Fischer RA, Wood JT (1979). Drought resistance in spring wheat cultivars III. Yield association with morphological traits. Aust. J. Agric. Res. 30: 1001-1020.
- Gavuzzi P, Rizza F, Palumbo M, Campaline RG, Ricciardi GL, Borghi B (1997). Evaluation of field and laboratory predictors of drought and heat tolerance in winter cereals. Can. J. Plant Sci. 77:523-531.
- Geravandi M, Farshadfar E, Kahrizi D (2011). Evaluation of some physiological traits as indicators of drought tolerance in bread wheat genotypes. Russ. J. Plant Physiol. 58(1):69-75.
- Gholinezhad E, Darvishzadeh R, Bernousi I (2014). Evaluation of Drought Tolerance Indices for Selection of Confectionery Sunflower (*Helianthus annuus* L.) Landraces under Various Environmental Conditions. Bot. Horti Agrobot. Cluj-Napoca 42(1):187-201.
- Golabadi M, Arzani A, Mirmohammadi Maibody SAM (2006). Assessment of drought tolerance in segregating populations in durum wheat. Afr. J. Agric. Res. 1(5):162-171.
- Guttieri MJ, Stark JC, Brien K, Souza E (2001). Relative sensitivity of spring wheat grain yield and quality parameters to moisture deficit. Crop Sci. 41:327-335.
- Hossain ABS, Sears AG, Cox TS, Paulsen GM (1999). Desiccation tolerance and its relationship to assimilate partitioning in winter wheat. Crop Sci. 30:622-627.
- İlker E, Tatar Ö, Aykut Tonk F, Tosun M, Turk J (2011). Determination of Tolerance Level of Some Wheat Genotypes to Post-Anthesis Drought. Turkish J. Field Crops 16(1):59-63.

- Jafari A, Paknejad F, Jami Al-Ahmadi M (2009). Evaluation of selection indices for drought tolerance of corn (*Zea mays* L.) hybrids. *Int. J. Plant Prod.* 3(4):33-38.
- Jilibene M (2009). Amélioration génétique du blé tendre au Maroc à l'aube du 21ème siècle. 80 pages, Editions INRA.
- Karamanos AJ, Papatheohari AY (1999). Assessment of drought resistance of crop genotypes by means of the water potential index. *Crop Sci.* 39:1792-1797.
- Karamanos AJ, Economou G, Papastavrou A, Travlos IS (2012). Screening of Greek wheat landraces for their yield responses under arid conditions. *Inter. J. of Plant Prod.* 6 (2):1735-8043.
- Khalili M, Kazemi M, Moghaddam A, Shakiba M (2004). Evaluation of drought tolerance indices at different growth stages of late-maturing corn genotypes. *Proceedings of the 8th Iranian Congress of Crop Science and Breeding.* Rasht, Iran. P 298.
- Khodarahmpour Z, Choukan R, Bihamta MR, Majidi Harvan E (2011). Determination of the Best Heat Stress Tolerance Indices in Maize (*Zea mays* L.) Inbred Lines and Hybrids under Khuzestan Province Conditions. *J. Agric. Sci. Technol.* 13:111-121.
- Kumar S, Dwivedi SK, Singh SS, Jha SK, Lekshmy S, Elanchezian R, Singh ON, Bhatt BP (2014). Identification of drought tolerant rice genotypes by analyzing drought tolerance indices and morpho-physiological traits. *SABRAO J. Breed. Genet.* 46(2):217-230.
- Lan J (1998). Comparison of evaluating methods for agronomic drought resistance in crops. *Acta Agric. Bor-occid Sinic* 7:85-87.
- Lin CS, Binns MR, Lefkovich LP (1986). Stability analysis: where do we stand? *Crop Sci.* 26: 894-900.
- Mevlüt A, Sait Ç (2011). Evaluation of drought tolerance indices for selection of Turkish oat (*Avena sativa* L.) landraces under various environmental conditions. *Zemdirbyste Agric.* 98(2):157-166.
- Mitra J (2001). Genetics and genetic improvement of drought resistance in crop plants. *Curr. Sci.* 80:758-762.
- Moghaddam A, Hadizadeh MH (2002). Response of corn hybrids and their parental lines to drought using different stress tolerant indices. *Seed Plant Improv. J.* 18(3):255-272.
- Mohammadi R, Farshadfar E, Aghaee M, Shutka J (2003). Locating QTLs controlling drought tolerance criteria in rye using disomic addition lines. *Cereal Res. Commun.* 31:257-263.
- Mohammadi R, Armion M, Kahrizi D, Amri A (2010). Efficiency of screening techniques for evaluating durum wheat genotypes under mild drought condition. *Int. J. Plant Prod.* 4(1):11-24.
- Mohammadi M, Karimizadeh R, Abdipour M (2011). Evaluation of drought tolerance in bread wheat genotypes under dryland and supplemental irrigation conditions. *Aust. J. Crop Sci.* 5(4):487-493.
- Moosavi SS, Samadi YB, Naghavi MR, Zali AA, Dashti H, Pourshahbazi A (2008). Introduction of new indices to identify relative drought tolerance and resistance in wheat genotypes. *Desert* 12:165-178.
- Moradi H, Akbari GA, Khorasani SK, Ramshini HA (2012). Evaluation of drought tolerance in corn (*Zea Mays* L.) new hybrids with using stress tolerance indices. *Eur. J. Sustain. Dev.* 1(3):543-560.
- Naghavi MR, Pouraboughadareh A, Khalili M (2013). Evaluation of drought tolerance indices for screening some of corn (*Zea mays* L.) cultivars under environmental conditions. *Not. Sci. Biol.* 5(3):388-393.
- Normand Moayed F, Rostami A, Ghanadha MR (2001). Evaluating drought tolerance indices in bread wheat. *Iran. J. Agric. Sci.* 32(4):795-805.
- Nouraein M, Abolghasem Mohammadi S, Aharizad S, Moghaddam M, Sadeghzadeh B (2013). Evaluation of drought tolerance indices in wheat recombinant inbred line population. *Ann. Biol. Res.* 4(3):113-122.
- Nouri A, Etmnan A, Teixeira da Silva JA, Mohammadi R (2011). Assessment of yield, yield-related traits and drought tolerance of durum wheat genotypes (*Triticum turjidum* var. durum Desf.). *Aust. J. Crop Sci.* 5(1):8-16.
- Passioura JB (2007). The drought environment: physical, biological and agricultural prospective. *J. Exp. Bot.* 58:113-117.
- Rahmani S, Farshadfar E, Jowhar MM (2013). Locating QTLs controlling yield based indicators of drought tolerance in agropyron using wheat agropyron disomic addition lines. *Int. J. Agric. Crop Sci.* 5(9):1028-1033.
- Rajaram S, Van Ginkle M (2001). Mexico, 50 years of international wheat breeding. In *The World Wheat Book: A History of Wheat Breeding.* Eds., Bonjean A.P. and W.J. Angus. Lavoisier Publishing, Paris, France. pp. 579-604.
- Ramirez P, Kelly JD (1998). Traits related to drought resistance in common bean. *Euphytica* 99:127-136.
- Richards RA (1996). Defining selection criteria to improve yield under drought. *Plant Growth Regul.* 20:157-166.
- Rosielle AA, Hamblin J (1981). Theoretical aspects of selection for yield in stress and non-stress environments. *Crop Sci.* 21:943-946.
- Shirinzadeh A, Zarghami R, Azghandi AV, Shiri MR, Mirabdulbaghi M (2010). Evaluation of Drought Tolerance in Mid and Late Mature Corn Hybrids Using Stress Tolerance Indices. *Asian J. Plant Sci.* 9(2):67-73.
- Singh BU, Rao KV, Sharma HC (2011). Comparison of selection indices to identify sorghum genotypes resistant to the spotted stemborer, *Chilo partellus* (Swinhoe) (Lepidoptera: Noctuidae). *Int. J. Trop. Insect Sci. Open Access ICRISAT, Patancheru.*
- Sio-Se Mardeh A, Ahmadi A, Poustini K, Mohammadi V (2006). Evaluation of drought resistance indices under various environmental conditions. *Field Crops Res.* 98(2-3):222-229.
- Talebi R, Fayaz F, Naji AM (2009). Effective selection criteria for assessing drought stress tolerance in durum wheat (*Triticum durum* Desf.). *Gen. Appl. Plant Physiol.* 35: 64-74.
- Travlos IS (2012). Reduced herbicide rates for an effective weed control in competitive wheat cultivars. *Int. J. Plant Prod.* 6:1-14.

