

*Full Length Research Paper*

# Effect of “eight-trigram boxing” exercise on blood oxidative status and intestine bifidobacterium and lactobacillus count in practicers

Yin Hai-li<sup>1</sup> and Niu Ai-jun<sup>2,3\*</sup>

<sup>1</sup>Institute of Physical Education, Ludong University, Yantai, Shandong Province, 264025, China.

<sup>2</sup>Wushu Department, Guangzhou Sport University, Guangzhou, 510500, China.

<sup>3</sup>History Department, Fudan University, Shanghai, 200433, China.

Accepted 03 March, 2018

**The aim of this study was to evaluate the effectiveness of “Eight-trigram boxing” in reducing blood oxidative status and improving intestine beneficial bacteria counts. Participants were randomly assigned to “Eight-trigram boxing” exercise. After 3 months, significant increase in GSH, SOD, CAT, GSH-Px and GR activities were observed. In addition, intestine beneficial bacteria bifidobacterium and lactobacillus count were markedly enhanced. It could be concluded that “Eight-trigram boxing” exercise was beneficial to practicers’ health.**

**Key words:** “Eight-trigram boxing”, antioxidant, bifidobacterium, lactobacillus.

## INTRODUCTION

Qigong has been practiced for thousands of years in China and in recent years, slowly but surely, it is spreading to the west (Yan et al., 1998). Millions of people, each year, are helping themselves get rid of all kind of diseases, ranging from minor aches and pains through to the most debilitating ones. It is becoming quite apparent that its future looks bright and that it can be beneficial to whoever practices it, regardless of sex, age, race, religion, level of education or status (Sakata et al., 2008).

The “Eight-trigram boxing”, one of Qigong, originally named Palm turning boxing, body shifting eight trigram boxing and interlocked eight trigram boxing, is characterized by changing positions of the palms and changing paces, somewhat resembling the Eight trigrams in the “Book of changes” (Pan and Feng, 2010). Through long period of exercise, the muscles are gently stretched and strengthened. The bones and ligaments become stronger. Organ stimulation occurs and blood flows. Balance and coordination are improved with an overall vitality and vigour towards life (Miao et al., 2009).

It is now accepted that free radical- mediated oxidation of biological molecules such as lipids, proteins, and DNA is involved in a variety of disorders and diseases (Sun et al., 2010; Aliyu et al., 2009). Above all, lipids are very susceptible to free radical attack and lipid peroxidation induces alterations in integrity, disturbances in fine structure, and functional loss of biomembranes.

Furthermore, lipid peroxidation mediated by free radicals proceeds by a chain mechanism, amplifying the damaging effect of free radicals and. Lipid peroxidation products are potentially cytotoxic and modify proteins and DNA (Omata et al., 2010; Hu et al., 2010; Liu, Wang and Liu, 2009). It is now accepted that lipid peroxidation is involved in the pathogenesis of various diseases, and consequently the role of free radical-scavenging antioxidants has received much attention (Kohri et al., 2009).

Bifidobacteria are considered as important probiotics and used in the food industry to relieve and treat many intestinal disorders. Bifidobacteria exert a range of beneficial health effects, including the regulation of intestinal microbial homeostasis, the inhibition of pathogens and harmful bacteria that colonize and/or infect the gut mucosa, the modulation of local and systemic immune responses, the repression of

\*Corresponding author. E-mail: niuaj967@yahoo.cn.

procarcinogenic enzymatic activities within the microbiota, the production of vitamins, and the bioconversion of a number of dietary compounds into bioactive molecules (Fu, 2009; Killer et al., 2010; Rahman et al., 2008). *Lactobacillus* is a genus of Gram-positive facultative anaerobic or microaerophilic bacteria. They are a major part of the lactic acid bacteria group, named as such because most of its members convert lactose and other sugars to lactic acid. They are common and usually benign. In humans they are present in the vagina and the gastrointestinal tract, where they are symbiotic and make up a small portion of the gut flora. Many species are prominent in decaying plant material. The production of lactic acid makes its environment acidic, which inhibits the growth of some harmful bacteria. Several members of the genus have had their genome sequenced (Singh et al., 2009; Pratumvinit et al., 2009; Giraffa et al., 2010).

In this study, we examined effect of “Eight-trigram boxing” exercise on blood oxidative status in practicers. Then, effect of “Eight-trigram boxing” exercise on intestine bifidobacterium and lactobacillus count were also investigated.

## **PARTICIPANTS AND METHODS**

### **Subjects and procedures**

Fifty-two subjects (38 women and 14 men aged  $37.07 \pm 5.16$  years with a body height of  $167.0 \pm 8.1$  cm, a body weight of  $61.21 \pm 6.38$  kg) with no previous injuries in the year before the study were recruited. All practicers do not have “Eight-trigram boxing” exercise experience. Subjects attended a 80 min “Eight-trigram boxing” exercise 3 times per day for 5 months, located at the community center. During practice, blood was taken from all subjects per month. Feces were also collected from all subjects per month.

### **Biochemical analysis**

TC, TG, LDL-c and HDL-c levels were determined using spectrophotometer and standard commercial kits. GSH level was determined by DTNB method (Nannelli et al., 2007). MDA level was examined with TBA fluoremetry (Hepgül et al., 2010). And the total SOD activity was determined by pyrogallol-NBT colorimetry (Cigremis et al., 2009). The CAT activity in liver extracts was measured spectrophotometrically at 240 nm by calculating the rate of degradation of  $H_2O_2$ , the substrate of the enzyme (wirsk-Korczała et al., 2004). Glutathione peroxidase (GSH-Px, EC 1.6.4.2) activity was measured by the method of Kocer et al. (2006). Glutathione reductase (GR) activity was measured using the procedure described by Błaszczuk et al. (2008).

### **Counting of bifidobacteria and lactobacillus in feces**

Feces of subjects were analyzed for the number of total bifidobacteria and lactobacillus using the procedure described by Kawase et al. (1982).

## **Statistical analysis**

Statistical analysis was performed using the SPSS-PC 11.0 software (SPSS Inc., Chicago, Illinois, USA). One-way ANOVA and LSD test were performed in order to determine whether there was any significant difference during different exercise time.

## **RESULTS**

### **Effect of “eight-trigram boxing” on MDA and GSH levels**

The results of the changes of MDA and GSH levels in practicers' blood are represented in Table 1. As can be observed, there was a decrease in MDA level and increase in GSH level, respectively. In both cases, the effect was statistically significant ( $P < 0.05$ ;  $P < 0.01$ ). These effects were strengthened with increasing exercise time.

### **Effect of “eight-trigram boxing” on TC, TG, LDL-c and HDL-c levels**

The results of the changes of TC, TG, LDL- c and HDL-c levels in practicers are represented in Table 2. As can be observed, there was a decrease in TC, TG, and LDL-c levels. On the contrary, HDL-c level increased significantly ( $P < 0.05$ ;  $P < 0.01$ ). In both cases, the effect was statistically significant ( $P < 0.05$ ;  $P < 0.01$ ). These effects were shown to develop progressively over exercise time.

### **Effect of “eight-trigram boxing” on SOD, CAT, GSH-Px and GR activities**

As demonstrated in Table 3, SOD, CAT, GSH-Px and GR activities in practicers were significantly ( $P < 0.05$ ;  $P < 0.01$ ) increased. The effect were significantly ( $P < 0.05$ ;  $P < 0.01$ ) increased in a time-dependent manner. Namely, the effect increased with increasing exercise time.

### **Effect of “eight-trigram boxing” on intestine bifidobacterium and lactobacillus count**

Table 4 showed the role of “Eight- trigram boxing” in modulating intestine bifidobacterium and lactobacillus count during exercise. Highly significant enhancement ( $P < 0.05$ ;  $P < 0.01$ ) in the count of intestine bifidobacterium and lactobacillus were obtained during exercise. Moreover, with increasing exercise time, intestine bifidobacterium and lactobacillus count accordingly increased.

**Table 1.** Effect of “Eight-Trigram Boxing” on MDA and GSH levels.

	0 (month)	1 (month)	2 (month)	3 (month)	4 (month)	5 (month)
MDA (mmol/mg)	12.38±1.07	12.15±1.13	9.02±1.11 <sup>a</sup>	7.47±0.63 <sup>b</sup>	7.05±0.66 <sup>b</sup>	6.53±0.54 <sup>b</sup>
GSH ( mol /g)	44.25±2.93	49.68±2.08 <sup>a</sup>	58.04±3.71 <sup>b</sup>	63.75±5.07 <sup>b</sup>	77.12±5.39 <sup>b</sup>	80.62±7.31 <sup>b</sup>

<sup>a</sup> P < 0.05, <sup>b</sup> P < 0.01, compared with control (0 month).

**Table 2.** Effect of “Eight-Trigram Boxing” on TC, TG, LDL-c and HDL-c levels.

	0 (month)	1 (month)	2 (month)	3 (month)	4 (month)	5 (month)
TC (mmol L <sup>-1</sup> )	1.73±0.18	1.68±0.11	1.54±0.08 <sup>b</sup>	1.48±0.11 <sup>b</sup>	1.32±0.06 <sup>b</sup>	1.28±0.10 <sup>b</sup>
TG (mmol L <sup>-1</sup> )	0.39±0.04	0.34±0.03	0.31±0.02 <sup>b</sup>	0.29±0.03 <sup>b</sup>	0.29±0.02 <sup>b</sup>	0.27±0.02 <sup>b</sup>
LDL-c (mmol L <sup>-1</sup> )	0.42±0.04	0.41±0.03	0.38±0.04 <sup>a</sup>	0.34±0.03 <sup>b</sup>	0.31±0.04 <sup>b</sup>	0.29±0.03 <sup>b</sup>
HDL-c (mmol L <sup>-1</sup> )	1.31±0.21	1.46±0.14 <sup>a</sup>	1.57±0.21 <sup>b</sup>	1.63±0.15 <sup>b</sup>	1.82±0.16 <sup>b</sup>	2.06±0.15 <sup>b</sup>

<sup>a</sup> P<0.05, <sup>b</sup> P<0.01, compared with control (0 month).

**Table 3.** Effect of “Eight-Trigram Boxing” on SOD, CAT, GSH-Px and GR activities.

	0 (month)	1 (month)	2 (month)	3 (month)	4 (month)	5 (month)
SOD(U/mg)	79.09±6.09	89.33±6.77 <sup>b</sup>	98.09±6.92 <sup>b</sup>	132.11±11.65 <sup>b</sup>	146.08±12.35 <sup>b</sup>	153.07±11.09 <sup>b</sup>
CAT(U/mg)	9.56±0.87	12.07±0.69 <sup>a</sup>	15.25±1.63 <sup>b</sup>	17.07±1.12 <sup>b</sup>	27.09±2.83 <sup>b</sup>	28.14±2.04 <sup>b</sup>
GSH-Px(U/mg)	18.95±1.43	24.75±2.09 <sup>b</sup>	30.09±1.79 <sup>b</sup>	32.11±2.48 <sup>b</sup>	34.09±2.16 <sup>b</sup>	35.01±1.95 <sup>b</sup>
GR(U/mg)	14.96±1.08	18.09±1.67 <sup>a</sup>	23.75±1.85 <sup>b</sup>	25.09±2.84 <sup>b</sup>	26.11±1.75 <sup>b</sup>	28.09±2.94 <sup>b</sup>

<sup>a</sup> P<0.05, <sup>b</sup> P<0.01, compared with control (0 month).

**Table 4.** Effect of “Eight-Trigram Boxing” on intestine bifidobacterium and lactobacillus count (log CFU /g, x ± s, n = 5).

	0 (month)	1 (month)	2 (month)	3 (month)	4 (month)	5 (month)
bifidobacterium	7.45±0.31	8.24±0.75 <sup>a</sup>	8.89±0.57 <sup>b</sup>	9.23±0.77 <sup>b</sup>	9.64±0.83 <sup>b</sup>	10.42±0.65 <sup>b</sup>
lactobacillus	11.57±0.93	12.57±0.58 <sup>a</sup>	13.08±1.08 <sup>b</sup>	13.65±1.32 <sup>b</sup>	14.01±2.09 <sup>b</sup>	15.19±1.94 <sup>b</sup>

<sup>a</sup> P<0.05, <sup>b</sup> P<0.01, compared with control (0 month).

## DISCUSSION

Age, diet, and strenuous exercise could all causes oxidative stress, resulting in lipid peroxidation and DNA damage (Błaszczuk et al., 2010; Qiao et al., 2009).

Oxidative injury and inflammatory response involving oxygen free radicals have been implicated to be of vital importance in healthy people and patients, which sometimes lead to higher morbidity (Jia et al, 2009; Fan et al, 2009).

This work presents the results from a controlled experiment where the effect of “Eight-trigram boxing” exercise on changes in plasma lipid peroxidation, capacity antioxidant and intestine bifidobacterium and lactobacillus count in practice. In these trained subjects, a significant increase in GSH levels, SOD, CAT, GSH- Px and GR activities was found in relation to exercise. In addition, a decrease in TC, TG, LDL-c levels and increase

in HDL-c were detected in practicers. Moreover, these beneficial effects were further improved with increasing exercise time. These suggest that “Eight-trigram boxing” exercise enhanced antioxidant status and reduced oxidative damage in practicers’ body.

Probiotic bacteria favorably alter the intestinal microflora balance, inhibit the growth of harmful bacteria, promote good digestion, boost immune function, and increase resistance to infection (Smirnov et al., 1993). People with flourishing intestinal colonies of beneficial bacteria are better equipped to fight the growth of disease-causing bacteria (Mel’nikova et al., 1993; De Simone et al., 1993). Lactobacilli and bifidobacteria maintain a healthy balance of intestinal flora by producing organic compounds—such as lactic acid, hydrogen peroxide, and acetic acid—that increase the acidity of the intestine and inhibit the reproduction of many harmful bacteria (Veldman et al., 1992; Kawase, 1982). Probiotic

bacteria also produce substances called bacteriocins, which act as natural antibiotics to kill undesirable microorganisms (Rasic, 1983). Our work showed that “Eight-trigram boxing” exercise could increase intestine bifidobacterium and lactobacillus count.

## Conclusion

Cumulative ROS generated in body could induce oxidative injury. “Eight-Trigram Boxing” exercise could activate antioxidant enzymes activities and reduced lipid peroxidation level. In additionm, “Eight-trigram boxing” exercise could still increased intestine bifidobacterium and lactobacillus count. Furthermore, the effect became better with prolonging exercise period. These results suggest that “Eight-trigram boxing” is useful for practicer’s health.

## ACKNOWLEDGEMENT

Academic Discipline Project from Ludong University; Shandong province Social Science Key Constructional Projects Foundation (09BTYJ01) supported the present study.

## REFERENCES

- Aliyu AB, Ibrahim MA, Musa AM, Ibrahim H, Abdulkadir IE, Oyewale AO (2009). Evaluation of antioxidant activity of leave extract of *Bauhinia rufescens* Lam. (Caesalpinaceae). *J. Med. Plants Res.*, 3(8): 563-567.
- Błaszczuk I, Grucka-Mamczar E, Kasperczyk S, Birkner E (2008). Influence of Fluoride on Rat Kidney Antioxidant System: Effects of Methionine and Vitamin E. *Biol Trace Elem. Res.*, 121: 51-59.
- Błaszczuk I, Grucka-Mamczar E, Kasperczyk S, Birkner E (2010). Influence of Methionine upon the Activity of Antioxidative Enzymes in The Kidney of Rats Exposed to Sodium Fluoride. *Biol. Trace Elem. Res.*, 133: 60-70.
- Cigremis Y, Turel H, Adiguzel K, Akgoz M, Kart A, Karaman M, Ozen H (2009). The effects of acute acetaminophen toxicity on hepatic mRNA expression of SOD, CAT, GSH-Px, and levels of peroxynitrite, nitric oxide, reduced glutathione, and malondialdehyde in rabbit. *Mol. Cell Biochem.*, 323: 31-38.
- De Simone C, Vesely R, Bianchi SB, Jirillo E (1993). The role of probiotics in modulation of the immune system in man and in animals. *Int. J. Immunother.*, 9: 23–28.
- Fan YJ, He XJ, Zhou SD, Luo AX, He T, Chun Z (2009). Composition analysis and antioxidant activity of polysaccharide from *Dendrobium denneanum*. *Int. J. Biol. Macromol.*, 45(2): 169-173.
- Fu YL (2009). Isolation, purification, and structural elucidation of a fructan from *Arctium lappa* L. *J. Med. Plants Res.*, 3(3): 171-173.
- Giraffa G, Chanishvili N, Widyastuti Y (2010). Importance of lactobacilli in food and feed biotechnology. *Res. Microbiol.*, 161(6): 480-487.
- Hepgül G, Tanrikulu S, Ünalp HR, Akguner T, Erbil Y, Olgaç V, Ademo lu E (2010). Preventive Effect of Pentoxifylline on Acute Radiation Damage via Antioxidant and Anti-inflammatory Pathways. *Digest Dis. Sci.*, 55: 617-625.
- Hu TT, Liu D, Chen Y, Wu J, Wang SS (2010). Antioxidant activity of sulfated polysaccharide fractions extracted from *Undaria pinnatifida* *in vitro*. *Int. J. Biol. Macromol.*, 46(2): 193-198.
- Jia ZQ, Zhu H, Li YB, Misra HP (2009). Cruciferous Nutraceutical 3H-1,2-dithiole-3-thione Protects Human Primary Astrocytes Against Neurocytotoxicity Elicited by MPTP, MPP+, 6-OHDA, HNE and Acrolein. *Neurochem Res.*, 34: 1924-1934.
- Kawase K (1982). Effects of nutrients on the intestinal microflora of infants. *Jpn. J. Dairy Food Sci.*, 31:A241–243.
- Killer J, Kope ný J, Mrázek J, Rada V, Dubná S, Marounek M (2010). Bifidobacteria in the digestive tract of bumblebees. *Anaerobe*, 16(2): 165-170.
- Kocer I, Taysi S, Ertekin MV, Karslioglu I, Gepdiremen A, Sezen O, Serifoglu K (2006). The effect of L-carnitine in the prevention of ionizing radiation-induced cataracts: a rat model. *Graef. Arch. Clin. Exp. Ophthalmol.*, 245: 588-594.
- Kohri S, Fujii H, Oowada S, Endoh N, Sueishi Y, Kusakabe M, Shimmei M, Kotake Y (2009). An oxygen radical absorbance capacity-like assay that directly quantifies the antioxidant’s scavenging capacity against AAPH-derived free radicals. *Anal. Biochem.*, 386(2): 167-171.
- Lin PP, Hsieh YM, Tsai CC (2009). Antagonistic activity of *Lactobacillus acidophilus* RY2 isolated from healthy infancy feces on the growth and adhesion characteristics of enteroaggregative *E. coli*. *Anaerobe*, 15(4): 122-126.
- Liu LH, Wang LS, Liu XM (2009). Comparison of antitussive, expectorant and antiasthmatic activities of different extracts from *Ficus microcarpa*. *J. Med. Plants Res.*, 3(8): 596-599.
- Mel’nikova VM, Gracheva NM, Belikov GP, Blatun LA, Shcherbakova EG (1993). The chemoprophylaxis and chemotherapy of opportunistic infections. *Antibiotiki i Khimioterapiia*, 38: 44–48.
- Miao FS, Liu Xiang Y, Li Y, Wang XC (2009). Effect of Fitness Qigong Baduanjin on plasma lipid and lipoprotein metabolism of patients with hyperlipidemia. *Journal of Shandong Institute of Physical Education and Sports*. 25(10): 46-48. (in Chinese).
- Nannelli A, Messina A, Marini S, Trasciatti S, Longo V, Gervasi PG (2007). Effects of the anticancer dehydrotarplatin on cytochrome P450 and antioxidant enzymes in male rat tissues. *Arch. Toxicol.*, 81: 479-487.
- Omata Y, Saito Y, Yoshida Y, Jeong BS, Serwa R, Nam TG, Porter NA, Niki E (2010). Action of 6-amino-3-pyridinolols as novel antioxidants against free radicals and oxidative stress in solution, plasma, and cultured cells. *Free Radic Biol. Med.*, 48(10): 1358-1365.
- Pan HS, Feng YC (2010). Clinical Observation of Rehabilitation Therapy with Health Qigong BaDuanJin on Grade 1 Hypertension of Old Patients. *Journal of Nanjing Institute of Physical Education (NaturalScience)*, 9(1): 4-6. (in Chinese).
- Pratumvinit B, Srisapoomi T, Worawattananon P, Oparkiatikul N, Jiratchariyakul W, Kummalue T (2009). In vitro antineoplastic effect of *Ficus hispida* L. plant against breast cancer cell lines. *J. Med. Plants Res.*, 3(4): 255-261.
- Qiao DL, Ke CL, Hu B, Luo JG, Ye H, Sun Y, Yan XY, Zeng XX (2009). Antioxidant activities of polysaccharides from *Hyriopsis cumingii*. *Carbohydr Polym.*, 78(2): 199-204.
- Rahman MM, Kim WS, Kumura H, Shimazaki KI (2008). In vitro effects of bovine lactoferrin on autoaggregation ability and surface hydrophobicity of bifidobacteria. *Anaerobe*, 14(2): 73-77.
- Rasic JL (1983). The role of dairy foods containing bifido and acidophilus bacteria in nutrition and health. *N Eur. Dairy J.*, 4: 80-88.
- Sakata T, Li QM, Tanaka M, Tajima F (2008). Positive effects of a qigong and aerobic exercise program on physical health in elderly Japanese women: an exploratory study. *Environ. Hlth. Preventive Med.*, 13: 162-168.
- Singh S, Goswami P, Singh R, Heller KJ (2009). Application of molecular identification tools for *Lactobacillus*, with a focus on discrimination between closely related species: A review. *LWT - Food Sci. Technol.*, 42(2): 448-457.
- Smirnov VV, Reznik SR, V’iunitskaia VA, Sorokulova IB, Samgorodskaja NV, Tofan AV (1993). The current concepts of the mechanisms of the therapeutic-prophylactic action of probiotics from bacteria in the genus *bacillus*. *Mikrobiolohichnyi Zhurnal* 55: 92-112.
- Sun YX, Kennedy JF (2010). Antioxidant activities of different polysaccharide conjugates (CRPs) isolated from the fruiting bodies of *Chroogomphus rutilus* (Schaeff.: Fr.) O. K. Miller. *Carbohydr. Polym.*, 82(2): 510-514.
- Veldman A (1992). Probiotics. *Tijdschrift voor Diergeneeskunde*, 117: 345-348.

Yan X, Lin H, Li HM, Traynor-Kaplan A, Xia, ZQ, Lu F, Fang Y, Dao M (1998). Structure and property changes in certain materials influenced by the external qi of qigong. *Mater Res. Innov.* 2: 349-359.

Zwirska-Korczala K, Adamczyk-Sowa M, Polaniak R, Sowa P, Birkner E, Drzazga Z, Brzozowski T, Konturek SJ (2004). Influence of

extremely-lowfrequency magnetic field on antioxidative melatonin properties in AT478 murine squamous cell carcinoma culture. *Biol. Trace Elem. Res.*, 102: 227-243.