

Full Length Research paper

Biodeterioration of the African star apple (*Chrysophyllum albidum*) in storage and the effect on its food value

N. A. Amusa*, O.A. Ashaye and M. O. Oladapo

Institute of Agricultural Research and Training Obafemi Awolowo University. Moor Plantation, PMB 5029 Ibadan. Nigeria.

Accepted 8 February 2021

The biodeterioration of the African star apple fruits in storage was investigated at Ibadan, southwestern Nigeria. Eight fungal isolates were found associated with the deteriorating fruits. The fungi are *Botryodiplodia theobromae*, *Rhizopus stolonifer*, *Aspergillus niger*, *A. tamarii*, *A. flavus*, *Fusarium spp*, *Penicillium spp* and *Trichoderma spp*. All the fungal isolates were pathogenic on the star apple fruits with the exception of *Trichoderma spp*. The African star apple fruits stored for up to 5 days were associated with severe fungal infections and had significantly reduced crude protein, crude fat and moisture content while dry matter, potassium, calcium and sodium increased compared to the freshly harvested fruits.

Key words: *Chrysophyllum albidum*, biodeterioration, fungal pathogens, storage.

INTRODUCTION

Chrysophyllum albidum (Linn.) belongs to the family Sapotaceae. It is primarily a forest tree species and its natural occurrences have been reported in diverse ecozones in Nigeria, Uganda, Niger Republic, Cameroon and Cote d'Ivoire (Bada, 1997). The plant often grows to a height of 36.5m though it may be smaller (Bada, 1997). The african star apple fruit is a large berry containing 4 to 5 flattened seeds or some times fewer due to seed abortion (Keay, 1989). The plant has in recent times become a crop of commercial value in Nigeria. The fleshy pulp of the fruits is eaten especially as snack and relished by both young and old (Cenrad, 1999). The African star apple fruit has been found to have highest content of ascorbic acid with 1000 to 3,330 mg of ascorbic acid per 100gm of edible fruit or about 100 times that of oranges and 10 times of that of guava or cashew (Asenjo, 1946). It is reported as an excellent source of vitamins, irons, flavours to diets and raw materials to some manufacturing industries (Adisa, 2000; Bada, 1997; Okafor and Fernandes, 1987; Umelo, 1997). In addition, its seeds are a source of oil, which is used for diverse purposes. The seeds are also used for local games (Bada, 1997). The fruits also contain 90% anardic acid, which is used industrially in protecting wood and as source of resin, while several other components of the tree including the roots and leaves are used for medicinal purposes (Adewusi, 1997; Bada, 1997).

C. albidum fruit is common in both urban and rural center especially during the months of December to April. The fruits are not usually harvested from the trees, but left drop naturally to the forest floor where they are picked. Allowing the fruits to drop before picking promotes fungal infections. Recent market survey revealed that the fruits often deteriorate within a very short period. According to Adebisi (1997), *C. albidum* actually becomes bad in a period of 5 days, with the deterioration starting with change of colour from uniform orange to one with patches, and followed by shrinking of the fruit. This study was undertaken to investigate the etiology of post-harvest biodeterioration of african star apple fruits in Ibadan, South Western Nigeria and the effects on its nutrient (food) value

MATERIALS AND METHODS

Twenty five african star apple trees from Moniya and Apata areas of Ibadan in south western Nigeria were used for the studies in 2000 and 2001. Ibadan (7° 20'N, 3° 50'E: 200 mm above sea level) is in a transition zone between the humid forest and derived savannah agro- ecologies of Nigeria. It has a mean annual rainfall of 1200 mm and mean daily temperature of between 24°C (min) and 34°C (max). Infected fruits were collected from above mentioned locations and brought to the laboratory for further studies.

The infected portions of the fruits were sliced into 2 mm² pieces, surface sterilized for 3 min with 1% NaOCl and rinsed in 4 successive changes of sterile distilled water. The surface-sterilized infected portions were then plated on sterile potato dextrose agar (PDA) in petri dishes and incubated for six days under alternating

*Corresponding author; E-mail: drart@infoweb.abs.net

12-hour light and dark periods at 26⁰C. The fungal isolates were examined under a stereo binocular microscope, and their identities determined using cultural, morphological, and descriptions in existing publications (Barnett. and Hunter, 1972; Booth, 1971; Webster, 1980).

Pathogenicity Test

Freshly harvested ripe african star apple fruits obtained from a tree at Moniya, Ibadan were surface sterilized by swabbing with 70% alcohol. A sterile cork-borer (4mm in diameter) was used to remove a tissue core from each of the surface-sterilized fruit. A second sterile cork -borer was used to cut discs of agar containing 3-day-old cultures of fungal mycelia of the isolates and used to inoculate the hole created by scooping out fruit tissue. The scooped out tissues were replaced to cover the inoculated portion of the fruit. The inoculated fruits were then enclosed in polythene bags containing moist cotton wool to maintain high relative humidity and incubated at 25⁰C in Gallenkamp incubators for 7 days. Four fruits were inoculated per isolate, while the control fruits were inoculated with sterile PDA agar discs. The extent of rot was determined by measuring the size of infection (mm). Wet mounts of hyphal/asexual structures obtained from these infected materials were stained with lactophenol in cotton blue and viewed under the compound microscope for the presence of the pathogen that was used in the inoculation.

Storage of the African star apple fruit

African star apple fruits collected as soon as they dropped naturally from the tree were kept in an oven-sterilized container lined with sterile filter paper and kept in the laboratory at a temperature of 26⁰C. Samples were observed for spoilage (deterioration) and nutrient composition for 10 days at an interval of every other day.

Nutrient Composition

Ten of the stored african star apple fruits were processed for nutrient analysis at 3 days interval for 9 days along with 10 freshly picked fruits. The fruits were kept in clean containers, de-seeded and weighed. The fleshy pulp was cut into pieces and dried in a hot air oven at 60⁰C for 3 days. The dried fruits pulp was ground into powder, and analyzed (in triplicates) for moisture, carbohydrate, ash, crude fibre, crude proteins, and crude fat according to AOAC (1984) procedure. Mineral analysis was also carried out according to standard AACC (1983) method.

RESULTS AND DISCUSSION

Eight fungi were isolated from the deteriorating african star apple fruits in Ibadan, South Western Nigeria. The fungi include *Botryodiplodia theobromae*, *Rhizopus stolonifer*, *Aspergillus niger*, *A. tamarii*, *A. flavus*, *Fusarium spp*, *Penicillium spp* and *Trichoderma spp*. However, *A. niger* was the most prevalent of all the isolates, followed by *Penicillium spp*, *A. flavus* while *B. theobromae* was the least encountered (Table 1). The stored fruits were found associated with fungal infections beginning from the 3rd day in storage. After about nine days in storage, deterioration in the fruit is manifested by shrinking and obvious fungal mycelia growth. The

predominant fungi on the 3rd day was *A. niger* and *R. stolonifer*, but on the 9th day all of the other eight fungi associated with the field infection with the exception of *Trichoderma spp* had colonised the deteriorating pulp (Table 2). *B. theobromae* which occurred least in stored african star apple fruit has been reported to be one of the most important fruit rot pathogen in southwestern Nigeria (Adisa, and Fajola, 1982), and has been observed to cause significant reduction in cashew yield (Olunloyo, 1979). All the isolates except *Trichoderma spp* were observed to be pathogenic to African star apple fruit (Table 2). *R. stolonifer* and *A. niger* which are usually present in the air (Webster, 1980), are probably secondary invaders.

Table 1. Incidence of occurrence and pathogenicity of fungi found associated with the African star apple fruits in Ibadan in 2000 and 2001.

Fungal isolates	Rate (%) of occurrence	Diameter of infection (mm)
<i>A. niger</i>	22	14
<i>A. flavus</i>	50	28
<i>Penicillium sp.</i>	65	14
<i>Fusarium sp.</i>	40	25
<i>R. stolonifer</i>	75	19
<i>A. tamarii</i>	28	25
<i>Trichoderma sp.</i>	16	6
<i>B. theobromae</i>	21	42

This study showed that 35% of the fruits picked were infected. These infected fruits when packed with non-infected caused increased deterioration of African star apple in transit and storage (Adebisi, 1997). The natural dropping of star apple fruits probably causes entry point for the fungi that were associated with fruit deterioration. It is also possible that insect vectors are involved in dissemination as reported by Adelaja (1997) indicating that fruit fly stings enhance the entry of *Colletotrichum spp* into African star apple fruits by their oviposition on the fruits.

Results of the nutrient analysis revealed that the freshly harvested African star apple fruits had crude protein contents (CP) of 8.75 %, carbohydrate content (CHO) of 29.6%, crude fat (CF) of 16.2% and moisture content (MC) of 42.1%. However, 9 days after harvesting, the CP, CHO and CF contents decreased to about 5.01%, 20.2% and 13.2%, respectively. The MC also decreased to 32.6 % within the same period of study (Table 3) . The nutrient analysis of the freshly harvested fruits revealed that Potassium (K), Calcium (Ca) and Sodium (Na) contents were 1.63%, 0.70% and 0.63 % respectively (Table 2). Nine days after harvesting, the K, Ca and Na contents of the fruits had

Table 2. The incidence of occurrence of fungal isolates on the African star apple fruits stored for nine days.

Fungi isolates	Days in storage/incidence of occurrence								
	1	2	3	4	5	6	7	8	9
<i>A. niger</i>			+	+	+	+	+	+	+
<i>A. flavus</i>				+	+	+	+	+	+
<i>Penicillium</i> sp.				+	+	+	+	+	+
<i>Fusarium</i> sp.					+	+	+	+	+
<i>R. stolonifer</i>			+	+	+	+	+	+	+
<i>A. tamari</i>					+	+	+	+	+
<i>Trichoderma</i> sp.									
<i>B. theobromae</i>						+	+	+	+

+ = present on the pulp

Table 3. Nutrient content of African star apple fruits in Ibadan, Southwestern Nigeria.

Days in storage	Moisture contents	% Crude protein	% Ether extract (fat)	% CHO	% K	% Na	% Ca
1 st Day	42.10	8.75	16.2	29.6	1.63	0.63	0.70
3 rd Day	39.6	8.39	16.0	29.2	1.88	0.63	0.70
6 th Day	34.4	6.27	14.6	23.4	2.50	1.13	0.70
9 th Day	32.6	5.01	13.2	20.2	2.75	1.25	0.72

Proximate composition (% dry matter)

increased to 2.75%, 1.25% and 0.75% respectively. However, fruit from the market had 28.2% carbohydrates, 8.02% protein, 15.8% fat and 39.1% moisture. Adelaja (1997) reported that the crude protein, carbohydrates, crude fat content of the African star apple fruits were 8.8%, 29.9% and 17.1% respectively.

It could be deduced from this study that, deterioration of the fruit by the pathogen might have led to an increase in the mineral contents and decrease in metabolic synthetates of the African star apple fruits. The industrial use of African star apple fruits in jam making has been experimentally demonstrated (Umelo, 1997) . Changes in nutrient composition caused by infection of the fruit will adversely affect the uses for jam and other food products. Because the fruit may become contaminated when picked from the forest floor, disinfecting before storage and use within three days of dropping/picking will prevent excessive infection of the pulp by fungal pathogens.

ACKNOWLEDGEMENTS

We are very grateful to the staff of the crop utilization laboratory unit of the Institute of Agricultural Research & Training (IAR& T) Moor plantation Ibadan for carrying out the proximate analysis of the African star apple. We also acknowledge the assistance of Prof. P. O. Oyekan of IAR&T Moor plantation Ibadan, Nigeria for reviewing the manuscript.

REFERENCES

- AACC (1983). Approved methods of the AACC, 3rd edition, American Association of Cereal Chemist, St. Paul, Minnesota.
- Adebisi AA (1997). Preliminary survey of post-harvest and marketing constraints of *Chrysophyllum albidum* (African Star Apple) in Nigeria In: Proceedings of a National workshop on the potentials of the star apple in Nigeria (eds) Denton OA, Ladipo DO, Adetoro MA, Sarumi MB, pp. 84-102.
- Adelaja BA (1997). Observations on the pests and diseases of *Chrysophyllum albidum* in Nigeria. In: Proceedings of a National workshop on the potentials of the star Apple in Nigeria (eds) Denton OA, Ladipo DO, Adetoro MA, Sarumi MB, pp. 117-121.
- Adewusi HA. (1997). The African Star Apple *Chrysophyllum albidum* indigenous knowledge IK from Ibadan, Southwestern Nigeria. In: Proceedings of a National workshop on the potentials of the star apple in Nigeria (eds) Denton OA, Ladipo DO, Adetoro MA, Sarumi MB, pp 25-33.
- Adisa SA (2000). Vitamin C, Protein and mineral contents of African Apple (*Chrysophyllum albidum*) In: Proceedings of the 18th annual conference of NIST (eds) Garba SA, Ijagbone IF, Iyagba AO Iyamu AO, Kilani AS, Ufaruna N, pp. 141-146.
- Adisa VA, Fajola AO (1982). Post harvest fruit rots of three species of citrus in southwestern Nigeria. Indian Phytopathol. 35:595-603.
- AOAC (1984). Association of Official Analytical Chemist, Official Method of Analysis 14th Edition, Washington, DC.
- Aserjo CF (1946). The high ascorbic acid content of the West Indian Cherry. Science 103: 219.
- Bada SO (1997). Preliminary information on the ecology of *Chrysophyllum albidum* G. Don, in west and central Africa In: Proceedings of a National workshop on the potentials of the star Apple in Nigeria (eds) Denton OA, Ladipo DO, Adetoro MA, Sarumi MB, pp. 16-25.
- Barnett HL, Hunter BB (1972). Illustrated Genera of Imperfect Fungi. Mineapolis: Burgess Publishing Company, Minneapolis MN 241 p.
- Booth C (1971). The genus *Fusarium*: Laboratory Guide to the Identification of major species: by the Commonwealth Mycological Institute. Kew, Surrey UK 231p.

- Cenrad (1999). Centre for Environmental Renewable Natural Resources Management, Research and Development, Jericho, Ibadan. Publication No CEN. 011/1999 85.
- Keay RWJ (1989). Trees of Nigeria. A revised version of Nigerian trees (Vol 1 and 2) (eds) Keay RWJ, Onoche CFA, Stanfield DP. Clarendon Press, Oxford. 476p.
- Okafor JC, Fernandes ECM (1987) Compound farms of south-east Nigeria: a predominant agroforestry home system with crops and small livestock. *Agroforestry Systems* 5:153-168.
- Olunloyo OA (1979). The role of fruit flies in rot disease of cashew apples in the plantation. Annual Rept. Cocoa Research Institute of Nigeria, Ibadan.pp. 101-105.
- Umelo R (1997). Potentials for utilization of African star apple (*Chrysophillum albidum*) for Jam making in Nigeria. In: proceedings of a National workshop on the potentials of the star Apple in Nigeria (eds) Denton OA., Ladipo DO , Adetoro MA, Sarumi MB, p. 103.
- Webster J (1980). Introduction to fungi, 2nd ed. Cambridge University Press, 242p.