

Full Length Research Paper

# Evaluation of methicillin resistance *Staphylococcus aureus* isolated from patients in Golestan province-north of Iran

Hamid Vaez, Alijan Tabaraei, Abdolvahab Moradi and Ezzat A. Ghaemi\*

Department of Microbiology, School of medicine, Infectious diseases research centre, Golestan University of Medical Science, Gorgan, Iran.

Accepted 17 January, 2022

Methicillin resistant *Staphylococcus aureus* (MRSA) is a main cause of nosocomial infections with consequence of increasing hospitalization, costs of treatment and rate of mortality. This study was aimed to demonstrate distribution of MRSA strains and their antibiotic resistance pattern. In this descriptive study, 185 clinical isolates of *S. aureus* that were collected from different infections during September 2008 to 2009 were tested by polymerase chain reaction (PCR) and micro dilution broth. All the MRSA and methicillin sensitive *Staphylococcus aureus* (MSSA) isolates were tested for antibiotic resistance pattern by disk diffusion method with 14 different antibiotics. Data were entered in SPSS software version 16 and analyzed by chi-square test. P value of <0.05 was considered significant. Of 185 tested *S. aureus*, 67(36.2%) strains were MRSA, which demonstrated 100% resistance to Penicillin, Ampicillin and CO-Amoxyclav and -80, 96.2 and 75% resistance to Cephotoxime, Nalidixic Acid and erythromycin, respectively. All *S. aureus* isolates was sensitive to vancomycin. All isolates with minimum inhibitory concentration (MIC) >8 µg/ml were *mecA* positive. MRSA is spreading worldwide with increasing levels of resistance, and accurate and early detection of these strains is encouraged.

**Key words:** *Staphylococcus aureus*, methicillin resistant *Staphylococcus aureus*, methicillin sensitive *Staphylococcus aureus*, minimum inhibitory concentration, *mecA*

## INTRODUCTION

Infections caused by *Staphylococcus aureus* were the major cause of mortality in different communities, before the discovery of Penicillin by Alexander Fleming (Srifuengfung, 1994). *S. aureus*'s infections, especially caused by methicillin-resistant *Staphylococcus aureus* (MRSA), are emerging as a major public health problem. MRSA has become a leading cause of nosocomial infections worldwide, since the first isolate was detected in 1961 in England (Srifuengfung, 1994; Zhang, 2001). *S.aureus* has a protein in its cell wall called penicillin binding protein (PBP), with trans-peptidase activity, play a key role in cell wall synthesis and are the target for -lactam antibiotics. In addition MRSA strains produce a

modified PBP called PBP2a or PBP2<sup>f</sup> with low affinity for -lactam antibiotics (Hiramatsu, 2001; Zhang, 2005). Resistance to methicillin mediated by *mecA* gene, responsible for production of PBP2a, which have been observed in MRSA. *mecA* located on a region of chromosome called SCCmec. Five different types of SCCmec have been characterized as I, II, III, IV, V (Hiramatsu, 2001; Zhang, 2005). *mecA* expression in *invitro* condition was variable and depends on different factors such as temperature, cation concentration, incubation period and pH (Forbes, 2007). MRSA showed resistance to other -lactam antibiotics such as Oxacillin and Nafcillin. MRSA is the main causes of nosocomial infections in the worldwide. Power of morbidity and simultaneously resistance to other antibiotics in MRSA strains is higher than methicillin sensitive *S. aureus* (MSSA). Regarding to increased prevalence of these strains in recent years and pathogenesis of *S. aureus*,

\*Corresponding author. E-mail: [eghaemi@yahoo.com](mailto:eghaemi@yahoo.com). Tel: (+) 989113711770.

accurate and early identification of these strains is very important (Chambers, 199; Gradie, 2001). MRSA strains can grow in the presence of 16 µg/ml or more of methicillin while sensitive strains are inhibited (Lewis, 2008; Gradie, 2001). Our recent investigation on healthcare staff showed 12.5% MRSA among *S. aureus* carriers (Unpublished). The aim of the present study was to demonstrate prevalence of MRSA by polymerase chain reaction (PCR) and micro dilution broth methods as well as their antibiotic resistance pattern.

## MATERIALS AND METHODS

185 clinical samples were taken from various sites of infection including blood, wound, sputum, urine and others collected from five major hospitals in Golestan province named as: Taleghani, 5th Azar, Dezyani, Shohada and Shahid Motahari during June, 2008 and May, 2009. *S. aureus* identification performed based on standard tests such as Gram stain, catalase, DNase, growth on manitol salt agar, slide and tube coagulase (Forbes, 2007). Demographic data for all patients such as sex, age and site of infections were recorded. Methicillin resistance and molecular detection of *mecA* gene and susceptibility of strains to other antibiotics were performed as below.

### Minimum inhibitory concentration (MIC)

MIC of methicillin was determined by micro dilution broth method, using Muller Hinton broth supplemented with 2% NaCl. Bacterium inoculation of  $5 \times 10^5$  and incubation at 35°C for 24 h was done according to Clinical and Laboratory Standards (CLSI) guide lines (Lewis, 2008; Gradie, 2001).

### Disk diffusion test

Resistance to other antibiotics evaluated by Kerby bauer disk diffusion method based on CLSI guide lines (Brown, 2005; Kohner, 1999). Antibiotic disks (Himedia-India) that were used in this study is listed as: Nalidixic Acid (30 g)-Ceftriaxone (30 g)-Penicillin(10 unit)- Sulphamethoxazol-Trimethoprim (50 g)-Ciprofloxacin(5 g)- Chloramphenicol(30 g)-Cefotaxime(30 g)-Ampicillin(10 g)- Vancomycin(30 g)Gentamicin (10 g)-Erythromycin(15 g)-Tetracycline (30 g)-Co-amoxyclave(10 g) and Ceftioxin (30 g).

### Molecular detection of *mecA* gene

DNA extraction was performed by lysing *S. aureus* as following procedure: one milliliter of overnight growing bacteria in BHI broth transferred to 2 ml sterile microtube and centrifuged at 8000 rpm. 300 µl TE (tris EDTA pH=8, sigma- Germany) buffer was added to pellet and vortexed vigorously. 25 µl lysozyme (10 mg/ml) was added to suspension and incubated at 37°C for 30 min. Re-incubation at 37 °C for 1 h was done by adding 300 µl sarcosyl 2% and 30 µl proteinase K (10 mg/ml). DNA were extracted and purified by phenol-chloroform-isoamylalcohol (25:24:1) and cold pure ethanol method.

PCR for detection of *mecA* gene was carried out by using primers as below: *mecA*-F: 5'-AAAATCGATGGTAAAGGTTGGC-3' and *mecA*-R: AGTTCTGCAGTACCGGATTTGC-3' (Cinnagen, Iran) for coding a 533 bp *mecA*. 50 µl PCR reaction mixture were made consist of 10 pmol of each primers, 200 µM Dntp (Roche-Germany), 2.5 µl (50 mM mgcl<sub>2</sub>), 0.5 µl Taq pol (2.5 u) (Roche-Germany), 5 µl PCR buffer 10x (Roche-Germany) and 5 µl DNA-template. *S. aureus* COL strain was used as a positive control. Cycling condition

performed in 40 cycles including, denaturation at 95°C for 30 s, annealing at 55°C for 30 s, extension at 72°C for 1 min and final extension at 72°C for 5 min (Nimmo, 2003; Louie, 2000). PCR products were visualized on 1.7% agarose gel with ethidium bromide dye under UV transilluminator. Amplicon of 533 bp were consistent with *mecA* gene amplification.

## RESULTS

In this study 185 *S. aureus* were isolated from patients. Median age of patients were  $29.2 \pm 28$  years (from less than one year to 84 years) that 96 (52%) of them were male. The clinical specimens, 57 (30.8%), 49 (26.5%), 42 (22.7%), 20 (10.8) and 17 (9.2%) of them were isolated from urine, wound, blood, sputum and other specimens such as abscess respectively.

Based on MIC value, 67(36.2%) of isolates were MRSA (MIC 16 µg/ml). In 5 (2.7%) strains MIC 256 µg/ml were seen with highly resistance to methicillin. In 82 (44.3%) MIC was between 4 and 8 µg/ml. Only 36 (19.5%) of isolates had MIC less than 2 µg/ml and showed full sensitivity to methicillin (Table 1).

We have found *mecA* gene in 65 (35.1%) isolate of *S. aureus* by PCR method (Figure 1). None of *mecA* containing isolates were shown MIC 2 µg/ml and in 8 (6.7%) *S. aureus* isolates without *mecA* gene with MIC 16 µg/ml were seen (Table 1).

Rate of MRSA strains isolation in wound and sputum were higher than other samples with 42.9% and 40% respectively (Figure 2).


According to the sex, rate of MRSA and MSSA strains among men and women were 37 (38.5%) and 30 (33.7%) respectively, showing significant relation ( $P < 0.05$ ). Frequency of antibiotic resistance in MRSA strains was significantly higher than MSSA strains ( $P < 0.05$ ). This high resistance was seen more in MRSA strains to Cephotoxim, Erythromycin, Ceftriaxon and Gentamycin in compare with MSSA isolates (Table 2). Our finding showed that multi drug resistant property among MRSA strains was high, so that highest level of resistance was observed with Penicillin, Ampicillin, Nalidixic Acid, Co-amoxyclave, Ceftriaxone and Erythromycin. In MSSA strains resistance to Penicillin, Ampicillin and Co-amoxyclave was high in comparison with other antibiotics (Table 2). All isolates were sensitive to vancomycin.

## DISCUSSION


Our study demonstrated that 36.2% of *S. aureus* isolates were MRSA, this prevalence are similar to some other studies performed in various region in Iran such as Tehran as well as neighbor countries like Saudi Arabia and Kuwait, and also European country, France and American nation, Brazil (Fatholahzadeh, 2008; Mdani, 2001; Udo, 2008; Oteo, 2004). Instead of the abundance of MRSA isolates across the world are not the same; the most MRSA isolates have been reported from Taiwan, Canada and

**Table 1.** Comparison of MIC and PCR results in MRSA detection.

PCR result \ MIC	MIC			Total (%)
	2 (%)	4-8 (%)	16 (%)	
mecA+	0 (0)	6 (9.2)	59 (31.9)	65 (35.1)
mecA-	36 (30)	76 (63.3)	8 (6.7)	120 (64.9)
Total	36 (19.5)	82 (44.3)	67 (36.2)	185 (100)



**Figure 1.** Gel image of representative PCR *mecA* gene products –line1-marker (100 bp), line 2 negative control, line 3 positive control (533 bp band) and line4 to 8 patient specimens.



**Figure 2.** Distribution of *mecA* positive MRSA in clinical samples.

**Table 2.** Antibiotics resistance pattern in MSSA and MRSA.

Isolate Antibiotic	MSSA (120)(65%)	MRSA (65)(35%)	Total (185)(100%)
Penicillin	115(95.6)	65(100)	180(97.3)
Ampicillin	115(95.6)	65(100)	180(97.3)
Co-amoxyclave	93(95.3)	63(97)	156(84.3)
Nalidixic Acid	40(33.3)	58(89.2)	98(53)
Cefotaxime	3(2.5)	48(73.8)	51(27.6)
Erythromycin	8(6.7)	46(70.8)	54(29.2)
Ceftriaxone	3(2.5)	44(67.7)	47(25.4)
Trimethoprim	16(13.3)	41(63.1)	57(30.8)
Ciprofloxacin	4(3.3)	38(58.5)	42(22.7)
Gentamicin	1(0.8)	37(56.9)	38(20.5)
Tetracycline	26(21.7)	32(49.2)	58(31.4)
Imipenem	1(0.8)	15(23.1)	16(8.6)
Chloramphenicol	8(6.7)	12(18.5)	20(16.7)
Vancomycin	0(0)	0(0)	0(0)

Australia as much as 66 to 77% (Fatholahzadeh, 2008; Hsueh, 2004; Merlino, 2002; Arbique, 2001). As our recent study showed that 24% of healthcare staff were *S. aureus* carriers. Of these 12.5% were MRSA showing 3% total prevalence of MRSA (unpublished). Also recent report about rate of MRSA in nosocomial infections in Isfahan, Iran showed that 67.2% of isolates were MRSA (Khorvash, 2008). Different epidemiological factors such as geographical, health system capability in running infection control program has role in variability of prevalence of MRSA. Most isolates of MRSA were observed in wound specimens (43%). Instead of many similar and independent studies that is not showing any relation between sex, age, site of infection and rate of MRSA. Only a few report has been shown increasing the rate of MRSA in elder people aged 64 and more significantly (Fatholahzadeh, 2008; Hsueh, 2004; Merlino, 2002; Arbique, 2001; Waness, 2010). It has been clear that age is a risk factor because of its role in long term hospitalization; lose of immunity and longer antibiotic therapy (Waness, 2010).

Multi drug resistance MRSA strains are made difficult treatment of MRSA infections. In our study 100% resistance to Penicillin, Co-amoxyclave and Ampicillin, and more than 70% resistance to Cefotaxime, Erythromycin, Ceftriaxone and Nalidixic Acid were showed. In compare, rate of MSSA sensitivity to other antibiotics was considerable. It is comparable with other study in our country showing such results (Fatholahzadeh, 2008; Khorvash, 2008).

We have found *mecA* gene in 65 (35.1%) isolates of *S. aureus* by PCR method. None of *mecA* containing isolates was shown MIC 2 µg/ml whilst in 8 (6.7%) *S. aureus* isolates without *mecA* gene MIC 16 µg/ml were seen (Table 1). We considered MIC as the reference or "gold standard" method for establishing the sensitivity and

specificity of each of PCR technique studied. We have obtained specificity and sensitivity of 94.89 and 88% for PCR method respectively.

It is determined that 82 (44.3%) of isolates showed MIC between 4 and 8 µg/ml and 76 (63.3%) of them were *mecA* negative and categorized as border line in susceptibility to methicillin named as BORSA. High rate of BORSA strains is important in drug resistancy and it has been suggested that under pressure of antibiotics this isolates can probably shift to fully resistant MRSA (Mathews, 2010; Khorvash, 2008; Nelson, 2006).

In conclusion MRSA is spreading worldwide, growing epidemic and increasingly claiming victims. Accurate and early detection of these strains in hospitals and community is encouraged.

## REFERENCES

- Arbique J, Forward K, Haldane D, Davidson R (2001). Comparison of the *velogene* MRSA id system for rapid identification of methicillin resistant *Staphylococcus aureus*. *Diag. Microbial. Infect. Dis.*, 40: 5-10.
- Brown FJ, Edward D, Hawkey P, Morrison D, Ridgway G, Towner K, Wren MWD (2005). Guideline for the laboratory diagnosis and susceptibility testing of Methicillin Resistant *Staphylococcus aureus*. *J. Antimicrob. Chemother.*, 56: 1000-1018.
- Chambers HF (1997). Methicillin Resistance in Staphylococci :molecular and biochemical basis and clinical implication. *Clin. Microbiol. Rev.*, 10(4): 781-791.
- Fatholahzadeh B, Emameini M, Gilbert G, Udo E, Aligholi M, Modarressi MH (2008). Staphylococcal cassette chromosome *mec* (SCCmec) analysis and antimicrobial susceptibility patterns of methicillin –resistant *Staphylococcus aureus* (MRSA) isolets in Tehran ,iran. *Microbial Drug Resis.*, 14(3): 217-222.
- Forbes BA, Sahn DF, Weissfeld AS (2007). *bailey&scotts Diagnostic microbiology*. 12<sup>th</sup> ed. USA; Elsevier, pp. 172-213.
- Gradie E, Valera L, Aleksunes S, Bonner D, Fung G (2001). Correlation between genotype and phenotypic categorization of staphylococci based on methicillin susceptibility and resistance. *J. Clin. Microbiol.*, 39(8): 2961-2963.
- Hiramatsu K (2001). Vancomycin Resistant *Staphylococcus aureus*: A

- new model of antibiotic resistance. *Lancet Infect Dis.*, 1: 147-154.
- Hsueh PR, Teng IG, Chen WH, Pan HJ, Chen MI, Chang SC, Luh KT, Lin FY (2004). Increasing prevalence of methicillin resistant *Staphylococcus aureus* causing nosocomial infections at a university hospital in Taiwan from 1986 to 2001. *Antimicrob. Agents Chemother.*, 48(4): 1361-1364.
- Khorvash F, Mostafavizadeh K, Mobasherizadeh S (2008). Frequency of *mecA* Gene and Borderline Oxacillin Resistant *Staphylococcus aureus* in Nosocomial Acquired Methicillin Resistance *Staphylococcus aureus* Infections. *Pak. J. Biol. Sci.*, 11(9): 1282-1285.
- Kohner P, Uhl J, Kolbert C, Persing D, Cockerill F (1999). Comparison of susceptibility testing method with *mecA* gene analysis for determining Oxacillin (Methicillin) resistance in clinical isolate of *Staphylococcus aureus* and coagulase negative *Staphylococcus* spp. *J. Clin. Microbiol.*, 37(9): 2952-2961.
- Lewis K, Salyers AA (2008). Bacterial resistance to antimicrobials. CRC Press, Taylor and Francis Group.
- Louie L, Matsumura SO, Choi E, Simor E (2000). Evaluation of three rapid method for detection of methicillin resistance in *Staphylococcus aureus*. *J. Clin. Micro. Biol.*, 38(6): 2170-2173.
- Mathews AA, Thomas M, Appalaraju B, Jayalakshmi J (2010). Evaluation and comparison of tests to detect methicillin resistant *S. aureus*; 53(1): 79-82.
- Mdani TA, AL-Abdollah NA, AL-Sanousi AA (2001). Methicillin –resistant *Staphylococcus aureus* in two tertiary-care centers in jaddah Saudi Arabia. *Control Hosp. Epidemiol.*, 22: 211-216.
- Merlino J, Watson J, Rose B (2002). Detection and expression of methicillin resistant *Staphylococcus aureus* in central Sydney, Australia. *J. Clin. Microb. Chemother.*, 49: 793-801.
- Nelson L, Cockram CS, Lui G, Lam R, Lam E, Lai R, Ip M (2006). *Emerg. Infect. Dis.*, 21(1): 172-174.
- Nimmo GR, Bell GM, Mitchel D, Gosbell IA, Perman JW, Turnidge JD (2003). Anti microbial resistance in *staphylococcus aureus* in Australian teaching hospital, 1989-1999. *Microb. Drug Resist.*, 9(2): 155-159.
- Oteo J, Baquero F, Vindel A, Campos J (2004). Antibiotic resistance in 3113 blood isolate of *Staphylococcus aureus* in 40 Spanish hospitals participating in the European antimicrobial resistance surveillance system. *J. Antimicrob. Chemother.*, 53: 1033-1038.
- Srifuengfung S (1994). Methicillin-resistant *Staphylococcus aureus* (MRSA): Review of problems and comments. *J. Infect. Dis. Antimicrob. Agents*, 11: 141-147.
- Udo EE, AL-Sweih R, Dahr TS, Dimitrov EM, Mokaddas M, Johny I A, Johny M, Al-Obaid IA, Gomaa HH, Mobasher LA, Rotimi VO, Al-Asar A (2008). Surveillance of antibacterial resistance in *Staphylococcus aureus* isolated in Kuwaiti hospitals. *Med. Prink Pract.*, 17: 71-75.
- Waness A (2010). Revisiting methicillin-resistant *Staphylococcus aureus* infections, 2(1): 49-56.
- Zhang HZ (2001). A proteolytic transmembrane signaling pathway and resistance to beta lactams in *Staphylococci*. *Science*, 291: 1962-1965.
- Zhang K, McClure JA, Elsayed S, Louie T, Conly JM (2005). Novel multiplex PCR assay for characterization and concomitant subtyping of *Staphylococcal* cassette chromosome *mec* types I to V in methicillin resistant *Staphylococcus aureus*. *J. Clin. Microbiol.*, 43(10): 5026-5033.