

*Full Length Research Paper*

# Salinity tolerance of rice lines related to endogenous abscisic acid (ABA) level synthesis under stress

Saeed Saeedipour

Department of Agronomy, Shoushtar Branch, Islamic Azad University, Iran. E-mail: saeeds79@gmail.com.

Accepted 26 October, 2020

As the plant hormone abscisic acid (ABA) is involved in responses to salinity stress. We tested its putative relationship with the degree of tolerance to this abiotic stress. For this purpose we have examined the responses of sensitive (IR29) and tolerant (IR651) varieties of indica rice (*Oryza sativa* L.) to a range of salinity (0 (control) and 100 mM NaCl). Shoot and root dry weight was reduced and leaf Na concentration increased in response to salinity for both cultivars with a bigger extent in sensitive. Tolerance of IR29 to saline stress was generally improved by ABA treatment and leaves Na content reduced to their respective control treatment. This ABA effect was evident in IR29 with low tolerance, as their ability to recover from stress increased up to seven fold. Independent of the saline treatment, the absolute endogenous leaf ABA content in sensitive variety was significantly more than tolerant one. However, upon stress, the increase in endogenous ABA synthesis was higher in tolerant than in sensitive varieties. These data together with those obtained by using Fluridone, an inhibitor of ABA synthesis, suggested first, there was differential sensitivity to ABA in the tolerant and sensitive leaves cultivars and enhanced concentrations at tolerant levels acted primarily to maintain root and shoot growth salt stress and second, the differences in the level of tolerance to saline stress is related to their different capacity of ABA synthesis under stress conditions.

**Key words:** Abscisic acid, fluridone, *Oryza sativa* L., salinity, stress tolerance.

## INTRODUCTION

Salinization of arable land is increasing, and could possibly have devastating global effects. Over 6% of the world's land and 20% of the world's irrigated land are currently affected by salinity (Rhoades et al., 1992; Munns, 2005). In such soils, NaCl concentrations typically exceed 40 mM, and much higher values are frequently found (Munns, 2005). Enhancing membrane permeability in salt stressed plants has been well documented (Hasegawa et al., 2000), resulting in Na<sup>+</sup> accumulation and increasing Na<sup>+</sup>/K<sup>+</sup>, conversely reducing K<sup>+</sup> in many plant species that is rice (Dionisio-Sese and Tobita, 1998) green bean (Yasar et al., 2006), winter wheat (Zheng et al., 2008), umbu plant (Da Silva et al., 2008). From some previous studies, the results showed that rice growth and development are reduced when exposed to the salinity stress (Morsy et al., 2007; Cha-um et al., 2007; Dionisio-Sese and Tobita, 1998; Shannon et al., 1998). However, the response of rice to salt stress depends on its salt tolerance abilities. During the last two decades, it has been well established that abscisic acid

(ABA) is a vital cellular signal that mediates the expression of a number of salt and water deficit-responsive genes. The direct relation between stress tolerance and increased levels of ABA does not always exist. Thus, in barley and wheat changes in endogenous ABA levels have been reported to be unrelated to freezing tolerance or to cold hardening (Kadlecová et al., 2000; Faltusová-Kadlecová et al., 2002). Among the comparisons between stress-tolerant and stress sensitive cultivars within the same species, some reports have shown higher ABA levels in tolerant cultivars, although the differences appeared in either unstressed, non-acclimated cultivars (Bravo et al., 1998) or following the stress (Chen et al., 2002; Moons et al., 1995). In rice, ABA contents increased in a semi salt-tolerant variety but only marginal changes were measured in salt-tolerant and salt sensitive ones (Bohra et al., 1995). Exogenous ABA treatments prior to subjecting the plants or tissues to adverse conditions have been reported to improve tolerance to cold temperatures (Borrmann and Jansson,

1980; Duncan and Widholm, 1987), osmotic stress (Nayyar and Kaushal, 2002; Nayyar and Walia, 2003) and rice (Asch et al., 1995). In the current study we tested the important role of ABA in tolerance to abiotic stresses, and also provide new insights into the mechanism of improved salt tolerance by the application of exogenous ABA on crops.

## MATERIALS AND METHODS

### Plant material and growth conditions

Two rice cultivars different in tolerance of salt stress during the vegetative stages (Moradi et al., 2003) were selected for this investigation. IR65192-4B-10-3 (abbreviated as IR651) is breeding lines tolerant of salt stress at both the seedling and reproductive stages, and IR29 is a cultivar sensitive to salt stress during both stages and is commonly used as a sensitive check in breeding nurseries. The experiment was carried out on the Agricultural Biotechnology Research Institute of Iran (48°20 N; 31°41E; 20 m above sea level), in the growing season of 2009 to 2010. This experiment was conducted in a growth cabinet with light intensity of about  $700 \mu\text{mol m}^{-2} \text{s}^{-1}$  and 12 h duration, 70% relative humidity and 29/22°C day/night temperature. Pre-germinated seeds of each cultivar were sown in holes made on Styrofoam sheets with a nylon net at bottom (Gregorio et al., 1997). Two seeds were sown per hole, with 50 holes per entry. The sheets were first floated on distilled water in 11 L plastic trays for 3 day, after which a nutrient solution (Yoshida et al., 1976) was used until the plants were 21 days old. A control treatment (nutrient solution) and the level of salt stress,  $12 \text{ dsm}^{-1}$  (100 mM) using NaCl was introduced thereafter. Four hours after imposing the salt treatment, half of the plants from both cultivars were exposed to spraying of Fluridone (50  $\mu\text{M}$ ) or ABA (20  $\mu\text{M}$ ). The youngest fully expanded leaf (leaves that had completed petioles), harvested at 0 (immediately before starting the salinity), 4, 12, 24, 48, 96 and 168 h after the start of the salt treatments for ABA measurements. Shoot and root dry matter,  $\text{K}^+$  and  $\text{Na}^+$  concentration of youngest fully expanded leaf were measured at the end of this experiment.

The experiment was designed as 2x2 (Cultivars x Salinity) factorials in Completely Randomized Design (CRD) with three replicates. The mean in each treatment was compared by Duncan's New Multiple Range Test (DMRT) at  $p < 0.01$  and analyzed by SPSS software (SPSS for Windows, SPSS Inc., Chicago, Illinois, USA). The culture solution was renewed once with the pH adjusted daily to 5.5 by adding either NaOH or HCl.

### Sodium and potassium measurements

Dried samples were ground to a fine powder. About 0.1 g was transferred to a test tube containing 10 ml 0.2 N acetic acid. Then test tubes heated in a water bath at 80°C for 2 h. The extracted tissue was cooled at room temperature and left overnight and then filtered by using Whatman filter paper number 42. Sodium and potassium concentrations were determined by using an atomic absorption spectrometer (Perkins Elmer, Norwalk, CT, USA).

### Leaf gas exchange measurement

Stomatal conductance ( $g_s$ ) was made on fully expanded youngest leaf of each plant using an open system LCA-4ADS portable infrared gas analyzer (Analytical Development Company, Hoddesdon, England). Measurements were performed with the

following specifications/adjustment: molar flow of air per unit leaf area  $403.3 \text{ mmol m}^{-2} \text{s}^{-1}$ , atmospheric pressure 99.9 kPa, water vapor pressure into chamber ranged from 600 to 890 Pa, Par at leaf surface was maximum up to  $1711 \mu\text{mol m}^{-2} \text{s}^{-1}$ , temperature of leaf ranged from 28.4 to 32.4°C, ambient temperature ranged from 22.4 to 27.9°C, and ambient  $\text{CO}_2$  concentration  $352 \mu\text{mol mol}^{-1}$ .

### Measurement of endogenous ABA levels

Endogenous ABA analyses were performed as described in Gómez-Cadenas et al. (2002), frozen rice leaves of 1.5 g were finely ground in liquid nitrogen and 10 ml of 80% methanol was added together with 0.01 g of ascorbic acid and 0.01 g polyvinylpyrrolidone (PVP) to prevent oxidation reactions during extraction. The homogenate was stirred overnight at 4°C. After centrifugation (4000 g, 15 min), the supernatant was recovered and adjusted to pH 8.0. The aqueous methanol was evaporated under reduced pressure at 35°C. The residue was dissolved in 5 ml of water. Then it was frozen and thawed for three cycles. After centrifugation (4000 g, 15 min), the supernatant was recovered and adjusted to pH 2.5 and partitioned against ethyl acetate. Then the solution with the free ABA in ethyl acetate was collected. This process was repeated thrice. After that, the collection was adjusted to pH 8.0 and dried. The resulting dried precipitate was dissolved in 1 ml of 3% methanol containing 0.1 M acetic acid, and was filtered through a 0.45 mm membrane filter. The extract (100  $\mu\text{l}$ ) was automatically injected and processed by high performance liquid chromatography (HPLC) (Agilent 1100 Series, USA) equipped with a reverse phase column (4.6 x 250 mm Diamonsic C18, 5  $\mu\text{m}$ ). It was eluted with a linear gradient of methanol (3 to 97%), containing 0.01% acetic acid, at a flow rate of 4 ml/min. The detection was run at 260 nm with a diode array detector. The retention time of ABA was 36.4 min and shifted 0.1 to 0.5 min. Quantification was obtained by comparing the peak areas with those of known amounts of ABA.

## RESULTS

### Effect of stress on K and Na concentration with impaired ability for ABA synthesis

A different approach was used to study the implication of ABA in the mechanism of stress-tolerance. This experiment was performed with Fluridone treated, an inhibitor of ABA biosynthesis. Irrespective of Fluridone application no statistically significant differences were measured between tolerance and sensitive cultivars in Na concentration in control treatment of both cultivars investigated (Figure 2A and B). Salinity treatment increased 2 fold the Na level in IR29 but became unchanged in IR651 with their respective control (Figure 2A). The Na accumulation in young leaves of both cultivars became more pronounced upon applied Fluridone in salinity treatment (Figure 2A and B) and cause to rapid raise of Na, 7.3 and 2.8 fold in sensitive and tolerant cultivars respectively, (contrast to controls treatments). Fluridone had a dramatic effect on their response to stress in both cultivars.

Thus, when plants of both cultivars were sprayed with Fluridone, they showed higher sensitivity to a subsequent salinity stress than stressed plants that had not been

**Table 1.** Stomatal conductance ( $\mu\text{mol m}^{-2} \text{s}^{-1}$ ) of rice varieties under control (0 mM NaCl = 1.56 ds.m<sup>-1</sup>) and salinity (100 mM NaCl = 12 ds.m<sup>-1</sup>) conditions.

Variety	Salinity	Sampling time					
		12	24	48	72	96	120
IR29	0	0.5±0.01	0.05±0.01	0.5±0.02	0.5±0.05	0.5±0.03	0.6±0.03
	12	0.1±0.02	0.2±0.02	0.2±0.04	0.2±0.02	0.2±0.01	0.2±0.03
IR651	0	1.2±0.03	1.1±0.1	1.4±0.3	1.4±0.2	1.3±0.1	1.4±0.3
	12	0.3±0.03	0.8±0.04	0.7±0.1	0.7±0.1	0.8±0.1	0.7±0.04

Each value is a mean of three replicates ± se.

sprayed with the inhibitor (Figure 2A and B). Salt stress induced an approximate 30% increase in endogenous leaf ABA content of tolerant cultivar, while spraying with 50  $\mu\text{M}$  Fluridone during the salinity treatment not only suppressed this increase but declined it more than 30% (Figure 3).

In a further experiment, plants from tolerant and sensitive cultivars were treated with exogenous ABA and subjected to salinity (Figure 2A and B). When plants were treated with ABA (20  $\mu\text{M}$ ), leaf tolerant cultivar did not change significantly their Na level response to the subsequent stress. By comparison, in IR29 plants application of exogenous ABA, lead to dramatically reduction the Na level and dropped the value to normal treatment (Figure 2B). It has to be pointed out that the effect of ABA was totally counteracted with the inhibitor. Regardless of the salinity, ABA or Fluridone treatments, a very similar changing pattern was observed for K concentration in leaves of both cultivars and no significant differences was observed (Figure 2C and D).

### Effect of stress on endogenous ABA levels


Endogenous ABA levels of tolerant and sensitive of both cultivars were determined in order to check whether the different degree of tolerance to stress was correlated with hormone contents. Irrespective of salinity treatment, leaves ABA concentration in IR29 was higher than IR651. Endogenous ABA levels of leaves under non stress conditions varied in the range of 13.4 to 14.10  $\mu\text{g g}^{-1}$  Fwt in IR29. Compare to sensitive cultivar, the tolerant cultivar had a lower ABA concentration, and the values varied in the range of 6 to 9.5  $\mu\text{g g}^{-1}$  Fwt in control treatment. Leaves ABA content of stressed IR651 plants was significantly bigger than that of control plant throughout sampling time and the differences became more pronounced and achieved a maximum value by end of experiment (Figure 3). Opposite to tolerant cultivar, leaves ABA levels in sensitive cultivar increased slightly but did not reach the level of statistical significance at the end of experiment compare to their respective control treatment (Figure 3).

### Stomatal conductance and biomass accumulation


Stomatal conductance under no stress condition was two fold in IR651 than IR29, and the differences keep constant throughout of experiment (Table 1). Salinity was significantly lowered the stomatal conductance, the rate of stomatal conductance of IR29 and IR651 plants was reduced by 50% (Table 1). Result regarding plant biomass showed inverse relationship with salinity. A general trend of decrease in dry weight of plant with salinity was noted in both cultivars. However, shoot and root weight reduction was largest in the susceptible variety compare to tolerant variety (Figure 1B and D). Fluridone application lead to drastically decline in root and shoot dry weight of both cultivars (Figure 1A and C). Regardless of salinity, shoot and root dry weight in IR651 was two fold than IR29.

### DISCUSSION


The plant hormone abscisic acid has been implicated in plant responses to abiotic factors causing water stress. In most studies, elevated ABA levels prior to stress, or as a first step in the response to adverse conditions, are involved in tolerance mechanisms. Under this rationale, it is to be expected that ABA increased in leaves of tolerant variety have a positive role in alleviating stress effects. In the present work differences in ABA contents were observed between sensitive and tolerant of the two varieties studied, and maybe we can ascribe their different response to stress to a different endogenous hormone level under control, non-stress conditions (Figure 2) when plants were subjected to stress we could detect differences in ABA synthesis in varieties with respect to non-stressed controls (Figure 3). Tolerant variety seems to have a higher ability for ABA synthesis than sensitive. This agrees with reported results comparing sensitive/tolerant cultivars of different species (Chen et al., 2002; Lee et al., 1993). Moons et al. (1995) reported that although salt-tolerant rice cultivars showed higher increases of ABA than the sensitive cultivars, there was not a direct correlation between



**Figure 1.** Shoot and root dry weight of two rice cultivars during the seedling stage under control and salt stress of  $12 \text{ dsm}^{-1}$ , with (A and C) or without (B and D) spraying Fluridone ( $50 \mu\text{M}$ ). The data are mean values of three replications with three subsamples per replication, and vertical bars are  $\text{LSD}_{0.01}$ .



**Figure 2.** Na and K concentration of youngest full expanded leaf of two rice cultivars during the seedling stage under control and salt stress of  $12 \text{ dsm}^{-1}$ , with spraying Fluridone ( $50 \mu\text{M}$ ) or ABA ( $20 \mu\text{M}$ ). The data are mean values of three replications with three subsamples per replication, and vertical bars are  $\text{LSD}_{0.01}$ .



**Figure 3.** ABA measured on the youngest fully expanded leaf of two rice cultivars during the seedling stage under control and salt stress of 12 dsm<sup>-1</sup>. The data are mean values of three replications with three measurements per replication, and vertical bars represent ±SE of the mean (n=3) where these exceed the size of the symbol. The values on the x-axis represent of hours after the start of salt stress treatment of 21-day-old seedlings. The *arrows* indicate the spraying of Fluridone.

absolute ABA content and degree of tolerance. Therefore, they pointed out the importance of both the rate of ABA increase as well as the absolute ABA levels. The fact that IR29 produce less ABA under stress, could explain their sensitivity response to stress (Figure 3). Further experiments, using Fluridone, confirm the involvement of ABA synthesis in the mechanism of response to stress. Irrespective of cultivar used, spraying 50 μM Fluridone, drastically increased leaf Na concentration the tolerance and sensitive varieties when exposed to saline stress (Figure 2). However, the extent of this increment varied as a function of the tolerance level of the varieties used. Thus, in tolerance variety the increment was of 3 fold, while in sensitive variety it reached 8 fold in their respective to control treatments (Figure 2). Thus, when plants of both cultivars were sprayed with Fluridone, they showed higher sensitivity to a subsequent saline stress than stressed plants that had not been sprayed with the inhibitor. This protective role of ABA is confirmed in the present work (Popova, 1998). While stress induced dramatically increased endogenous ABA, spraying with 50 μM Fluridone during the salinity

treatment suppressed this increase (Figure 3). We observed that an ABA treatment during saline stress nearly to 7 times reduced the leaf Na content of sensitive cultivar to recover from stress in comparison to untreated plants subjected to the same stress conditions (Figure 2). In contrast, tolerance cultivar was not affected by the hormonal treatment. In agreement with this result, Bohra et al. (1995) have also shown that exogenous ABA treatments improved tolerance to salinity in sensitive but not in tolerant rice cultivars. It is interesting to note that non-stressed barley varieties with lower ABA contents were more sensitive to freezing and responded better to exogenous ABA treatments than other varieties with higher hormone levels (Bravo et al., 1998). In fact, growth rate and leaf Na concentration of plants were significantly affected by this inhibitor (Figures 1 and 2), which interfere with ABA action. Similar results have been obtained by Bianco-Trinchant and Le Page-Degivry (1998) in protoplasts from different plant origin. Our results demonstrate the importance of ABA synthesis in their tolerance to stress, tolerance variety with a higher capacity of ABA synthesis (Figure 3). In summary, our

study lead to the following conclusions:

1. The degree of stress tolerance of a given variety does not appear to be related to endogenous ABA levels under control conditions, but to the different rate of ABA synthesis under stress.
2. The fact that cells from IR651 produce more ABA when subjected to stress, could explain their better response to salinity condition.
3. The beneficial effect of exogenous ABA on the tolerance to saline stress is mainly restricted to IR29 with low tolerance (sensitive cultivar).
4. There was a different threshold irritability for ABA content, because opposite to the bigger leave ABA level in sensitive variety, more deleterious effect observed in this cultivar.
5. Regardless to present or absent of salinity treatment, IR651 shows a higher growth rate, and low accumulation of Na with a Fluridone application in contrast to IR29, suggesting that improved growth probably alleviating Na+ toxicity by diluting, this finding consistent with Nooden (1988).

## ACKNOWLEDGEMENT

The author gratefully acknowledge the funding from the Islamic Azad University Shoushtar branch through Grant.

## REFERENCES

- Asch F, Dorffling K, Dingkuhn M (1995). Response of rice varieties to soil salinity and air humidity: A possible involvement of root-borne ABA. *Plant Soil*, 177: 11-19.
- Bianco-Trinchant J, Le Page-Degivry MT (1998). ABA synthesis in protoplasts of different origin in response to osmotic stress. *Plant Growth Regul.*, 25: 135-141.
- Bohra JS, Dörffling H, Dörffling K (1995). Salinity tolerance of rice (*Oryza sativa* L.) with reference to endogenous and exogenous abscisic acid. *J. Agron. Crop Sci.*, 174: 79-86.
- Bornman CH, Jansson E (1980). *Nicotiana tabacum* callus studies. X. ABA increases resistance to cold damage. *Physiol. Plant*, 48: 491-493.
- Bravo LA, Zúñiga GE, Alberdi M, Corcuera LJ (1998). The role of ABA in freezing tolerance and cold acclimation in barley. *Physiol. Plant*, 103: 17-23.
- Cha-um S, Supaibulwatana K, Kirdmanee C (2007). Glycinebetaine accumulation, physiological characterizations and growth efficiency in salt-tolerant and salt-sensitive lines of indica rice (*Oryza sativa* L. ssp. indica) in response to salt stress. *J. Agron. Crop Sci.*, 193: 157-166.
- Chen S, Li J, Wang T, Wang S, Polle A, Hüttermann A (2002). Osmotic stress and ion-specific effects on xylem abscisic acid and the relevance to salinity tolerance in poplar. *J. Plant Growth Regul.*, 21: 224-233.
- Da Silva EC, Nogueira RJMC, De Araujo FP, De Melo NF, De Azevedo Neto AD (2008). Physiological responses to salt stress in young umbu plants. *Environ. Exp. Bot.*, 63: 147-157.
- Dionisio-Sese ML, Tobita S (1998). Antioxidant responses of rice seedlings to salinity stress. *Plant Sci.*, 135: 1-9.
- Duncan DR, Widholm JM (1987). Proline accumulation and its implication in cold tolerance of regenerable maize callus. *Plant Physiol.*, 83: 703-708.
- Faltusová-Kadlecová Z, Faltus M, Prásil I (2002). Abscisic acid content during cold hardening of barley and wheat cultivars with different freezing tolerance. *Rostlinná Výroba*, 48: 490-493.
- Gómez-Cadenas A, Pozo OJ, García-Agustín P, Sancho JV (2002). Direct analysis of abscisic acid in crude plant extracts by liquid chromatography/electrospray-tandem mass spectrometry. *Phytochem. Anal.*, 13: 228-243.
- Gregorio GB, Senadhira D, Mendoza RD (1997). Screening rice for salinity tolerance. International Rice Research Institute discussion paper series No. 22.
- Hasegawa PM, Bressan RA, Zhu J, Bohnert HJ (2000). Plant cellular and molecular responses to high salinity. *Anal. Rev. Plant Physiol. Plant Mol. Bio.*, 51: 463-499.
- Kadlecová Z, Faltus M, Prásil I (2000). Relationship between abscisic acid content, dry weight and freezing tolerance in barley cv. Lumet. *J. Plant Physiol.*, 157: 291-297.
- Lee TM, Lur HS, Chu C (1993). Role of abscisic acid in chilling tolerance of rice (*Oryza sativa* L.) seedlings. I. Endogenous abscisic acid levels. *Plant Cell Environ.*, 16: 481-490.
- Moons A, Bauw G, Prinsen E, Montagu MV, Van Der Straeten D (1995). Molecular and physiological responses to abscisic acid and salt in roots of salt-sensitive and salt-tolerant indica rice varieties. *Plant Physiol.*, 107: 177-186.
- Moradi F, Ismail AM, Gregorio GB, Egdane JA (2003). Salinity tolerance of rice during reproductive development and association with tolerance at the seedling stage. *Indian J. Plant Physiol.*, 8: 276-287.
- Morsy MR, Jouve L, Hausman JF, Hoffmann L, Mcd Stewart J (2007). Alteration of oxidative and carbohydrate metabolism under abiotic stress in two rice (*Oryza sativa* L.) genotypes contrasting in chilling tolerance. *J. Plant Physiol.*, 164: 157-167.
- Munns R (2005). Gene and salt tolerance: bringing them together. *New Phytol.*, 167: 645-663.
- Nayyar H, Kaushal SK (2002). Alleviation of negative effects of water stress in two contrasting wheat genotypes by calcium and abscisic acid. *Biol. Plant*, 45: 65-70.
- Nayyar H, Walia DP (2003). Water stress induced proline accumulation in contrasting wheat genotypes as affected by calcium and abscisic acid. *Biol. Plant*, 46: 275-279.
- Nooden LD (1988). Abscisic acid, auxin and other regulators of senescence. In: L. D. Nooden ed. *San Diego. Senescence and aging in plants: Academic Press Inc*, pp. 329-368.
- Popova L (1998). Fluridone- and light-affected chloroplast ultrastructure and ABA accumulation in drought-stressed barley. *Plant Physiol. Biochem.*, 36: 313-319.
- Rhoades JD, Kandiah A, Mashali AM (1992). The use of saline water for crop production (FAO irrigation and drainage paper 48). FAO of the United Nations, Rome.
- Shannon MC, Rhoades JD, Draper JH, Scardaci SC, Spyres MD (1998). Assessment of salt tolerance in rice cultivars in response to salinity problems in California. *Crop Sci.*, 38: 394-398.
- Yasar F, Uzal O, Tufenkci S, Yildiz K (2006). Ion accumulation in different organs of green bean genotypes grown under salt stress. *Plant Soil Environ.*, 52: 476-480.
- Yoshida S, Forno DA, Cock JK, Gomez KA (1976). Laboratory manual for physiological studies of rice. International Rice Research Institute, Manila, Philippines. p. 124.
- Zheng Y, Wang Z, Sun X, Jia A, Jiang G, Li Z (2008). Higher salinity tolerance cultivars of winter wheat relieved senescence at reproductive stage. *Environ. Exp. Bot.*, 62: 129-138.