

Full Length Research Paper

Physico-chemical properties of Chakiya variety of Amla (*Emblica officinalis*) and effect of different dehydration methods on quality of powder

Poonam Mishra^{1*}, Vijeyta Srivastava¹, Deepmala Verma¹, O. P. Chauhan² and G. K. Rai¹

¹Centre of Food Technology, University of Allahabad, Allahabad, India.

²Defence Food Research Laboratory, Siddarthanagar, Mysore, India.

Accepted 09 October, 2019

In present study a comparative study of wild variety and cultivated variety of *Amla* were carried out; simultaneously effect of different drying techniques on physicochemical properties of *Amla* powder were also studied. The fruits of wild variety were found smaller than the cultivated variety. In case of wild variety, the total phenolic contents were found to be 32.32 g/100 g of gallic acid equivalent (dwb), whereas Chakiya variety had 24.50 g/100 g of gallic acid equivalent (dwb). The fruits of Chakiya variety were used to prepare the powder by different techniques like freeze drying, sun drying, spray drying, hot air drying and vacuum drying. Powder yield varied with type of drying method as sun drying (10.11%), tunnel drying (8.78%) and vacuum drying (12.48%), spray drying (4.90%) and freeze drying (2.23%). Significant differences ($p < 0.05$) in chemical composition of *Amla* powder were observed when prepared by different techniques. The freeze dried powder had the highest ascorbic acid content followed by spray dried powder. The lowest concentration of ascorbic acid was found in sun dried powder. Freeze dried samples showed maximum mineral contents in terms of calcium (79.6 mg/100 g), phosphorus (12.38 mg/100 g) and iron (88.03 mg/100 g).

Key words: Amla, dehydration methods, quality, polyphenols.

INTRODUCTION

Amla (*Emblica officinalis*, family Euphorbiaceae) is a native of India, Ceylon, Malaya and China. The fruit is used as a major constituent in several Ayurvedic preparations such as *Chyavanprash* and *Rasayana* which promotes health and longevity (Rajkumar et al., 2001). *Amla* is one of the richest sources of Vitamin C known. In addition to this, potent antioxidant, several active tannoid principles (Emblucannin A, Emblucannin B, Punigluconin and Pedunculagin) have been identified which appear to account for its health benefits (Rastogi 1993; Rao et al., 1985). *Amla* has been reported to possess expectorant, purgative, spasmolytic, antibacterial, hypoglycemic (Jamwal et al., 1959; Jayshri and Jolly, 1993), hepatoprotective and hypolipidemic (Thakur and

Mandal, 2004) activity. The aqueous extract has been reported to have anti-pyretic laxative and tonic properties and also showed antibacterial activity (Vinaygamoorthy, 1982). The ascorbic acid content of fresh *Amla* fruit can range up to 950/100 gm which is said to be highest among all fruits next only to Barbados cherry (Shankar, 1969).

Several value added products have been reported from *Amla*. *Amla* was used to prepare ready-to-serve beverage (Deka et al., 2001), candy, powder (Tripathi et al., 1988), sauce (Chauhan et al., 2005) etc. The quality of final product depends on the nature of the raw material used for its preparation. There are several varieties of *Amla* grown throughout different parts of India. However, these varieties are yet to be researched in terms of physico-chemical properties for value addition (Kalra, 1988). In addition, storage conditions also play important role in postharvest quality of *Amla* fruits. Chakiya is an important variety of *Amla* which is grown

*Corresponding author. E mail: mpoonam1@rediffmail.com.
Tel: 09415081554.

Table 1. Physical properties of fruits of wild and Chakiya variety of Amla (n=15).

Particular	Chakiya variety	Wild variety
Average length, cm	3.0 ^a	2.72 ^b
Average diameter, cm	3.8 ^a	3.0 ^b
Average weight, g	28.88 ^a	17.66 ^b
No. of segments per fruit	6 ^a	6 ^a
Juice yield, %	41.9 ^b	48.3 ^a

Values in same row with different superscripts differ significantly ($p < 0.05$).

in the Eastern regions of India (Kumar and Khurdiya, 2002). However, little work has been carried out with respect to physicochemical characterization of this variety and powder preparation from the fruits of this particular variety. Therefore, present work was undertaken to study the physicochemical attributes of the fruits of Chakiya varieties as well as a wild variety and preparation of powder. Effect of different drying methods such as sun drying, tunnel drying, spray drying, freeze drying and vacuum drying were also studied with respect to the quality of Amla powder.

MATERIALS AND METHODS

Raw material and preparation of Amla powder

Amla of Chakiya variety (cultivated) was procured from local market while wild variety of was obtained from forest of Panna (Madhya Pradesh, India).

The Amla fruits were cleaned and cut into pieces and immediately pressed to obtain juice using a small laboratory manual press. For the preparation of spray dried powder, juice was evaporated up to 50% using rotary evaporator. Maltodextrin (5%, w/v of initial juice) was added to the concentrated juice and stirred for 5 min using mechanical stirrer followed by spray drying (Model LU 228 advance, Labultima, Mumbai, India) at 200°C inlet and 150°C outlet temperature and 40% aspiration speed to dry the sample. For freeze drying 200 ml of juice was subjected to freezing at -35°C for 3 h followed by freeze dehydration in a freeze dryer (Model Alpha 1-4, Martin Christ, Germany) at 60°C for 16 h. Sun dried powder was prepared from grated Amla dried under sun followed by grinding in a mixer grinder and filtration using muslin cloth. The powder was also prepared by drying the grated Amla in a vacuum oven (KC Instruments, Lucknow, India) at 50°C and in a tunnel drier (B. Sen Berry and Co. India) at 70°C followed by grinding and filtration as described earlier.

ANALYTICAL METHODS

Physical properties

Average fruit weight, length, diameter and number of segments per fruit were measured taking 15 samples of both the varieties. The juice yield was calculated by pressing the fruit segments after removal of seeds and the data were converted on percentage basis.

Moisture, crude fat and mineral content

The moisture and crude fat content of the samples were determined as per AOAC (1997) procedures; whereas minerals (Ca, P and Fe) were estimated as per the AOAC (1990) procedures.

Total phenolic content

Total polyphenols were estimated as per procedure described by Jayprakash et al. (2001) where 250 mg sample was taken in 10 ml of methanol and water (70:30 v/v) solution in a graduated test tube and heated on water bath at 70°C for 10 min. The sample was brought to room temperature, centrifuged at 3500 rpm for 10 min.

The supernatant (0.2 ml) was made up to 10 ml with distilled water. This solution was diluted 10 fold. Sample solution (5 ml) was mixed with saturated sodium carbonate (0.5 ml) and Folin-Ciocalteau reagent (0.2 ml) and made up to 10 ml with distilled water. The absorbance was read at 765 nm after 60 min by UV visible double beam spec-trophotometer (Model Evolution 600, Thermo Electron, US).

Ascorbic acid

Sample solution equivalent to 0.2 mg ascorbic acid/ml was prepared in water containing 3% (w/v) metaphosphoric acid. It was titrated against standard 2, 6 dichlorophenol indophenol (2,6 DCIP) solution of 0.5 mg/ml concentration until the pink color developed completely. The operation was repeated with a blank (Indian Pharmacopodia).

Tristimulus colour

Tristimulus color in terms of Hunter L, a, b values was measured using X-Rite spectrophotometer (USA) using D-65 illuminant and 10° observer. 'L' value represents lightness, 'a' value shows redness-greenness and 'b' value indicates blueness-yellowness of the samples.

Statistical analysis

The data obtained were analyzed statistically for analysis of variance (ANOVA) using completely randomized design with least significant difference (LSD) at $p \leq 0.05$ using Co.Stat 6.303, CoHort software (USA).

RESULTS AND DISCUSSION

Physical properties of wild and cultivated Chakiya variety of Amla fruits is given in Table 1. The fruits of wild variety were found smaller than the cultivated variety. There was significant difference ($p < 0.05$) in weight of the fruits of both the varieties. An average weight of the fruits of the wild variety of Amla was 17.66 g, whereas cultivated variety showed an average weight of 28.88 g. The fruits of both the varieties have six segments. Juice yield was found to be significantly ($p < 0.05$) higher in wild variety (48.3%) as compared to Chakiya variety (41.9%). In case of wild variety, the total phenolic contents were found to be 32.32 g/100 g of gallic acid equivalent (dwb), whereas, Chakiya variety had 24.50 g/100 g of gallic acid

Table 2. Chemical properties of fruits of wild and Chakiya variety of *Amla* (n=3).

Particular	Chakiya variety	Wild variety
Ascorbic acid (g/100 g)	463.0 ± 6.83 ^b	560.0 ± 9.42 ^a
TSS (^o Brix)	9.3 ± 0.23 ^b	13.1 ± 0.105 ^a
Total phenols in residue (g/100g, dwb)	4.0 ± 0.23 ^b	12.1 ± 0.35 ^a
Total phenols in juice (mg/100 gm, fwb)	174.0 ± 4.7 ^a	176.6 ± 6.18 ^a
Total Phenols in fresh <i>Amla</i> (g/100g, dwb)	24.5 ± 1.11 ^b	32.3 ± 1.35 ^a

Values in same row with different superscripts differ significantly (p < 0.05).

Table 3. Mineral content and some chemical properties of *Amla* powder (n=3).

Drying method	Ca	P	Fe (mg/100gm)	Ascorbic acid	Moisture (%)	Fat
Freeze dried	79.6 ± 0.72 ^a	12.38 ± 0.32 ^a	88.03 ± 0.75 ^a	5432.75 ± 16.11 ^a	6.78 ± 0.03 ^a	0.24 ± 0.005 ^d
Sun dried	70.3 ± 0.72 ^b	7.90 ± 0.24 ^c	54.20 ± 0.72 ^b	3006.00 ± 14.40 ^e	5.76 ± 0.19 ^c	0.39 ± 0.009 ^c
Vacuum dried	64.0 ± 0.27 ^c	9.16 ± 0.83 ^b	53.87 ± 0.66 ^b	3339.88 ± 15.14 ^d	6.53 ± 0.12 ^b	0.42 ± 0.094 ^b
Spray dried	55.0 ± 0.94 ^d	5.10 ± 0.25 ^d	49.26 ± 0.91 ^c	4058.44 ± 11.32 ^b	5.05 ± 0.91 ^e	0.23 ± 0.021 ^e
Tunnel dried	50.1 ± 1.23 ^e	7.85 ± 0.43 ^c	53.67 ± 0.97 ^b	3401.00 ± 13.31 ^c	5.59 ± 0.16 ^d	0.59 ± 0.089 ^a

Values in same column with different superscripts differ significantly (p < 0.05).

equivalent (dwb) (Table 2). The total phenolic contents were also found high in the juice and residue (left after extraction of juice) of wild variety of *Amla* than the Chakiya variety. The juices had total phenolic concentrations in the range of 174.0 and 176.6 mg/100 g of juice (fwb) in Chakiya and wild variety, respectively.

The phenolic contents in juices are low compared to fruits due to loss of phenolics during squeezing process. Another reason may be that polyphenols are involved in specific physicochemical interactions with the solid part of the fruits, especially the cell wall material (Ranaurd et al., 2001). This was reconfirmed by the polyphenolic estimation of residue of juice. Ascorbic acid content was also found to be high in wild variety of *Amla* compared to the cultivated variety (Table 2). Kaur and Kapoor (2002) reported more than 70% antioxidant activities as percentage inhibition of oxidation in *Amla* fruits which was correlated positively with total phenols. The total soluble solids in juice prepared from cultivated variety (9.3^oBrix) of *Amla* were also lower than that of wild variety (13.1^oBrix).

Effect of drying methods on the quality of powder

The fruits of Chakiya variety were used to prepare the powder. Powder yield varied with type of drying method as sun drying (10.11%), tunnel drying (8.78%) and vacuum drying (12.48%), spray drying (4.90%) and freeze drying (2.23%). Significant differences (p<0.05) in chemical composition of *Amla* powder were observed when prepared by different techniques. The total ascorbic acid content ranged from 3.00 to 5.43 g/100 g of powder. The freeze dried powder had the highest ascorbic acid

content followed by spray dried powder. The lowest concentration of ascorbic acid was found in sun dried powder (Table 3). When *Amla* fruits were cut and dried the oxidation process already took place and this may be the reason of low ascorbic acid content in sun dried powder. The rate of oxidation was high during sun drying, hence the sun dried sample showed lowest amount of ascorbic acid than other samples. Moisture content varied from 5.05 to 6.78% whereas fat content varied from 0.23 to 0.59% in the powder prepared by using different dehydration methods. Freeze dried samples showed maximum mineral contents in terms of calcium (79.6 mg/100 g), phosphorus (12.38 mg/100 g) and iron (88.03 mg/100 g). Tristimulus colour values of the powder on Hunter scale in terms of L, a, b values are given in Table 4. Significant differences (p < 0.05) were observed in L, a, b values of the samples. Freeze dried sample was found lighter in color as compared to other powder samples, whereas the sun dried sample was found to be darkest in color which can be observed by higher L value in freeze dried (61.64 and lower L value in sun dried (45.78) samples. Higher redness was observed in sun dried sample followed by tunnel, vacuum, freeze and spray dried samples. Where-as, b value was maximum in tunnel dried sample and minimum in spray dried sample. On the basis of L, a, b co-lour values, the freeze dried sample was considered better in terms of colour values followed by spray dried one.

Conclusion

Amla (var. Chakiya) can be used for development of value added products. This variety had a lower content of

Table 4. Tristimulus colour profile of *Amla* powder (n=3).

Drying method	L	a	b
Freeze dried	61.64 ^a	-0.16 ^d	15.02 ^b
Sun dried	45.78 ^e	2.49 ^a	14.37 ^c
Vacuum dried	46.18 ^d	1.58 ^c	12.77 ^d
Spray dried	53.84 ^c	-0.30 ^e	7.88 ^e
Tunnel dried	59.17 ^b	1.67 ^b	15.15 ^a

Values in same column with different superscripts differ significantly ($p < 0.05$).

phenolic content, total soluble solids and ascorbic acid as compared to wild variety. However, in terms of average weight, length and diameter the fruit of this variety was found better than wild variety. The powder developed from this variety showed better retention of nutrients and colour in the case of freeze dried samples followed by spray dried ones.

REFERENCES

- AOAC (1990). Official methods of analysis of Association of Analytic Chemists, Washington DC.
- AOAC (1997). Official methods of analysis of Association of Analytic Chemists, Washington DC.
- Chauhan OP, Srivastava S, Pandey P, Rai GK (2005). A Study on the Development of *Aonla* Blended Sauce. *Beverage Food World*. 32 (5): 31-32.
- Deka BC, Sethi V, Prasad R, Batra PK (2001). Application of mixtures methodology for beverages from mixed fruit juice/pulp. *J. Food Sci. Technol.*, 38(6): 615-618.
- Jamwal KS, Sharma IP, Chopra L (1959). Pharmacological investigations on the fruits of *Emblica Officinalis*. *J. Sci. Ind. Res.* 18 c: 180-181.
- Jayprakash GK, Singh RP, Sakariah KK (2001). Antioxidant activity of grape seed extracts on peroxidation models *in vitro*. *Food Chem.*, 73:285-290.
- Jayshri S, Jolly CI (1993). Phytochemical antibacterial and pharmacological investigations on *Monordica chirantia* and *Emblica officinalis*. *Ind. J. Pharm. Sci.* 1: 6-13.
- Kalra CL (1988). The chemistry and technology of *Amla* (*Phyllanthus emblica*) - a resume. *Indian Food Packer*. 42(4): 67-82.
- Kaur C, Kapoor HC (2002). Anti-oxidant activity and total phenolic content of some Asian vegetables. *International J. Food Sci. Technol.*, 37(2): 153-161.
- Kumar JS, Khurdiya DS (2002). Studies on juice extraction of aonla (*Emblica officinalis* Gaertn.) cv. 'ÁChakaiya'. *J. Food Sci. Technol.*, 39(5): 515-516.
- Rajkumar NV, Theres M, Kuttan R (2001). *Emblica officinalis* fruits afford protection against experimental gastric ulcers in rats. *Pharmaceut. Biol.*, 39 (5):375-380.
- Ranaurd CMGC, Baron A, Guyot S, Drilleau JF (2001). Interactions between apple cell walls and native apple polyphenols: quantification and some consequences. *Int. J. Biol. Macromol.*, 29: 115-125.
- Rao TS, Kumari KK, Netaji B, Subhokta PK (1985). *Ayurveda Siddha*. *J Res.* 6:213-224.
- Rastogi RP (1993). *Compendium of Indian Medicinal plants*, CDRI, Lucknow and ID, New Delhi 1: 530.
- Shankar G (1969). *Aonla* for your daily requirement of Vitamin C. *Indian Hort.* 13: 9-15.
- Thakur CP, Mandal K (1984). Effect of *Emblica Officinalis* on Cholesterol Induced atherosclerosis in rabbits. *Ind. J. Med. Res.*, 79: 142-146.
- Thakur CP, Mandal K (1984). Effect of *Emblica officinalis* on cholesterol induced atherosclerosis in rabbits. *Ind. J. Med. Res.* 79:142-146.
- Tripathi VK, Singh MB, Singh S (1988). Studies on comparative compositional changes in different preserved products of *Amla* (*Emblica officinalis* Gaertn) var. Banarasi. *Indian Food Packer.*, 42(4): 60-66.
- Vinayagamoothy T (1982). Antibacterial activity of some medicinal plants of Srilanka Ceylon. *J. Sci. Biol.*, 11: 50-55.