

Full Length Research Paper

Effect of different pesticide application methods on spray deposits, residues and biological efficacy on strawberries

Nigar Yarpuz-Bozdogan^{1*}, Ekrem Atakan², Ali Musa Bozdogan³, Huseyin Yilmaz⁴, Nebile Daglioglu⁵, Tunahan Erdem³ and Ebru Kafkas⁶

¹Vocational School of Technical Sciences, University of Çukurova, Adana/Turkey.

²Department of Crop Protection, Faculty of Agriculture, University of Çukurova, Adana/Turkey.

³Department of Agricultural Machinery, Faculty of Agriculture, University of Çukurova, Adana/Turkey.

⁴NetKim Laboratory Equipment Marketing Co., Adana/Turkey.

⁵Department of Forensic Medicine, Faculty of Medicine, University of Çukurova, Adana/Turkey.

⁶Biotechnology Research and Experimental Center, University of Çukurova, Adana/Turkey.

Accepted 31 August, 2019

In agricultural food production, positive results can be obtained by using appropriate pesticides through the application of appropriate methods. In this study, we aimed to determine the effects of different spraying methods on spray deposits and drift, pesticide residues and biological efficacy on strawberries. For this purpose, strawberries were sprayed with dicofol by broadcast and band spraying. The effectiveness of different application methods was studied on 09 May, 2007 and 16 April, 2008. Broadcast sprayings were applied via hollow cone nozzles (HC) and air-assisted spinning cage nozzles (ASC). Band spraying was applied via flat fan nozzles (FF). Pesticide deposits on leaf surfaces, ground, drift and pesticide residues on strawberries were analyzed with Gas Chromatography/Electron capture detector (GC/ECD). Strawberries were analyzed based on dicofol residues up to 6 days after spraying. The results obtained showed that the highest pesticide deposits on leaf surfaces and also biological efficiency were obtained with FF in 2007 and 2008. The lowest pesticide deposits on ground were obtained by HC and the highest deposit of drift was obtained by ASC in 2007 and 2008. According to Commission Directive 2000/42/EC, the MRL's of dicofol on strawberry is 0.02 mg/kg. In this study, residues on strawberries for all types of nozzles were below the suggested maximum residue level (MRL) of dicofol by Commission Directive 2000/42/EC in 2007 and 2008.

Key words: Broadcast spraying, band spraying, hollow cone nozzle, flat fan nozzle, air-assisted spinning cage nozzle, biological efficacy, red spider mites, gas chromatography, electron capture detector (ECD).

INTRODUCTION

In modern quality agricultural food production, efficient results can be obtained by the use of suitable pesticides and their application methods. Broadcast spraying, which is used by farmers due to easy application, covers the pesticides in all field area. On the other hand, band spraying is defined as covering the pesticides in all band width where pesticides are applied on crops for insecticides and fungicides, and on soil between crops

row spacing for herbicides. Some farmers use band spraying method for strawberry pest management. Band spraying is regarded to be more economical compared to broadcast spraying, since the row application kit ensures that the pesticide goes where it is directed, resulting in a more effective application (Landers, 2007). Moreover, band spraying has advantages such as reduction in pesticides and water use per hectare (Debear et al., 2005).

Different nozzles such as hollow cone, flat fan, and spinning disc/cage (with/without air-assistance) are some of the main types used in broadcast spraying method. Hollow cone (HC) and flat fan (FF) nozzles can be used

*Corresponding author. E-mail: nyarpuzbozdogan@cu.edu.tr.
Tel/Fax: (90) 322 - 338 65 18.

in high, normal and low volume applications. Spinning disc/cage nozzles with/without air assistance can be used in low and ultra-low volume applications. In spraying, using spinning disc/cage nozzles with air assistance improves leaf coverage through the plant canopy, provides better coverage to the underside of leaves, enables more penetration into crop canopy, and covers more acres per load (Taylor and Andersen, 1989; Panneton et al., 1996; Holownicki et al., 2000; Panneton et al., 2000a, Panneton et al., 2000b; Bayat and Yarpuz-Bozdogan, 2005). In Turkey, hollow cone nozzles are widely used in broadcast spraying.

Pesticide residue on strawberries is very important and has negative affects on human health (Goodwin et al., 1984; Kilmer et al., 2001; Safi et al., 2002). Some researchers use gas chromatography-electron capture detector (GC-ECD) for pesticide residues on vegetables and fruits (Stensvand and Christiansen, 2000; Safi et al., 2002; Yenisoay-Karakas, 2006; Guardia-Rubio et al., 2007).

There are many harmful insects and mite species causing injuries to strawberry roots, leaves and flowers. Red spider mites (RSM) (Acarina: Tetranychidae) are often considered as major pest acari of strawberry, destroying the mesophyll cells of leaves and high population densities that can result in the plant being dwarfed or killed (Yigit and Erkiliç, 1992). Damage is expressed as stippling and bronzing of the leaves and leave's veins. Plants that sustain greater numbers of RSM may become severely weakened and appear stunted, dry, and red in coloration. *Tetranychus urticae* Koch and *Tetranychus cinnabarinus* (Boisd.) were more common RSM species in Turkey and they were recognized as pest (Oncag and Cengiz, 1978). There are a few studies regarding use of various spraying techniques for controlling of RSM on strawberry. Labanowska et al. (1996) compared two spraying techniques for the control of *T. urticae* and grey mold *Botrytis cinerea* on strawberry by testing three different spray liquid volumes 260, 400 and 550 per hectare and two doses of pesticides. They found that all spray volumes sufficiently controlled the *T. urticae*, but full recommended doses of the pesticides must be used.

In this study, different pesticide application methods were used against strawberry pests. The aim of this study was to determine the effects of different pesticide application methods on spray deposits, spray drift, pesticide residues and biological efficacy on strawberries. Air assisted spinning cage nozzles (ASC) were utilized as an alternative to the hollow cone (HC) nozzles in the broadcast method and the flat fan (FF) nozzles were used in the band spraying method.

MATERIALS AND METHODS

Location

In this study, strawberries (*Fragaria X ananassa*, Duch, Rosaceae) were grown in research experimental fields of the Yaltir Corporation

in Adana which is in the Mediterranean region of Turkey. The spraying experiment was designed as a complete randomized block with four replicates; also 10 plants of cultivar camarosa species were randomly selected in each replicate. Each plot covers 10 m in length and 5 m in width. The effectiveness of different application methods was studied on 09 May, 2007 and 16 April, 2008.

Pesticide

In this study, 0.292 kg a.i. ha⁻¹ dicofol/Kelthane™ was used for strawberries. Maximum residue level (MRL) value of dicofol is 0.02 mg/kg (EC, 2000).

Spraying methods

Two different types of spraying methods were used: (1) broadcast spraying method with hollow cone nozzles (HC) (200 l/ha) and air-assisted spinning cage nozzles (ASC) (20 l/ha), and (2) band spraying method with flat fan nozzles (FF) (200 l/ha) (Table 1). Only one field sprayer was used. The following options were applied onto the field sprayer:

- 1) Hollow cone (HC) nozzles (Tara™, D4-45, 50 cm above the crop) as broadcast spraying.
- 2) Air-assisted spinning cage (ASC) nozzle (Micron™, fan/atomizer speed 4000 rpm) as broadcast spraying. ASC nozzles were situated towards ground plane with an angle of 45°.
- 3) Row application kit with flat fan (FF) nozzles (Arag™, 11001, one directed towards the centre and two towards the sides of the crop) as band spraying.

The forward speed of the tractor was 6.5 km/h and the power-take-off (PTO) rotational speed was 540 rpm. The spray boom of the field sprayer (HC) covered 4 rows of the strawberry plantation in one swath. On the other hand, the spray boom (single-sided) of the field sprayer with FF and ASC nozzles covered two rows of the strawberry plantation in one swath. Each trial was repeated four times.

Meteorological conditions

During the trials, temperature, wind velocity and relative humidity were recorded. They were measured by digital Hotwire Anemometer CE (Lutron™ AM-4204 HA) and digital Humidity/type k Thermometer CE (Lutron™ HT-3006 HA). Details of meteorological conditions during experiments were shown in Table 2.

Artificial sampling methods of spray deposit on plant and the ground and drift measurements

Spray deposits were evaluated by fixing filter papers (Schleicher and Schuell, Germany), 30 × 30 mm dimension, on the upper and lower leaf surfaces at four leaves (left, right, front and back) on strawberry plants. For each replication, 10 plants were used. Spray deposits on the ground were collected on three wooden wedges to which filter papers (30 × 40 mm) were attached and placed between rows. The drift measurement was laid perpendicular to the line of travel of the sprayer and parallel to the wind direction. Using the method described by Rowinski et al. (1994), spray drift were placed horizontally along the line of measurement at distances of 1, 2, 3, 4, 5, 7.5, 10, 15, 30 and 50 m respectively, from the end of the boom. The average deposits on plant and the ground, and drift deposits were evaluated separately. After spraying, the filter paper samples were collected from field and placed in bottles (height: 9

Table 1. Details of pesticide application methods.

| Pesticide application methods | Pressure (bar) | Flow rate (L min ⁻¹) | Application rate (L ha ⁻¹) | VMD (µm) | Air-Assisted velocity (ms ⁻¹) |
|-------------------------------|----------------|----------------------------------|--|----------|---|
| Broadcast spraying | | | | | |
| HC | 9.5 | 1.07 | 200 | 250 | - |
| ASC | 1.2 | 0.13 | 20 | 100-120 | 25 |
| Band spraying | | | | | |
| FF | 1.2 | 0.22 | 200 | 200 | - |

Table 2. Details of meteorological conditions during experiments.

| Year | Pesticide application methods | Spraying time (h) | Temperature (°C) | Relative humidity (%) | Wind speed (m/s) |
|------|-------------------------------|-------------------|------------------|-----------------------|------------------|
| 2007 | Broadcast spraying | | | | |
| | HC | 09:55 – 10:03 | 24.3 - 25.5 | 59.1 – 61.9 | 2.7 – 3.6 |
| | ASC | 15:14 – 15:26 | 30.1 – 31.1 | 45.8 – 47.2 | 0.8 – 1.4 |
| 2007 | Band spraying | | | | |
| | FF | 11:24 – 11:43 | 29.3 – 32.8 | 42.6 – 50.0 | 1.5 – 2.7 |
| 2008 | Broadcast spraying | | | | |
| | HC | 10:40 – 10:48 | 24.0 – 25.6 | 58.1 – 63.0 | 0.3 – 1.3 |
| | ASC | 08:39 – 08:57 | 21.1 – 22.6 | 64.4 – 68.6 | 0.2 – 1.5 |
| | Band spraying | | | | |
| | FF | 09:56 – 10:14 | 21.1 – 22.8 | 65.1 – 70.1 | 0.4 – 1.1 |

cm; diameter; 4.5 cm) and taken to laboratory.

Spray deposits on leaf surfaces of strawberry plants

For spray deposit on leaf surfaces of strawberry plants, four leaves (left, right, front and back) were randomly picked for each replication. For each replication, 10 plants were used. After spraying, leaves were collected from field and placed in bottles (height: 9 cm; diameter: 4.5 cm) and taken to laboratory.

Pesticide residues on strawberries

Strawberries were collected after the first hour (accepting as zero) then first, second, third, fourth, fifth and sixth days after spraying for the residue analysis. They were randomly sampled 1 kg in each plot. The samples were kept in plastic bags.

Extraction

Liquid- liquid extraction method was used. Strawberry samples and leaves were weighed and transferred to a steel blender (Waring Commercial Blender™). 100 ml acetone was added then blended. The sample was filtered with suction. The collected extract was poured in 250-ml suction flask. Later 40 ml sample extract was placed in 1 L separator funnel for liquid/liquid partitioning; 50 ml petroleum ether and 50 ml methylene chloride were added into the

separator. The lower layer was transferred to a glass flask, while the upper layer was filtered through anhydrous sodium sulfate to eliminate residual water. The gained mixture was poured into evaporation tube (200 ml) of TurboVap System (Caliper Life Sciences™). The solution in the evaporation tubes was evaporated under nitrogen by TurboVap System. The resulting solution was ready for injection into the gas chromatograph.

Analytical equipment

An Agilent (6890 – 7683B auto-sampler) gas chromatograph equipped with an electro-capture-detector (ECD) and a HP-5MS capillary column (Agilent Technologies), stationary phase of 5% Phenyl Methyl Siloxane (30 m × 0.25 mm i.d., 0.25 µm film thickness) and nitrogen as the carrier gas (60 ml/min) were used. A split/splitless injector was used in the pulse splitless mode. The chromatographic conditions were: detector temperature was 290°C, capillary column temperature 325°C. The analytical procedure was applied according to GC methods (Pesticide Analytical Manual, 1999). The maximum residue levels (MRL) were adopted from Commission Directive 2000/42/EC.

Sampling of red spider mites

Ten plants were randomly selected in plots representing each spraying technique and one middle leaflet near to bottom section was picked from each selected strawberry plant. These leaflets

were wrapped in packing paper (250 g), placed in insulated cool containers and transferred to laboratory. Specimens of red spider mites on the leaflet samples were counted under a stereomicroscope with $\times 45$ magnification on the same day (Nelson and Show, 1975). Larvae, nymphs and adult stages were pooled into one category as mobile mites (MM). Samples were taken in the periods March 27 - May 22 (9 weeks) in 2007 and March 17 - May 12 in 2008 (9 weeks).

Statistical analysis

All data were analyzed with randomized block design methods using an SPSS package computer programme (SPSS, 1993). Significant differences between mean values were based on Duncan's multiple range tests (Duncan, 1955). Residues on strawberry and deposits on plant, ground and drift were used to compare the results of the different application methods. Comparisons of the mean numbers of red spider mite were quantified by the one-way analysis of variance (ANOVA) at $P < 0.05$. Their mean values were grouped by the Tukey's reliability test at $P < 0.05$. Henderson-Tilton formula (Henderson and Tilton, 1955) was used to calculate the corrected mortality (percentage).

RESULTS AND DISCUSSION

Artificial sampling methods of spray deposit on plant and the ground and drift measurements

The highest total average deposits were obtained from FF nozzles in 2007. Contrary to this, the results in 2008 showed that the highest total average deposits obtained from HC nozzles (Table 3). For trials conducted in 2007, the wind speed in HC nozzles was higher than other nozzles. For this reason, the total average deposits from HC nozzles in 2007 were lower than HC nozzles in 2008. Although pesticide deposits on upper and lower leaf surfaces of plants in ASC nozzles were lower than HC nozzles, no statistical difference was observed in 2007. However, in 2008, pesticide deposits on upper leaf surfaces in FF nozzles were lower than ASC and HC nozzles. In statistical analysis, pesticide deposits in FF, HC and, ASC nozzles were different from each other. Average pesticide deposits on upper leaf surfaces were found to be higher than average pesticide deposits on lower leaf surfaces of strawberries in HC both in 2007 and 2008.

Gan-Mor et al. (1996) reported that the air-streams enhanced spray deposition on the underside of leaves. In this study, air assist (ASC) improved the average deposit of lower leaf surfaces in 2008. Sumner and Herzog (2000) evaluated that the air-assisted sprayers offered better coverage than the other sprayers on the undersides of leaves and good coverage on the topsides. Manor et al. (1989) reported that the lower leaves droplet coverage and defoliation results were better with air-sleeve sprayer than with hydraulic sprayer. Similar results were found in this study. However, average pesticide deposits on lower leaf surfaces were similar to average pesticide deposits of upper leaf surfaces in ASC in 2008.

Also, average pesticide deposits on lower leaf surfaces were lower than upper leaf surfaces in ASC in 2007.

The highest pesticide deposits on ground were obtained with FF in 2007 and 2008 (Table 4). The highest spray drift was obtained at the first point (1 m) and the lowest spray drift was obtained at the last point (5 m) in all pesticide application methods in 2007 and 2008 (Figure 1). Thus, spray drift reduced when the distance was increased.

The highest spray drift was obtained with ASC in all pesticide application methods both in 2007 and 2008. ASC obtained smaller droplets than HC and FF. Small droplets can drift long distances because of their light weight (Wolf, 1997; Ozkan and Derksen, 1998; Matthews, 2000; PISC, 2002).

Spray deposits on leaf surfaces of strawberry plants

The highest pesticide deposits were obtained with FF nozzles in 2007 and 2008 (Table 5). Yet, in 2007 no statistical difference was observed in FF and HC. In statistical analysis, pesticide deposits in FF and HC were similar in 2008 (Table 5).

Pesticide residues on strawberries

Pesticide residues on strawberries of dicofol applied at different pesticide spraying methods are shown in Table 6. Residues were detected on the strawberry samples in all spraying methods. The highest residues in 2007 and 2008 were recorded for HC ranging from 0.0347 - 0.0164 mg/kg and 0.1479 - 0.0189 mg/kg, respectively. The lowest residues were recorded for ASC, ranging from 0.0218 - 0.0086 mg/kg in 2007 and finally for FF, ranging from 0.0797 - 0.0069 mg/kg in 2008. The 10 fold increase in spray volume between the LV (20 l ha^{-1}) and MV (200 l ha^{-1}) applications gave a greater increase in residue compared to the 2.5 fold reduction in volume median diameter (VMD). Cross et al. (2000) obtained similar results in efficacy of strawberry spraying.

Average pesticide residues on strawberries of dicofol applied through different pesticide application methods are shown in Table 7. When the application methods are compared, the highest pesticide residues were obtained in HC in 2007 and 2008. In statistical analysis, pesticide residue in HC was different from others. Although pesticide residue in ASC was lower than FF, no statistical significant difference was observed in 2007. However, in 2008, pesticide residue in FF was lower than ASC and in statistical analysis; pesticide residue in FF was different from ASC. Nordmark (1994) reported that a band sprayer resulted in better uniformity and less contamination of the environment than air-assisted boom sprayer. These pesticide application methods were more efficient than broadcast boom sprayer with hollow cone nozzles in vegetables and strawberries.

Table 3. Average pesticide deposit ($\mu\text{g}/\text{cm}^2$) on upper and lower leaf surfaces of plants.

| Pesticide application method | Average pesticide deposit ($\mu\text{g}/\text{cm}^2$)* | | | | | |
|------------------------------|--|---------|--------|---------|---------|-------|
| | 2007 | | | 2008 | | |
| | Upper | Lower | Total | Upper | Lower | Total |
| Broadcast spraying | | | | | | |
| HC | 0.466 a | 0.029 a | 0.495 | 6.289 c | 0.177 a | 6.466 |
| ASC | 0.301 a | 0.020 a | 0.321 | 2.203 b | 2.601 c | 4.804 |
| Band spraying | | | | | | |
| FF | 16.065 b | 2.090 b | 18.155 | 1.370 a | 1.685 b | 3.055 |

* Different letters in the same column indicate significant differences ($P < 0.05$) by Duncan test.

Table 4. Average pesticide deposit on ground.

| Pesticide application methods | Average pesticide deposit ($\mu\text{g}/\text{cm}^2$) | |
|-------------------------------|---|-------|
| | 2007 | 2008 |
| Broadcast spraying | | |
| HC | 0.906 | 0.654 |
| ASC | 0.989 | 0.789 |
| Band spraying | | |
| FF | 2.423 | 1.101 |




Figure 1. Spray drift in horizontally.

Parameters for pesticide decline of strawberries were shown in Table 8. The equations were used for other parameters such as K_{dec} , R_0 , R_p , etc. These parameters led to determination of the consuming day of strawberries following pesticide application. The MRL of dicofol on strawberry fruit was 0.02 mg/kg (EC, 2000). In all experiments when the recommended rate of dicofol was applied, the residue levels were above MRL after approximately 1 day in ASC, 2 days in FF and 4 days in HC (Table 8). On the spraying day in 2007 and 2008,

residues on strawberries in all applications were higher than the MRL with 1.74 mg/kg and 7.39 mg/kg for HC, 1.21 mg/kg and 6.14 for FF, and 1.10 and 3.98 for ASC, respectively. When HC were used, residues reached the MRL value in approximately 4 days and 7.5 days after spraying in 2007 and 2008. On the other hand, when ASC and FF were used, their residues reached the MRL value earlier than HC: approximately 1 day for ASC and 2 days for FF in 2007 and, approximately 5 days for ASC and FF in 2008. The residues on strawberries were below

Table 5. Average pesticide deposit on leaf surfaces of strawberry plants.

| Pesticide application methods | Average pesticide deposit ($\mu\text{g}/\text{cm}^2$)* | |
|-------------------------------|--|----------|
| | 2007 | 2008 |
| Broadcast spraying | | |
| HC | 0.777 b | 0.726 bc |
| ASC | 0.329 a | 0.585 ab |
| Band spraying | | |
| FF | 0.929 b | 1.223 c |
| Control | 0.004 a | 0.096 a |

* Different letters in the same column indicate significant differences ($P < 0.05$) by Duncan test.

Table 6. Pesticide residues on strawberries following pesticide application (Mean values \pm SD of four replicates).

| Year | Pesticide application methods | Active Ingredient (mg/kg) (Days after treatment) | | | | | | |
|------|-------------------------------|--|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|
| | | 0 | 1 | 2 | 3 | 4 | 5 | 6 |
| 2007 | Broadcast spraying | | | | | | | |
| | HC | 0.0347 \pm 0.000 | 0.0269 \pm 0.005 | 0.0261 \pm 0.002 | 0.0218 \pm 0.008 | 0.0204 \pm 0.002 | 0.0175 \pm 0.007 | 0.0164 \pm 0.003 |
| | ASC | 0.0218 \pm 0.000 | 0.0225 \pm 0.002 | 0.0196 \pm 0.009 | 0.0159 \pm 0.006 | 0.0143 \pm 0.001 | 0.0134 \pm 0.002 | 0.0086 \pm 0.001 |
| | Band spraying | | | | | | | |
| | FF | 0.0241 \pm 0.011 | 0.0232 \pm 0.007 | 0.0214 \pm 0.007 | 0.0198 \pm 0.003 | 0.0148 \pm 0.003 | 0.0136 \pm 0.006 | 0.0130 \pm 0.002 |
| | Untreated control | 0.0071 \pm 0.001 | 0.0062 \pm 0.001 | 0.0062 \pm 0.001 | 0.0060 \pm 0.001 | 0.0056 \pm 0.001 | 0.0052 \pm 0.002 | 0.0051 \pm 0.001 |
| 2008 | Broadcast spraying | | | | | | | |
| | HC | 0.1479 \pm 0.028 | 0.1044 \pm 0.007 | 0.0611 \pm 0.020 | 0.0564 \pm 0.021 | 0.0581 \pm 0.013 | 0.0353 \pm 0.012 | 0.0189 \pm 0.003 |
| | ASC | 0.1228 \pm 0.006 | 0.0812 \pm 0.006 | 0.0618 \pm 0.006 | 0.0412 \pm 0.004 | 0.0280 \pm 0.006 | 0.0148 \pm 0.003 | 0.0055 \pm 0.004 |
| | Band spraying | | | | | | | |
| | FF | 0.0797 \pm 0.012 | 0.0592 \pm 0.003 | 0.0489 \pm 0.006 | 0.0348 \pm 0.004 | 0.0266 \pm 0.001 | 0.0184 \pm 0.001 | 0.0069 \pm 0.002 |
| | Untreated control | 0.0873 \pm 0.010 | 0.0202 \pm 0.006 | 0.0088 \pm 0.001 | 0.0061 \pm 0.001 | 0.0022 \pm 0.001 | 0.0011 \pm 0.001 | 0.0002 \pm 0.001 |

the MRL value in 1 (for ASC), 2 (for FF), and 4 (for HC) days after spraying day in 2007 and, approximately 5 (for ASC and FF) and 7.5 (for HC) days after spraying day in 2008. On the other hand, the MRL of dicofol on strawberry fruit is

1.00 mg/kg and suggested consuming day is 7 days after pesticide application according to WHO and FAO. In this study, the residues on strawberries were below the legal limits (MRL) value (according FAO and WHO) in 1 day after

spraying in 2007 and 2008.

When HC, FF, and ASC were used, the suggested consuming days after pesticide application for these nozzles were 4.1 – 7.5 for HC, 2.2 – 5.2 for FF, and 0.9 – 5.1 days for ASC.

Table 7. Average pesticide residues on strawberries.

| Pesticide application methods | Average pesticide residues (mg/kg)* Mean±SD | |
|-------------------------------|---|----------------|
| | 2007 | 2008 |
| Broadcast spraying | | |
| HC | 0.0234 ± 0.007 c | 0.0689±0.043 d |
| ASC | 0.0166 ± 0.006 b | 0.0508±0.039 c |
| Band spraying | | |
| FF | 0.0186 ± 0.006 b | 0.0383±0.025 b |
| Untreated control | 0.0059 ± 0.001 a | 0.0180±0.029 a |

*Means with the same letter are not significantly different at the 5% level of least significant difference.

Table 8. Parameters for pesticide decline of strawberries.

| Pesticide application methods | Equation | R ² | K _{dec} (days ⁻¹) | R ₀ (mg/kg) | R _p (mg/kg) | Decline R ₀ to R _p (%) | Suggested consuming day after pesticide application according to MRL (0.02) |
|-------------------------------|---------------------------|----------------|--|------------------------|------------------------|--|---|
| 2007 | | | | | | | |
| Broadcast spraying | | | | | | | |
| HC | $R = 0.0326.e^{-0.1194t}$ | 0.97 | 0.1194 | 0.0326 | 0.0159 | 51.2 | 4.1 |
| ASC | $R = 0.0226.e^{-0.1353t}$ | 0.88 | 0.1353 | 0.0226 | 0.0100 | 55.8 | 0.9 |
| Band spraying | | | | | | | |
| FF | $R = 0.0257.e^{-0.1152t}$ | 0.95 | 0.1152 | 0.0257 | 0.0128 | 50.2 | 2.2 |
| 2008 | | | | | | | |
| Broadcast spraying | | | | | | | |
| HC | $R = 0.1905.e^{-0.2997t}$ | 0.92 | 0.2997 | 0.1905 | 0.0315 | 83.5 | 7.5 |
| ASC | $R = 0.2376.e^{-0.4826t}$ | 0.95 | 0.4826 | 0.2376 | 0.0131 | 94.5 | 5.1 |
| Band spraying | | | | | | | |
| FF | $R = 0.1345.e^{-0.3674t}$ | 0.92 | 0.3674 | 0.1345 | 0.0148 | 89 | 5.2 |

R: Residue (mg/kg); t: Time (day); R²: Coefficient of determination; K_{dec}: Decline constant (days⁻¹); R₀: Initial residue (mg/kg); R_p: Residues at pre-harvest interval of pure a.i. (mg/kg).

Therefore, we suggest that strawberries can be harvested after a minimum of 5 days when the spraying was done with HC, and after 5 days with FF and ASC 5 days when the spraying was done with HC, FF and ASC. The results that this study yielded can well be utilized in the cultivation of strawberries in specific, and in other crops and fruits in general.

Biological efficiency

Population trends in abundance of red spider mites were very similar in 2007 and in 2008 with detections of some unimportant fluctuations in plots of different spraying methods and control plots in both years (Figures 2 and 3). Mean numbers of eggs and mobile mites regularly

increased at the beginning of first weeks of April in 2007. Their numbers increased up to the highest population densities on same sampling date, May 8, 2007 (Figure 2). In 2008, earlier infestations of red spider mites in all the plots were detected on 24 May and numbers of eggs and MM stage started to increase after March 31 (Figure 3). Their numbers peaked on April 21 or April 28. There was a statistically meaningful difference among the treatments for numbers of both eggs and MM on day 3 after spraying dicofol, with detection of the similar and significantly lower numbers of RSM in HC, ASC and FF compared to numbers found in control plot in 2007 (Table 9) (Egg: F = 27.909; d.f. = 3, 156; P < 0.001; MM: F = 27.909; d.f. = 3, 156; P < 0.001). Day 6 and day 10 after spraying, mean numbers of eggs (F = 20.642; d.f. = 3, 156, P < 0.01 and F = 6.453; d.f. = 3, 156, P < 0.01, respectively) and MM in




Figure 2. Mean numbers of *T. cinnabarinus* on strawberry leaflets in plots of HC (a) FF (b) ASC (c) and control (d) in 2007.




Figure 3. Mean numbers of *T. cinnabarinus* on strawberry leaflets in plots of HC (a) FF (b) ASC (c) and control (d) in 2008.

plots of HC and ASC were similar and significantly larger than FF ($F = 178.292$; d.f. = 3, 156, $P < 0.01$ and $F = 29.318$; d.f. = 3, 156, $P < 0.01$, respectively). The numbers of eggs and mobile stages apparently declined to low levels in all plots and this was probably due to rainfall. On day 14, numbers of red spider mites were

lower in plots characterized by the different spraying techniques and also in control plot. However, number of eggs and MM was significantly lower in band spraying than those found in others (Eggs: $F = 2.297$; d.f. 3, 156; $P = 0.033$; MM: $F = 2.207$; d.f. 3, 156; $P = 0.037$, respectively). Similar to previous experimental year, in

Table 9. Results of experiments to control red spider mite, *T. cinnabarinus* Boids., on strawberry plantations, with different acaricides (results before treatment and after applications, 2007)^a.

| Treatment | Pre-treatment | | | | Time application (day, date) | | | | | | | | | | |
|-----------|-------------------|--------|----------------|---------|------------------------------|--------|--------|-----------------|--------|--------|-----------------|-------|--------|------|--|
| | (8 May) | | Day 3 (11 May) | | Day 6 (14 May) | | | Day 10 (18 May) | | | Day 14 (18 May) | | | | |
| | MM | E | MM | E | CM | MM | E | CM | MM | E | CM | MM | E | CM | |
| HC | 82.2 ^b | 95.3a | 27.7 b | 57.3 c | 56.3 | 7.8 b | 44.2 a | 63.5 | 9.7 b | 8.6 b | 61.0 | 2.0 a | 2.4 a | 50.0 | |
| ASC | 90.2 | 108.3a | 38.3 b | 93.0 b | 44.7 | 9.4 b | 21.2 b | 46.4 | 10.0 b | 11.3 b | 43.2 | 1.8 a | 2.1 a | 47.8 | |
| FF | 81.5 | 103.2a | 35.2 b | 61.3 c | 54.5 | 4.1 c | 26.6 b | 70.3 | 1.2 c | 4.4 c | 83.0 | 0.8 b | 0.6 b | 62.0 | |
| Control | 89.0 | 106.5a | 102.0 a | 136.7 a | | 46.3 a | 53.2 a | | 20.0 a | 18.2 a | | 2.5 a | 1.1 ba | | |

^a Numbers of mobile mites (MM) or eggs per leaflet, or corrected percentage mortality (CM); ^b means in the same column followed by the same letter do not differ significantly ($P < 0.05$) by Tukey's honest test.

Table 10. Results of experiments to control red spider mite, *T. cinnabarinus* Boids., on strawberry plantations, with different acaricides (results before treatment and after applications, 2008)^a.

| Treatment | Pre-treatment | | | | Time application (day, date) | | | | | | | | | | |
|-----------|---------------|-------|------------------|-------|------------------------------|-------|--------|-------------------|-------|--------|----------------|-------|--------|------|--|
| | (14 April) | | Day 3 (18 April) | | Day 6 (21 April) | | | Day 10 (25 April) | | | Day 14 (6 May) | | | | |
| | MM | E | MM | E | CM | MM | E | CM | MM | E | CM | MM | E | CM | |
| HC | 38.8a | 51.8a | 13.7b | 65.4b | 21.2 | 17.8b | 76.4b | 27.4 | 22.6b | 56.1b | 51.1 | 29.6c | 51.7b | 48.3 | |
| ASC | 49.6a | 78.5a | 15.3b | 60.6b | 36.8 | 15.7b | 65.8b | 49.7 | 29.1b | 54.5b | 56.4 | 47.7b | 86.8a | 28.7 | |
| FF | 32.5a | 62.2a | 14.0b | 69.0b | 22.7 | 10.2c | 48.5c | 62.4 | 14.0c | 34.3c | 72.7 | 25.9c | 48.3b | 56.5 | |
| Control | 35.0a | 66.2a | 41.5a | 80.9a | | 62.1a | 113.8a | | 70.4a | 129.0a | | 90.0a | 102.8a | | |

^a Numbers of mobile mites (MM) or eggs per leaflet, or corrected percentage mortality (CM); ^b means in the same column followed by the same letter do not differ significantly ($P < 0.05$) by Tukey's honest test.

2008 significantly clear differences in numbers of both eggs and MM among the treatments were found on day 3 (Table 10). Mean numbers of MM in FF were significantly lower than numbers found in HC and ASC treatments and in control on day 6, 10 and 14 ($F = 28.193$; d.f. 3, 156; $P = 0.010$; $F = 21.280$; d.f. 3, 156; $P < 0.001$ and $F = 24.394$; d.f. 3, 156; $P = 0.001$, respectively). Mean numbers of eggs in FF were significantly lower than numbers found in HC and ASC treatments, in control on day 6, 10 and 14 after spraying ($F = 3.178$; d.f. 3, 156; $P = 0.010$; $F = 6.335$; d.f. 3,

156; $P < 0.001$ and $F = 5.547$; d.f. 3, 156; $P = 0.001$, respectively).

Dicofol did not effectively control the RSM populations in two study-years. The corrected mortality rates in band spraying were greater with rates of 70.3 and 83.0% on days 6 and 10 in 2007, respectively. These were followed by the rates of 63.5 and 61.0% obtained in hollow cone treatment on days 6 and 10, respectively. In 2008 the highest corrected mortality rate was obtained in FF with rate of 72.7% on day 10.

Dicofol was not able to suppress the RSM

populations effectively in both years. Mortality rates in RSM populations in the plots ranged from 42 to 83% in 2007 and 21 to 72% in 2008. Dicofol had given excellent control of *T. cinnabarinus* on strawberry in the same ecological region of Turkey (Yi it and Erki liç, 1992). In this study, it is found that lower efficacy of dicofol in controlling the mite populations is due to a resistance to dicofol. Some previous studies indicated that *T. urticae* developed resistance to dicofol in various crops (Dennehy et al., 1988; Ferguson-Kolmes et al., 1991; Wilson et al., 1995). In Turkey dicofol

resistance to *T. cinnabarinus* in cotton has been reported by Da li and Tunç (2001).

Mean numbers of eggs and mobile stages (pooled) in band spraying were significantly lower compared to numbers obtained in the treatments of the other two spraying techniques in both years (Tables 9 and 10). HC and ASC were found to be less effective in reducing the population densities of both eggs and MM of RSM. In the present study, it is found that detection of the lower mortality rates in plots of HC and ASC are due to lower deposits of dicofol on the leaf surfaces compared with FF. Different results between this study and previous ones may be related to different factors. In present study, greater spray drifts in horizontally distances were measured from ASC. Smaller drops of <80 µm can cause the spray drifts (Miller, 1993).

Although dicofol did not exhibit desirable control of this pest mite in both years, higher corrected mortality rates of RSM were recorded in the plots of band spraying. Band spraying may effectively reduce the total amount of pesticides used on per hectare basis, which, in turn, reduces costs and minimizes the potential for pesticide carryover, allowing the survival of the arthropods including beneficial insects inhabiting the untargeted area, that is, refugia and field margins. Field margins bearing weedy plants have ecologically important roles in conservation and augmentations of the natural enemies (Welling et al., 1988; Boatman and Wilson, 1988; Boatman, 1994; Denys and Tscharrntke, 2002).

Conclusions

1. Both highest pesticide deposits on leaf surfaces and best biological efficiency was obtained from FF nozzles in 2007 and 2008.
2. Average pesticide deposits on upper leaf surfaces were found to be higher than average pesticide deposits on lower leaf surfaces of strawberries in all pesticide application methods in 2007 and 2008.
3. The highest spray drift was obtained from ASC in all pesticide application methods in 2007 and 2008.
4. The highest pesticide deposits on ground were obtained from FF in 2007 and 2008.
5. When HC, FF, and ASC were used, the suggested consuming days after pesticide applications for these nozzles were 4.1 – 7.5 for HC, 2.2 – 5.2 for FF, and 0.9 – 5.1 days for ASC. Therefore, we suggest that strawberries can be harvested after 5 days when the spraying was done with HC, FF and ASC.

ACKNOWLEDGEMENTS

The work represented part of a research carried out supported by The Scientific and Technical Research Council of Turkey (TUBITAK TOVAG 106 O 757). This

support is greatly acknowledged. The authors wish to thank Prof. N. Kemal Koc, Head of Subtropical Fruits Research and Experimental Center, University of Cukurova, for allowing access to use the laboratory during our experimental studies; Prof. Halil Elekcioğlu, Head of Department of Crop Protection, Yalex Co. for allowing access to strawberries field; Mehmet Yaltir, Mehmet Ali Unlu and Arzu Kiminsu. Invaluable technical assistance, in designing the spray boom was given by Mr. Fevzi Sahbaz and Mr. B. Mert Kaplan. For field and laboratory studies, special thanks to Miss. Melike Arslan and Mr. Fevzi Sahbaz. The authors wish to thank Mrs. Asli Abdulvahitoglu for her effort in correcting the English text.

REFERENCES

- Bayat A, Yarpuz-Bozdogan N (2005). An air-assisted spinning disc nozzle and its performance on spray deposition and reduction of drift potential. *Crop Prot.*, 24: 951-960.
- Boatman ND, Wilson PJ (1988). Field edge management for game and wildlife conservation. *Aspects Appl. Biol.*, 16: 56-61.
- Boatman N (1994). Field margins – Integrating agriculture and conservation. British Crop Protection Council monograph no.58. British Crop protection Council, Farnham.
- Cross JV, Berrie AM, Murray RA (2000). Effect of drop size and spray volume on deposits and efficacy of strawberry spraying. *Aspects Appl. Biol.*, 57: 313-320.
- Da li F, Tunç I (2001). Dicofol resistance in *Tetranychus cinnabarinus*: resistance and stability of resistance in populations from Antalya, Turkey. *Pest Mgt. Sci.*, 57: 609-614.
- Debear C, Broers N, Denruyter L, Jaeken P (2005). Effect of band spraying on deposit and biological efficacy in strawberries. VIII workshop on spray application techniques in fruit growing. Barcelona, Spain, p. 8.
- Dennehy TJ, Nyrop Reissig WP, Weires RW (1988). Characterization of dicofol in spider mites (Acari: Tetranychidae) from New York apple orchards. *J. Econ. Entomol.*, 81: 1551-1561.
- Denys C, Tscharrntke T (2002). Plant -insect communities and predator-prey ratios in field margin strips, adjacent crop fields, and fallows. *Oecol.*, 130: 315-324.
- Duncan DB (1955). Multiple range and multiple F tests. *Biometrics*, 11: 1-42.
- Ferguson-Kolmes LA, Scot JG, Dennehy TJ (1991). Dicofol resistance in *Tetranychus urticae* (Acari: Tetranychidae): Cross-resistance and Pharmacokinetics. *J. Econ. Entomol.*, 84: 41-48.
- EC (2000). Official Journal of the European Communities, Commission Directive 2000/42/EC Official Bulletin, p. 25.
- Gan-Mor S, Grinstein A, Beres H, Yehudit R, Zur I (1996). Improved uniformity of spray deposition in a dense plant canopy: Methods and equipment. *Phytoparasitica*, 24: 57-67.
- Goodwin S, Ahmad N, Newell G (1984). Dimethoate spray residues in strawberries. *Pesticide Sci.*, 16: 143-146.
- Guardia-Rubio M, Marchal -Lopez RM, Ayora-Canada MJ, Ruiz-Medina A (2007). Determination of pesticide in olives by gas chromatography using different detection systems. *J. Chromatogr. A*, 1145: 195-203.
- Henderson CF, Tilton EW (1955). Test with acaricides against the brown wheat mite. *J. Econ. Entomol.*, 48: 157-161.
- Holownicki R, Doruchowski G, Godyn A, Swiechowski W (2000). Variation of spray deposit and loss with air-jet directions applied in orchards. *J. Agric. Engr. Res.*, 77: 129-136.
- Kilmer R, Andre AM, Stevens TJ (2001). Pesticide residues and vertical integration in Florida strawberries and tomatoes. *Agribus.*, 17: 213-226.
- Labanowska BH, Doruchowski G, Goszczy ski W, Godyn A (1996). Comparison of two spraying techniques in the control of twospotted spider mite (*Tetranychus urticae*) and grey mold (*Botrytis cinerea*) on

- strawberry. *Acta Horticult.* (ISHS), 422: 387-388.
- Landers A (2007). Modern technology to improve spraying efficiency. Online: <http://www.aben.cornell.edu/extension/pestapp/publications/mod.tech.html>. 22/10/2007.
- Manor G, Hofner R, Or R, Phishler G, Epstein Y, Nakash T, Jacobi M (1989). Air Stream Facilitated Application of Cotton Foliage Treatments. *Trans. ASAE*, 32(1): 37-40.
- Matthews GA (2000). *Pesticide Application Methods*. Third Edition. Blackwell Science Ltd. UK, p. 432.
- Miller PCH (1993). *Application Technology for Crop Protection*. CAB Int., pp. 101-122.
- Nelson RD, Show ED (1975). A novel group of miticides containing the cyclopropane moiety: laboratory experiments and field studies in strawberries on the two spotted spider mite. *J. Econ. Entomol.*, 68: 261-266.
- Nordmark L (1994). Spray technology for improving in field vegetables and strawberries. *Acta Horticult.*, 372: 103-110.
- Oncag G, Cengiz F (1978). Ege bölgesi çilek alanlarında böcek faunası (Zararlı ve faydalı) tesbiti üzerinde ara tırmalar. *Zirai Mücadele Ara tırma Yılı ı*, p. 61. (In Turkish).
- Ozkan HE, Derksen RC (1998). Effectiveness of Turbdrop® and Turbo Teejet® Nozzles in Drift Reduction. *AEX*, 524-98.
- Panneton B, Phillion H, Theriault R (1996). Spray chamber of air assisted spraying. *ASAE Paper No. 96-1078*, ASAE, St. Joseph, MI 49085.
- Panneton B, Phillion H, Theriault R, Khelifi M (2000a). Spray chamber evaluation of air-assisted spraying on broccoli. *Crop Sci.*, 40: 444-448.
- Panneton B, Phillion H, Theriault R, Khelifi M (2000b). Spray chamber evaluation of air-assisted spraying on potato plants. *Trans. ASAE*, 43: 529-534.
- Pesticide Analytical Manual (1999). E1 Extraction with acetone, liquid-liquid partitioning with petroleum ether/methylene chloride, 1: 302.
- PISC (Primary Industries Standing Committee) (2002). *Spray Drift Management, Principles, Strategies and Supporting Information*. Australya, CSIRO Publishing, p. 71.
- Rowinski RS, Arvidsson T, Hagenvall H (1994). A Standart Method for the Investigation of Drift. *AgEng*, International Conference on Agricultural Engineering. 29th – 1st September, pp. 722-723.
- Safi JM, Abou-Foul NS, El-Nahhal YZ, El-Sebae AH (2002). Monitoring of pesticide residues on cucumber, tomatoes and strawberries in Gaza Governorates, Palestine. *Nahrung/Food*, 46: 34-39.
- SPSS (1993). *SPSS for Windows Release, 6.0* (June 17, 1993). Copyright SPSS Inc., 1989–93, New York, USA.
- Stensvand A, Christiansen A (2000). Investigation on fungicide residues in greenhouse-grown strawberries. *J. Agric. Food Chem.*, 48: 917-920.
- Sumner HR, Herzog GA (2000). Assessing the Effectiveness of Air-Assisted and Hydraulic Sprayers in Cotton via Leaf Bioassay. *J. Cotton Sci.*, 4: 79-83.
- Taylor WA, Andersen PG (1989). The use of air-assistance in field crop sprayer to reduce drift and modify drop trajectories. *Proc. Brighton Crop Prot. Conf. Weeds*, 2: 631-639.
- Yenisoy-Karakas S (2006). Validation and uncertainty assessment of rapid extraction and clean-up methods for the determination of 16 organochlorine pesticide residues in vegetables. *Anal. Chicia Acta*, 571: 298-307.
- Yi it A, Erkiliç L (1992). Studies on the chemical control of *Tetranychus cinnabarinus* Boisd. (Acarina: tetranychidae) a, pest of strawberry in the east Mediterranean region of Turkey. *Crop Prot.*, 11: 433-438.
- Welling M, Pötzl RA, Jürgens D (1988). Untersuchungen in Hessen über Auswirkung und Bedeutung von Ackerschonstreifen. 3. Epigäische Raubarthropoden. *Mitt. Biol. Bundesanst.*, 247: 55-63.
- Wilson LJ, Herron GA, Leigh TF, Rophail J (1995). Laboratory and field evaluation of the selective acaricides dicofol and propargite for control of *Tetranychus urticae* Koch (Acari: Tetranychidae). in Australian cotton. *J. Austr. Entomol. Society*. 34: 247-252.
- Wolf R (1997). *Equipment to Reduce Spray Drift*. The Cutting Edge. 97-7.