

Full length Research Paper

Sequence analysis of the *groEL* gene and its potential application in identification of pathogenic bacteria

Hu Yushan^{1,2}, Luo Lei^{1,2}, Liu Weijia^{1,2} and Chen Xiaoguang^{1*}

¹Department of Pathogenic Biology, School of Public Health and Tropical Medicine, Southern Medical University, Guangzhou 510515, China.

²The Center for Disease Control and Prevention of Guangzhou, Guangzhou 510080, China.

Accepted 22 July, 2017

This study aims to develop a rapid method for identifying pathogenic bacteria based on the sequence difference of pathogenic bacteria *groEL* gene. A pair of universal degenerate primers was designed to amplify the *groEL* genes of 34 major species of 16 genus's of pathogenic bacteria. The conservation, variation and the interspecies phylogenetic relations of the *groEL* gene sequences were informatically analysed. Our results showed that the *groEL* gene fragments of the selected pathogenic bacteria could be amplified using the degenerate primer. The results of the sequence comparative study indicated that the *groEL* gene is characterized by conservation and variation, while the variation regions are distributed separately among the conserved regions, just like the mosaic. This conservation and variation coexistence of pathogenic bacteria *groEL* genes could provide a solid basis for the rapid identification of pathogenic bacteria and the development of a diagnostic microarray.

Key words: *groEL*, homology, phylogenesis.

INTRODUCTION

Although most infectious diseases were controlled to a low level with the development of medical science, the new and re-emerging infectious diseases continue to appear. And infectious diseases remain the major cause of death worldwide and the leading cause of illness (Franklin et al., 2004). In order to control infectious diseases more effectively, laboratory methods for the rapid and accurate identification of emerging and reemerging pathogens must be presented. The routine approaches for pathogen identification are cultivation, biochemistry reaction and serologic or immunological test. Those approaches are trivial, time-consuming and labor-intensive, especially for those slow-growing and hard-to-culture pathogens. Recently, typing methods began to evolve from phenotype-based methods toward genotype-based methods, including restriction endonuclease analysis (REA) of the total bacterial genome, pulsed-field gel electrophoresis (PFGE), arbitrarily primer PCR (AP-PCR) and PCR ribotyping (George et al.,

2008). Ribosomal 16S rRNA, 23S rRNA, *slpA*, *selC*, and the *pheU* tRNA locus are usually selected as the target genes for successful ribotyping subgroups of bacteria (Gelsen et al., 1994; Gurtler et al., 1996; Anthony et al. 2000; Hayden et al., 2001; Corinne et al., 2008; Ghebremedhin et al., 2008). The *groEL* gene, which encodes a 60 kDa subunit (known as HSP60, 60 kDa chaperonin and heat shock protein), also has the potential to serve as a general phylogenetic marker because of its ubiquity and conservation in nature. The heat shock protein is of great importance for maintaining cellular normal physiological function and is also the major antigen protein of many bacterial pathogens. It has been well documented that the *groEL* gene is one of the most conservative systems in nature (Kwok et al., 1999; Rebecca et al., 2002; Karuna et al., 2003; Jung-Hee et al., 2003; Yu-Hsiu et al., 2003; Eric et al., 2004; Sensus et al., 2004; Giuseppe et al., 2008). Despite the conserved nature of the *groEL* gene, the level of interspecies variation of *groEL* sequence is greater. Therefore, the *groEL* gene may be a good target gene for species classification. In the present study, *groEL* genes of 34 major species of 16 genus bacterial pathogens were analyzed for homology, variation and gene distribution.

*Corresponding author. E-mail: xgchen@fimmu.com,
xgchen2001@hotmail.com.

This study may provide better resolution for the rapid identification of pathogenic bacteria.

MATERIALS AND METHODS

Bacterial strains

34 species of 16 genera of bacterial pathogens of medical interest were selected, and their *groEL* gene sequences were extracted from GenBank. The bacterial strains used in this study are listed in Table 1.

The conservation analysis of *groEL* gene

Given conservative sequence length parameter and variation value, *groEL* gene sequences of the bacteria strains were edited and compared to find out the conservative region of the gene and analyze conservative distribution according to bio-software ClustalX and Bio-edit.

Homology analysis of *groEL* gene

The *groEL* gene sequences of bacteria strains were analyzed for homology using bio-software DNASTar. The phylogenetic tree was drawn based on the homology analysis of *groEL* gene. The phylogenetic relationship among the bacterial classification and the importance of rapid diagnosis were analyzed at the same time.

The variation analysis of *groEL* gene

The variation analysis of *groEL* gene was conducted based on the homology analysis of *groEL* gene. The map of conservative sequence and variant sequence distribution was drawn accordingly. Furthermore, the variation rule of *groEL* gene sequence was analyzed.

PCR amplification of *groEL* gene

Using bio-software primer premier 5.0 (PREMIER Biosoft International, USA), a proper pair of universal primers were designed based on the homology analysis of the conserved sequence of *groEL* gene. Because of the lower homology within the region, a degenerate upstream primer was designed. The sequences of primers were P1: -5'AGTTACCCT XGG YCCZ AAAG-3', X is C or T, Y is T or C, Z is A or G; P2: -5'CAGCAAC-CACGCCTTCTTC-3', with an amplified fragment of approximately 600 bp in length. The template used for amplification was prepared as follows: bacteria grown in nutrient broth for 24 h were plated on selective agar plate and single colony was picked up after cultivation for 24 – 36 h. Bacteria suspension was made with 100 μ l of re-distilled water, heated in 95°C for 10 min, centrifugated at 12000 rpm for 1 min and the supernatant was used as PCR template. The PCR amplification was performed in 50 μ l reaction volume of mixture containing 5 μ l of 10 \times PCR buffer, 4 μ l of 20 mM dNTP, 3 μ l of 2 mM, MgCl₂, 1 μ l of 25 pmol forward and reverse primers, 5 μ l of template DNA, and 1 μ l of 5U *Taq* DNA polymerase (Takara Bio). The condition for PCR thermal cycling was as follows: the PCR mixture was held at 95°C for 4 min prior to the 35 cycles of PCR amplification in a thermal cycler (Eppendorf, Hamburg, Germany), the first 5 cycles are consisted of denaturation at 94°C for 40 s, annealing at 46°C for 50 s and extension at 72°C for 1.5 min and followed the other 30 cycles: denaturation at 94°C for 30 s, annealing at 53°C for 45 s and extension at 72°C for 1.5 min.

RFLP analysis of *groEL* gene

The *groEL* genes were amplified, purified and then digested with restriction enzymes Hae III (Takara Bio). The digested products were electrophoresed with a 1% TBE agarose gel in 0.5x TBE and visualized under UV light.

Hybridization analysis of *groEL* gene in oligonucleotide membrane array

The nucleic acid hybridization oligonucleotide probes were selected from the target region of different species, respectively (Table 2). All probes were synthesized in Takara Bio. Positive charge nylon membrane (Takara Bio) was used as array base. A 3 \times 3 mm grid was formed with blunt pencil. The nylon membrane was immersed in distilled water for 10 min and then dried in Whatman paper. Each probe was suspended to make a 25 pmol/ μ l solution. The solution was heated at 95°C for 3 min and 1 μ l of the solution was spotted on a corresponding position of the nylon membrane. Membranes with the probes were immersed in a Petri dish containing 0.5 ml DIG Easy Hyb solution (Roche, Indianapolis, USA) pre-warmed at hybridization temperature and pre-hybridization was performed at 50°C for 30 min with gentle shaking. 5 μ l PCR products were heated at 95°C for 5 min, immediately cooled on ice and then added to newly pre-warmed hybridization solution. The membranes were hybridized in the solution at 50°C for 2 h with gentle shaking. After hybridization, the membranes were washed 4 times with 0.25 \times SSC- 0.1% SDS for 2 min at 37°C. Before blocking, the membranes were washed in washing buffer (Roche) for 1 min, then immersed in 10 ml blocking buffer (Roche) for 30 min, and put to react with anti-digoxigenin antibody for another 30 min with gentle shaking. Color development was made with NBT-BCIP for 30–60 min in the dark without shaking. The reaction was stopped with tap water. The resulting images were visible and photographed.

RESULTS

The conservative sequences of *groEL* gene

The *groEL* gene sequences of pathogenic bacteria were analyzed by the bio-software ClustalX and Bioedit for the presence of the conservative regions in the genomes. It was found that the distribution of the conserved regions showed a characteristic pattern, that is, the mutant regions inserted into the conserved regions, just like mosaic (Figure 1). The conserved regions were mainly distributed in C region (nt900 - nt1500) and D region (nt1500 - nt1900), in which most conserved regions with high conservation were distributed in D region.

Variation of *groEL* gene

By software analysis, the variation map of *groEL* gene was obtained (Figure 1). As demonstrated in this map, it is evident that the *groEL* gene appears to be conserved. However, there are still considerable variations exist, especially in the A region (nt1- nt600). And these variations are found to be inserted among the conserved

Table 1. Bacteria strain resources used in the study.

Bacteria strain	Genbank accession No.	Source
<i>E. coli</i>	M11294	Guangzhou CDC
<i>S. enterica</i>	AY044102	Guangzhou CDC
<i>S. typhi</i>	U01039	Guangzhou CDC
<i>S. typhimurium</i>	AY044105	Guangzhou CDC
<i>S. flexneri</i>	AY044103	FuTian CDC
<i>S. sonnei</i>	AY044104	FuTian CDC
<i>S. boydii</i>	AY044101	FuTian CDC
<i>E. aerogenes</i>	AF306521	NanShan CDC
<i>E. amnigenus</i>	AB008140	NanShan CDC
<i>E. asburiae</i>	AB008137	NanShan CDC
<i>K. pneumoniae</i>	U81143	FuTian CDC
<i>Listeria monocytogenes</i>	AY922346	FuTian CDC
<i>Vibrio parahaemolyticus</i>	AF230952	Guangzhou CDC
<i>Vibrio cholerae</i>	AF230940	Guangzhou CDC
<i>Yersinia enterocolitica</i>	X59367	ShenZhen CDC
<i>Yersinia pestis</i>	AB064592	ShenZhen CDC
<i>Campylobacter jejuni</i>	AY628401	ShenZhen CDC
<i>Helicobacter pylori</i>	AJ558222	ShenZhen CDC
<i>Aeromonas hydrophila</i>	AY922365	GuangDong CDC
<i>Aeromonas jandaei</i>	AY922357	GuangDong CDC
<i>Mycobacterium marinum</i>	AF271346	GuangDong CDC
<i>Brucella melitensis</i>	L09273	GuangDong CDC
<i>S. aureus</i>	AF053568	GuangDong People Hospital
<i>S. epidermidis</i>	AF029245	GuangDong People Hospital
<i>S. haemolyticus</i>	U92809	GuangDong People Hospital
<i>S. saprophyticus</i>	AF029246	GuangDong People Hospital
<i>S. warneri</i>	AF053569	GuangDong People Hospital
<i>S. anginosus</i>	AY344497	GuangDong People Hospital
<i>S. bovis</i>	AY346155	FuTian CDC
<i>S. mutans</i>	AY344496	FuTian CDC
<i>S. pneumoniae</i>	AY344504	GuangDong People Hospital
<i>S. sanguinis</i>	AY344503	GuangDong People Hospital
<i>N. gonorrhoeae</i>	U64996	GuangDong People Hospital
<i>N. perflava</i>	AY837569	GuangDong People Hospital

regions, just like a mosaic with serrated appearance.

Homology of *groEL* gene

According to the comparative analysis of homology in different species of pathogenic bacteria, the phylogenetic tree was drawn using DNASTAR (Figure 2). The phylogenetic relationships derived from comparisons of the *groEL* sequences are presented in Figure 2. The phylogenetic analysis revealed that the sequences of the *groEL* genes from bacteria species were divided into 5 clusters. From this phylogenetic tree, it is apparent that the highest homology exists in the same genus of bacteria with closest interrelationship, such as *Shigella*, *Salmonella*

and *Staphylococcus* of the same genus. The homology of bacteria belonging to the same family is higher than those of bacteria of different families, such as, the interrelationship of *Salmonella* and *Shigella* in the same family is closer than that of *Salmonella* with *Mycobacterium* in the different family. However, intercross in interrelationship occurs among different genus of bacteria, such as *Staphylococcus* and *Streptococcus*. *Coccus* and *Bacillus* belong to different branch of bacteria. The phylogenetic tree also showed that the relationship between bacteria *groEL* genes and eukaryotic *groEL* genes is close. For example, *Aspergillus clavatus*, *Aspergillus fumigatus*, *Drosophila melanogaster* and *Mycobacterium marinum* are highly related. But *Cryptococcus neoformans* and *Plasmodium vivax* are relative far to

Table 2. Probes used in the study.

Probe name detection range	Probe sequence (5' to 3')	Genbank accession number
Universal probes	CTAAAGCGATTGCTCAGGTTG	X62914
<i>Salmonella</i> spp	TGAACCCGATGGACCTGAAACG	AY044102
<i>Salmonella enterica</i>	TGAACCCGATGGACCTGAAACG	AY044102
<i>Salmonella typhi</i>	AAGGCGTGGTTGCTGGTGGTGG	U01039
<i>Salmonella typhimurium</i>	CTCCGCTAACTCCGACGAAACC	AY044105
<i>Shigella</i> spp	ACCATCTCCGCTAACTCCGACG	AY044103
<i>Shigella flexneri</i>	TCTCCGCTAACTCCGACGAAA	AY044103
<i>Shigella sonnei</i>	ACCATCTCCGCTAACTCCGACG	AY044104
<i>Shigella boydii</i>	ACCATCTCCGCTAACTCCGACG	AY044101
<i>Streptococcus</i> spp.	CAAGTAGGTGCGATTTCTGC	AF053568
<i>S. aureus</i>	ATCGTGCTAAACCGTATGCGTG	AF053568
<i>S. haemolyticus</i>	AAACCGTATGCGTGGAAC	U92809
<i>S. epidermidis</i>	TAAACCGTATGCGTGGAAC	AF029245
<i>Streptococcus</i> spp.	GAGGATGCTCTAAATGCCACA	AY344504
<i>S. pneumoniae</i>	TCCAATCGGTATTGTCGTG	AY344504
Positive control	-	
Negative control	-	

Postive control is digoxigenin labeling plasmid *pBR328/BamHI*. Negative control is plasmid *pBR328/BamHI*.




Figure 1. Analysis of *groEL* gene divergence. There are 4 variant regions in *groEL* gene. A region is the most mutable one; D region is the least mutable one.




Figure 2. Phylogenetic tree of bacteria and eukaryotics *groEL* gene.

bacteria.

PCR amplification of *groEL* gene sequence

The *groEL* gene in intestinal pathogens, such as *E. coli*, *Salmonella enterica*, *Shigella flexneri*, *Klebsiella pneumoniae*, *Vibrio parahaemolyticus*, *Campylobacter jejuni* was successfully amplified by PCR and a DNA fragment of 600bp could be obtained from all the pathogenic bacteria (Figure 3), indicating that the universal degenerate primer can amplify the *groEL* gene effectively.

PCR-RFLP analysis of *groEL* gene products

The PCR-amplified products of *groEL* gene of *S. enterica*,

S. typhi, *S. typhimurium*, *S. flexneri*, *Shigella sonnei* and *Shigella boydii* were identified by RFLP analysis. It is apparent that the RFLP patterns were rather clear with definite DNA bands in size except *S. typhi* and *S. typhimurium* because of their similar patterns (Figure 4). So this method can be used for partially identification of pathogenic bacteria.

Hybridization analysis of *groEL* gene in oligonucleotide membrane array

The probes were designed, synthesized and dotted on the oligonucleotide membrane (Figure 5). The PCR products were then hybridized with probes on the membrane followed by color development under appropriate condition. Some bacterial species were tested with the




Figure 3. Amplification of *groEL* gene fragment of some pathogenic bacteria. Lanes 1–6 are PCR products. 1: *E. coli*; 2: *S. enterica*; 3: *S. flexne*; 4: *K. pneumonia*; 5: *V. parahaemolyticus*; 6: *C. jejuni*; Lane M: DNA molecular marker.

oligonucleotide array method. The results showed that high sensitivity and specificity of hybridization results were obtained with these species of bacteria, including *S. flexneri*, *S. sonnei*, *S. pneumoniae*, *S. aureus*, *Staphylococcus haemolyticus*, *Staphylococcus epidermidis*, *S. enterica*, *S. typhimurium*, *S. typhi*. As to *S. boydii*, we found poor signals because of the low amount of oligonucleotide dotted on the membrane (Figure 6).

DISCUSSION

The traditional methods involved in the discovery of the infective agents are cultivation, isolation, biochemical identification and the serological tests to the suspected pathogens. However, these methods are more or less

trivial and time-consuming with a rather low detection rate. This is disadvantageous especially during the outbreak of an epidemic infection when the pathogens should be rapidly identified to meet the requirement for effective control of infection or disease. Molecular biological technology provides a better resolution in the rapid identification of pathogens.

At present, various DNA-based techniques have been developed to identify pathogenic bacteria, such as PCR-RFLP, PCR-SSCP, APPCR, PFGE, multiple PCR and PCR aiming at specific gene. In addition, the high through-out biochip is now being available for rapid identification of microbes (Hyun-Hee et al., 2002; Bangxing et al., 2004; Carl et al., 2004; Gerard et al., 2004; Kensuke et al., 2004; Rousseaux et al., 2004; Nonglak et al., 2005; Ana et al., 2006; Il-Jin et al., 2007; Ling-Xiang et al., 2007; Zhengshan et al., 2008). The




Figure 4. RFLP analysis of *groEL* gene fragment of some pathogenic bacteria. Lanes 1 – 6 are RFLP analysis patterns. 1: *S. enteric*; 2: *S. typhi*; 3: *S. typhimurium*; 4: *S. flexneri*; 5: *S. sonnei*; 6: *S. boydii*; Lane M: DNA molecular marker

p	b	j	f	q
	c	k	g	
a	d	l	h	
	e	m	i	
p		o	n	q

Figure 5. The probe position of the oligonucleotide array. The letter in the grids is probe name as indicated in Table 2.

conserved genes of pathogenic micro-organisms are usually used as the target gene in the identification of pathogens, such as 16S rRNA gene, 23S rRNA gene, 16-23S rRNA space region. In these conserved genes, 16-23S rRNA space region is a short gene fragment with large extent of variations; 23S rRNA also has great variation in bacterial species, but its sequence is incomplete; 16S rRNA is most conservative among species and its sequences is the most complete which can be used for the classification of bacteria (Gelsen et al., 1994; Gurtler et al., 1996; Anthony et al., 2000; Hayden et al., 2001; Corinne et al., 2008; Ghebremedhin et al., 2008). Heat shock protein is a highly conserved protein, whose encoding gene *groEL* constitutes to be the most conserved component in evolution. The *groEL* gene

has been used as the target gene in the typing and identification of *Salmonella*, *C. jejuni* and *Staphylococcus* on account of its complete database (Goh et al., 1996; Satheesh et al., 2002; Karenlampi et al., 2004). Thus, in this study, we select the *groEL* gene as the target gene to investigate its significance in the identification of pathogenic bacteria.

It was found that the *groEL* gene appeared to be conserved and mutable, and therefore can be used as a general phylogenetic marker to construct the phylogenetic tree. In this way, the phylogenetic tree of various pathogenic bacteria can be drawn based on *groEL* gene using the bio- software and the phylogenetic tree constructed was found to be consistent with those based on 16S rRNA gene and 23S rRNA gene. Even though, the *groEL* gene analysis has several characteristics different from those of analysis of other genes. Compared with the 16S rRNA gene and 23S rRNA gene, the *groEL* sequences have higher divergence for species. The *groEL* gene is more heterogeneous than 16S rRNA gene and 23S rRNA gene and it is useful in intraspecies differentiation. So the *groEL* sequence analysis is useful not only in interspecies differentiation but also in intraspecies differentiation. The species were characterized by phylogenetic analysis based on *groEL* gene sequences revealed that species within a group are highly related. For example, *S. flexneri*, *S. sonnei* and *S. boydii* are highly related. Similarly, *S. enterica*, *S. typhimurium* and *S. typhi* are highly related. From this phylogenetic tree, it is apparent that the homology of bacteria belonging to the same family is higher than those of bacteria of different families, such as, the interrelationship of *Salmonella* and *Shigella* in the same family is closer than that of *Salmonella* with *Mycobacterium* in the different family. From the result of variance analysis on *groEL* gene, it is evident that its variation is rather high, especially among interspecies. There are 4 variant regions in *groEL* gene, in which the A region appears to be the most mutable one, while the D region is the least mutable one. According to the distribution of *groEL* gene, we have designed the PCR universal primers and by the use of a pair of primers thus synthesized, all the corresponding gene fragments in pathogenic bacteria could be amplified by PCR under the proper amplification conditions. Meanwhile, the detection probe for genus and species of bacteria can be also designed based on the variant region.

Subsequently, we can use PCR-RFLP or PCR-SSCP in the classification of pathogenic bacteria as well, and can also take advantage of special probes for spot hybridization detection. Above all, we can still design genus, species or type probes based on the variant region as well as construct the high throughput, high effective and parallel disposal microarray for bacterial identification from genus to species and type in the same membrane array. Our study showed that satisfactory hybridization results with good sensitivity and specificity were obtained with these tested bacteria. *S. boydii* gave weak signal




Figure 6. Membrane array hybridization assay results of some pathogenic bacteria. Lanes 1 - 10 are hybridization results of different bacterial species. 1: *S. flexneri*; 2: *S. boydii*; 3: *S. sonnei*; 4: *S. typhimurium*; 5: *S. typhi*; 6: *S. enterica*; 7: *S. aureus*; 8: *S. haemolyticus*; 9: *S. epidermidis*; 10: *Streptococcus pneumoniae*.

because of the low amount of oligonucleotide dotted on the membrane. Proper design of probes is the key factor for successful oligonucleotide array hybridization. Hybridization conditions and bacteria gene mutations often affect the sensitivity and specificity of the detection. In addition to probes, different labeling methods also influence the result of detection.

In conclusion, the *groEL* gene is useful for the identification and characterization of pathogens.

ACKNOWLEDGMENTS

This study was supported by Guangzhou Scientific and Technological Project (No.2007J1-C0201).

REFERENCES

- Ana CRM, Beatriz ECG, Marina BM (2006). Can the *flhC* PCR-Restriction Fragment Length Polymorphism Technique Replace Classic Serotyping Methods for Characterizing the H Antigen of Enterotoxigenic *Escherichia coli* Strains? *J. Clin. Microbiol.*, 44: 1453-1458.
- Anthony RM, Brown TJ, Freng GL (2000). Rapid diagnosis of bacteremia by universal amplification of 23S ribosomal DNA followed by hybridization to an oligonucleotide array. *J. Clin. Microbiol.*, 38: 781-788.
- Bangxing H, Lifang J, Yushan H, DanYun F, HuiYu G (2004). Application of oligonucleotide array technology for the rapid detection of pathogenic bacteria of foodborne infections. *J. Microbiol. Methods*, 58: 403-411.
- Carl RH, Sacha L, Karyn PR, Udo W, Tracy JE, Jay CDH, Michael JG, Arjan N (2008). A short-oligonucleotide microarray that allows improved detection of gastrointestinal tract microbial communities. *BMC Microbiol.*, 8: 195.
- Corinne M, Laëtitia LD, Marcelo G, Marylène K (2006). Molecular characterization of *Streptococcus suis* strains by 16S–23S intergenic spacer polymerase chain reaction and restriction fragment length polymorphism analysis. *Can. J. Vet. Res.*, 70: 94-104.
- Eric G, Christophe H, Chi ZL, Carol AG (2004). A chaperone network controls the heat shock response in *E. coli*. *Genes Dev.*, 18: 2812-2821.
- Franklin R, Cockerill III, Thomas FS (2004). Response of the Clinical Microbiology Laboratory to Emerging (New) and Reemerging Infectious Diseases. *J. Clin. Microbiol.*, 42: 2359–2365.
- Gelsen K, Loeffelhol M, Purohit A, Leong D (1994). PCR Primers and probes for the 16S rRNA Gene of Most species of Pathogenic Bacteria, Including Bacteria Found in cerebrospinal Fluid. *J. Clin. Microbiol.*, 32: 335-351.
- George K, Angela T, Stuart J, Jon B, Ed K, Jacques P, Eric HF, Paul S, Brad N, Renate J, vanden B, Haru K, Susan PS, Walter Z, Christopher W, Brandi L, Dale NG, Clifford M (2008). Comparison of Seven Techniques for Typing International Epidemic Strains of *Clostridium difficile*: Restriction Endonuclease Analysis, Pulsed-Field Gel Electrophoresis, PCR-Ribotyping, Multilocus Sequence Typing, Multilocus Variable-Number Tandem-Repeat Analysis, Amplified Fragment Length Polymorphism, and Surface Layer Protein A Gene Sequence Typing. *J. Clin. Microbiol.*, 46: 431-437.
- Gerard A, Robert JF, Kaeryn, Lewis NL, Devon LR, Ketan SS, Sparrow JM, Stefan VG (2004). Evaluation of a High-Throughput Repetitive-Sequence-Based PCR System for DNA Fingerprinting of *Mycobacterium tuberculosis* and *Mycobacterium avium* Complex Strains. *J. Clin. Microbiol.*, 42: 2685-2693.
- Ghebremedhin B, Layer F, König W, König B (2008). Genetic Classification and Distinguishing of *Staphylococcus* Species Based on Different Partial gap, 16S rRNA, hsp60, rpoB, sodA, and tuf Gene Sequences. *J. Clin. Microbiol.*, 46: 1019-1025.
- Giuseppe B, Vincenzina F, Danilo E, Maria A, Olimpia P, Francesco V (2008). *Lactobacillus* Strain Diversity Based on Partial hsp60 Gene Sequences and Design of PCR-Restriction Fragment Length Polymorphism Assays for Species Identification and Differentiation. *Appl. Environ. Microbiol.*, 74: 208-215.
- Goh SH (1996). *groEL* gene sequences as universal targets for microbial species identification: studies with coagulase-negative staphylococci. *J. Clin. Microbiol.*, 34: 818-823.
- Gurtler V, Stanisich VA (1996). New approaches to typing and identification of bacteria using the 16S-23S-rDNA-spacer region. *Microbiology*, 142: 3-16.
- Hayden RT, Kolbert CP, Hopkins MK, Persing DH (2001). Characterization of culture-derived spiral bacteria by 16S ribosomal RNA gene sequence analysis. *Diagn. Microbiol. Infect. Dis.*, 39: 55-59.
- Hyun-Hee K, Ji-Yeol S, Hak-Sun Y, Kim J, Hahn TW, Hahn YH, Chung

- DI(2002). Mitochondrial DNA Restriction Fragment Length Polymorphism (RFLP) and 18S Small-Subunit Ribosomal DNA PCR-RFLP Analyses of *Acanthamoeba* Isolated from Contact Lens Storage Cases of Residents in Southwestern Korea. *J. Clin. Microbiol.*, 40: 1199-1206.
- Il-Jin K, Hio CK, Sang GJ, Sun AA, Hyunju Y, Jaegahb P (2007). Development and Applications of a BRAF Oligonucleotide Microarray. *J. Mol. Diagn.*, 9: 55-63.
- Jung-Hee L, Hyo-Soon P, Won-Jong J (2003). Differentiation of *Rickettsiae* by groEL Gene Analysis. *J. Clin. Microbiol.*, 41: 2952-2960.
- Karenlampi RI, Tolvanen TP, Hanninen ML (2004). Phylogenetic Analysis and PCR-Restriction Fragment Length Polymorphism Identification of *Campylobacter* Species Based on Partial groEL Gene Sequence. *J. Clin. Microbiol.*, 42: 5731-5738.
- Karuna P, Yasuyuki N, Timothy DR, Artem C, Jeffrey K, Caixia S, Colleen CN, Robert CB (2003). Molecular Analysis of the Multiple GroEL Proteins of *Chlamydiae*. *J. Bacteriol.*, 185: 1958-1966.
- Kensuke S, Jun T, Toshio S, Kazuhiko N, Kazumichi T, Haruo W, Yoshifumi T, Shinji Y (2004). Development of a PCR-Restriction Fragment Length Polymorphism Assay for the Epidemiological Analysis of Shiga Toxin-Producing *Escherichia coli*. *J. Clin. Microbiol.*, 42: 5205-5213.
- Kwok AY, Su SC, Reynolds RP, Wilson JT, Coulthart M, Mutharia L, Chow AW (1999). Species identification and phylogenetic relationships based on partial groEL gene sequences within the genus *Staphylococcus*. *Int. J. Syst. Bact.*, 49: 1181-1192.
- Ling-Xiang Z, Zhi-Wei Z, Can W, Hua-Wei Y, Di J, Qiong Z, Keith M, Jing C (2007). Use of a DNA Microarray for Simultaneous Detection of Antibiotic Resistance Genes among Staphylococcal Clinical Isolates. *J. Clin. Microbiol.*, 45: 3514-3521.
- Nonglak Y, Colleen O, Michael FG, Chulabhorn Pr, Sumalee P (2005). M protein typing of Thai group A streptococcal isolates by PCR-Restriction fragment length polymorphism analysis. *BMC Microbiol.*, 5: 63.
- Rebecca S, Wong Y, Chow AW (2002). Identification of enteric Pathogenes by heat shock protein 60kDa (GroEL) gene sequences. *FEMS Microbiol. Lett.*, 206: 107-111.
- Rousseaux S, Olier M, Lemaitre JP, Guzzo J (2004). Use of PCR-Restriction Fragment Length Polymorphism of inlA for Rapid Screening of *Listeria monocytogenes* strains Deficient in the Ability to Invade Caco-2 Cells. *Appl. Environ. Microbiol.*, 70: 2180-2185.
- Satheesh N, Thong KL, Tikki P, Martin A (2002). Characterization of *Salmonella* Serovars by PCR Single-Strand Conformation Polymorphism Analysis. *J. Clin. Microbiol.*, 40: 2346-2351.
- Sensu L, Hyo-Soon P, Jung-Hee L, Seongeun K, Tae-Kyu P, Won-Jong J, Kyung-Hee P, Bum-Joon K, Yoon-Hoh K (2004). Evaluation of groEL Gene Analysis for Identification of *Borrelia burgdorferi*. *J. Clin. Microbiol.*, 42: 1270-1273.
- Yu-Hsiu C, Yung-Hui S, Hung-Chi L, Hwan-Wun L (2003). PCR Assay of the groEL Gene for Detection and Differentiation of *Bacillus cereus* Group Cells. *Appl. Environ. Microbiol.*, 69: 4502-4510.
- Zhengshan Z, Gerardo A, Francois JP, Ann H, Éric L, Johanne F, Guy B, Teodor V, Michel MD, Michel GB (2008). Plastic Polymers for Efficient DNA Microarray Hybridization: Application to Microbiological Diagnostics. *J. Clin. Microbiol.*, 46: 3752-3758.