

Full Length Research Paper

Effects of manure on morphological qualities in *ALOE VERA*

Bagheri G.Hatami and Ahmed Arsham and Arya Hesamedin

Department of Horticulture, Faculty of Agriculture, Urmia University, Urmia, West Azerbaijan Province, Iran

Accepted 8 April, 2013

This experiment was conducted at the Institute of National Genetic Engineering and Biotechnology in Iran to evaluate the effect of different amount of fertilizers on the leaf and plant characteristics as well as yield of *ALOE VERA*. There were 6 different treatments viz., T₁ = 100% soil (control), T₂ = 100 PK (50% P + 50% K), T₃ = 50% K + 50% soil, T₄ = 50% N + 50% soil, T₅ = 150% NPK (50% N + 50% P + 50% K, T₆ = 50% P + 50% soil. It was observed that the plant produced highest mature leaf length and number of tillers plant⁻¹ and maximum leaf weight, as well as weight of largest leaves with application of 50% P+ 50% K (T₂). Different plant characters such as mature leaf breadth and breadth of the largest leaf was also found to be highest with T₅ treatment over control (100% soil). The number of leaves was also significantly affected by different fertilizer treatments where maximum effect was noticed at early stages with T₄ (50% N). It was revealed that T₃ had a little effect on number of leaves of *A. VERA* over chemical fertilizer.

Key words: *Aloe vera*, fertilizer, morphological traits, leaf growth.

INTRODUCTION

Aloe genus (family Liliaceae) consists of at least 600 known species, many of which have been used as botanical medicines in many countries for centuries (Okamura et al., 1996). Species of *Aloe* which have been used as folk medicine includes: Curacao Aloe (*Aloe barbadensis* or *Aloe vera*), Cape Aloe (*Aloe ferox*), and Socotra Aloe (*Aloe perryi*). This plant species can be easily propagated from cuttings and is probably the most widely cultivated species of the genus in the world. It is widely used in world, not only as a folk remedy for gastrointestinal complaints, skin injuries and burns, but also as an ingredient in health foods and cosmetics (Capasso et al., 1998). Products containing *A. vera* are used for the treatment of minor cuts and burns and to heal wounds. They are also contained in a variety of cosmetics including skin creams, lotions and shampoos. Aloe gel, among other things, enhances immunity,

improves liver function, prevents asthma and has anti-inflammatory, anti-ulcerous, anti-diabetic and anti-hypertensive properties (Dagne et al., 2000). Also epidemiological data suggest that the intake of *A. vera* prevents human lung cancer (Sakai, 1989). The cultivation of *A. vera* has acquired great commercial importance for medicinal products and cosmetics processing but information are scarce about agronomic management of this crop.

The land of Iran is very fertile as well as seasons are variable and favorable for various medicinal plants to grow. *A. vera* is cultivated in many places in Iran country but not in wide range. Cultivation of *A. vera* is expanding day by day in the area as it provides quick and regular income to the farmers. Farmers are not using any recommended farming practices for *A. vera* cultivation which resulted poor yield. Fertility management in *A. vera* field may be one of the strategies for increasing of the yield of *A. vera* (Saha et al., 2005). As *A. vera* is a succulent plant, it is more responsive to nutrient. However, the excess doses of chemical nutrient as well as improper sources can show negative effect of quality.

*Corresponding author. E-mail: Baghari16@gmail.com

Table 1. Geographical origins of aloe genotypes.

Ecotype no	Region originated	Climate ^a	Latitude, N	Longitude, E	Altitude (m)
1-10	Boushehr	Warm temperate-humid	50 28 'N	27 59 'E	5
11-20	Borazjan	Warm-arid	29 15 'N	51 12 'E	94
21-30	Sar korreh	Warm-arid	28 53 'N	51 17 'E	43
31-40	Hormozgan	Warm temperate-humid	27 75 'N	56 05 'E	9
41-50	Bashagerd	Warm -arid	50 31 'N	29 34 'E	10

^a Yearly mean temperature in warm, temperate and cool climates are respectively 30 to 50 and 0 to 24.5°C. Yearly mean rainfalls are warm- arid and warm temperate-humid climates are respectively 100 to 300 mm, 300 to 400 mm.

Table 2. Botanical traits of Iranian Aloe genotypes.

Location	Heightof plant(cm)	Breadthof plant(cm)	No. ofleaves plant ⁻¹	Matureleaf length(cm)	Matureleaf breadth(cm)	Lengthofthe largestleaf(cm)	Breadthofthe largestleaf(cm)	Singlemature leafweight(g)	Weight of the largest leaf (g)	No. of tillers plant ⁻¹
Borazjan	61.20 ^{ab}	20.80 ^a	25.20 ^{ab}	44.40 ^a	9.60 ^a	5.50 ^b	1.72 ^b	1.77 ^a	2.15 ^b	44.60 ^a
Sarkorreh	51.50 ^b	10.80 ^b	17.60 ^{abb}	40.70 ^a	7.83 ^{ab}	45.60 ^b	2.12 ^a	1.54 ^{ab}	1.74 ^c	42.80 ^a
Boushher	55 ^b	11.20 ^b	29.50 ^a	45.20 ^a	7.95 ^{ab}	49.10 ^b	2.26 ^a	1.65 ^{ab}	1.85 ^{bc}	44.70 ^a
Hormozgan	55.80 ^b	23.9 ^a	25.30 ^{ab}	40.80 ^a	8.80 ^{ab}	46.30 ^b	2.15 ^a	1.47 ^b	1.83 ^{bc}	42.50 ^a
Bashagerd	69.70 ^a	20.40 ^a	14.40 ^b	40.40 ^a	7.10 ^b	67.30 ^a	2.07 ^a	1.77 ^a	2.95 ^a	44 ^a
Total	58.64	17.43	22.40	42.30	8.25	51.76	2.06	1.64	2.10	43.72

Optimum chemical fertilizers are more effective in *A. vera* growth and yield. In addition chemical fertilizers enhance a good leaf quality. So, it may be necessary to find out a suitable recommendation for fertilization in *A. vera* farming. The present study was carried out to determine the effect of fertilization levels on the growth and yield of *A. vera*.

MATERIALS AND METHODS

A total of 50 Aloe ecotypes were collected from different areas. This sampling was done according to the information of local agricultural extension offices and producers from all over Iran. Geographical origins of the 50

Aloe ecotypes are listed in Table 1. This experiment was conducted at the Institute of National Genetic Engineering and Biotechnology in Iran, during 2010. The climate of this area is subtropical. The soil of the experimental site was clay loam with pH of 6.5. The experiment was laid out by Randomized Completely Block Design (RCBD) with 4 replications comprising six different treatments viz. T₁ = 100% soil (control), T₂ = 100% PK (50% P +50% K), T₃ = 50% K + 50% soil, T₄ = 50% N + 50% soil, T₅ = 150% NPK (50% N, 50% P, 50% K), T₆ = 50% P + 50% soil. Plants of collected from 5 regions (Borazjan, Sarkorreh, Boushehr, Hormozgan and Bashagerd) were planted in the greenhouse and determined morphological traits (Table 2). Distance between plants was 20 cm. Chemical fertilizer was used in this experiment as per treatment (Hasanuzzaman et al., 2008). The field was irrigated whenever necessary. Data were recorded every 15 days interval starting from 15 days after planting (DAP) for

measuring leaf length and breadth. Final data were recorded at harvest to measure plant characters. Flexible tap and scale were used to measure leaf. Weighing was done by digital balance (Kaifeng Group Co., Ltd., China). The data were analyzed following analysis of variance (ANOVA) technique and mean separations were adjusted by the multiple comparison test using the statistical computer program MSTAT-C v.1.2 (MSTAT-C, 2000). Means were compared by using LSD test at 5% level of significance.

RESULTS

Different combination of fertilization significantly affected the leaf characteristics of *A. vera* in this experiment (Table 3). The highest number of leaf

Table 3. Duncan s multiple range tests analysis for mean comparison of evaluated traits *Aloe vera* plant at harvest 60 (DAP).

Treatment	Mean amount of fertilizer	Mean low vigor sucker	Mean med vigor sucker	Mean high vigor sucker	Mean total No. of sucker	MeanNo. of leaves plant ⁻¹	Mean mature leaf length (cm)	Mean mature leaf breadth (cm)	Mean length of the largest leaf (cm)	Mean breadth of the largest leaf (cm)	Mean single mature leaf weight (g)	Mean weight of the largest leaf (g)	Mean No. of tillers plant ⁻¹
1 (0% control)	7.47 ^{ab}	2.73 ^a	3.15 ^a	1.63 ^a	5.31 ^{ab}	13.31 ^a	25 ^b	6.24 ^a	30.94 ^a	7.05 ^a	7463.2 ^a	8846.8 ^a	43.9 ^{ab}
2 (100% pk)	8.2 ^a	3/20 ^a	4 ^a	1 ^a	4 ^b	13 ^a	33 ^a	5.88 ^a	31.60 ^a	6.60 ^a	7960 ^a	9020 ^a	45.80 ^a
3 (50% k)	7.22 ^b	3 ^a	2.66 ^a	1.55 ^a	4.66 ^b	12.55 ^a	29.67 ^{ab}	6.47 ^a	34.77 ^a	6.50 ^a	7955.6 ^a	9000 ^a	42.7 ^{ab}
4 (50% N)	7.44 ^{ab}	3.77 ^a	2.44 ^a	1.22 ^a	7.22 ^a	14.22 ^a	30.33 ^{ab}	6.32 ^a	34.55 ^a	7.02 ^a	7100 ^a	8344.4 ^a	41.55 ^b
5 (150% NPK)	7.44 ^{ab}	3.55 ^a	3.33 ^a	0.55 ^a	4.44 ^b	13.11 ^a	27.44 ^{ab}	6.64 ^a	36.11 ^a	7.16 ^a	7344.4 ^a	8977.8 ^a	44 ^{ab}
6 (50% P)	7.33 ^b	2 ^a	4 ^a	1.33 ^a	5.88 ^{ab}	14.11 ^a	27.88 ^{ab}	6.33 ^a	37.22 ^a	6.61 ^a	7577.8 ^a	9011.1 ^a	44.8 ^{ab}

Table 4. Variance analysis for evaluated traits *aloe vera*.

	df	Meansquare													
		Amount of fertilizer	Low vigor sucker	Med vigor sucker	High vigor sucker	Total No. of sucker	No. of leaves plant ⁻¹	Mature leaf length (cm)	Mature leaf breadth (cm)	Length of the largest leaf (cm)	Breadth of the largest leaf (cm)	Single Mature leaf weight (g)	Weight of the largest leaf (g)	No. of tillers plant ⁻¹	
Treatment	6	0.67 ^{ns}	3.74 ^{ns}	3.37 ^{ns}	1.63 [*]	1.75 [*]	10.9 ^{ns}	3.74 ^{ns}	54.29 ^{ns}	0.44 [*]	31.07 [*]	0.76 [*]	926224.5 [*]	605484 ^{ns}	17.2 ^{ns}
Error	54	0.54	2.96	2.53	1.93	2	4.42	3.95	38.35	0.87	35.67	1.23	1060670.5	513094	12.71
CV%	-	9.9	57.74	49.72	106.96	36.60	39.30	14.84	21.88	14.80	16.71	16.15	13.68	8.09	8.15

per plant was observed from T₄ = 50% N (50% N + 50% soil) followed by T₆ = 50% P (50% P + 50% soil). Concerning total leaf fresh weight per plant, the data in the same table indicate that all treatments increased significantly leaf fresh weight as compared to control plant.

Using 100% PK fertilizer was the most effective in increasing leaf fresh weight per plant (7960 g). Different fertilization treatment also significantly affected the plant characteristics of *A. vera* in this study (Table 3). The highest weight of plant at harvest (7920 g) was observed in the treatment where 100% PK fertilizer was applied (T₂) which

was statistically superior to other treatments. The lowest plant weight (7100 g) was observed in (T₄) 50% N (Table 4). In case of leaf type no. 1 (from base), continuous and highest growth rate of leaf was observed in the treatment where 100% PK fertilizer was applied, But length of the largest leaf was observed in T₆ (50% P). The leaf length was also highest in this case (Figure 1). This trend was also found in the treatments where comparatively more fertilizer was applied. The growth rate in length decreased with the decrease in fertilizer percentage (Figure 1). The lowest growth rate was observed in control treatment where no

fertilizer was applied. The highest of number of suckers was observed in 50% N. There were negative correlation between high-vigor, med vigor and low vigor of suckers with all treatment. Growth rate of mature leaf breadth has been presented in Figure 2. In this case the highest growth rate was observed in the treatments where higher amount of fertilizer was used. The lowest growth rate was observed in T₂ (100% PK) and control treatment. Higher leaf breadth and breadth of the largest leaf trend was also observed with the treatment T₅ where 150% NPK fertilizer was applied.




Figure 1. Leaf length of *Aloe vera* leaf type no.1 at different days after planting.




Figure 2. Leaf breadth of *Aloe Vera* leaf type no. 2 at different days after planting.

DISCUSSION

In this experiment application of 50% N as recommended doses did not showed superior results than fertilizer application which was supported by Saha et al. (2005) and Nobel et al. (1991). This trend of increased production due to increased application of fertilizer were also observed in case of single leaf weight, leaf length, number of tillers and weight of the largest leaf. The application of nutrient matter were increased the cell division and elongation without hampering the nutrient

uptake process which provided the better results due to better nutrition. Guerrero et al. (2001) found that nutrient matter addition is a suitable technique for accelerating the natural recovery process of burned soils. An increased trend of plant weight was observed with the increase of fertilizer amount. It was due to the beneficial effect of nutrient matter in soil properties and plant growth (Uyanoz et al., 2002; Dexter, 1988; Tisdall and Oades, 1982). More or less similar trend was observed in case of total plant weight along with tillers. Number of tillers per plant was highest with the treatment T₂ (50% P + 50% K)

which were followed by T₅ and T₆. These results were supported by Hernández-Cruz et al. (2002).

The lowest growth rate was observed in case of control treatment where no fertilizer was applied. The growth rate of the leaves was highest at early stages of growth which declined gradually. The growth of control treatment was lower but consistent up to 60 DAP. These findings were also in agreement with the results of Van Schaik et al. (1997) and Chatterjee et al. (1979). The highest length of leaves was observed in case of the treatment where 100% PK, but highest length of the largest leaf was observed in 50% P fertilizer was applied. This dose of fertilizer improved the *A. vera* plant growth by providing the essential nutrient which results the maximum cell growth and turgidity which influenced the leaf growth. Pichgram (1987) also observed similar results in *A. vera*. The highest percentage of increase in mature leaf breadth was observed in the early stage of growth irrespective of treatments. Some regular increase in growth rate of mature leaf breadth was observed in control treatment at later stage. Van Schaik et al. (1997) observed similar results. In case of leaf type no. 2 growth rate of leaf breadth was higher with the treatments where higher percentage of fertilizer was used (Figure 2). The percentage of increased rate was highest in the early 15 days in all the treatments. This increase was gradually decreased in the later stages. The decreasing rate was highest in T2 and control treatment and treatments with lower doses of fertilizer. It was due to the failure of *Aloe* plant to produce the cell expanse. This result was supported by Saha et al. (2005) and Bates (1971).

REFERENCES

- Capasso F, Borrelli F, Capasso R (1998). *Aloe* and its therapeutic use. *Phytother Res.*, 12: 124-127.
- Chatterjee BN, Singh KI, Pal A, Maiti S (1979). Organic Fertilizer as a Substitute for Chemical Fertilizers for High-Yielding Rice Varieties. *Indian J. Agric. Sci.*, 49: 188-192.
- Dagne E, Bisrat D, Viljoen A, Van-Wyk BE (2000). Chemistry of *Aloe* species. *Curr. Org. Chem.*, 4: 1055-1078.
- Dexter AR (1988). Advances in characterization of soil structure. *Soil Tillage Res.*, 11: 199-238.
- Guerrero C, Gomez I, Moral R, Mataix-Solera J, Mataix-Beneyto J, Hernandez T (2001). Reclamation of a burned forest soil with municipal waste compost: macronutrient dynamics and improved vegetation cover recovery. *Bioresour. Technol.*, 73(3): 221-227.
- Hernández-Cruz LR R, Rodríguez DJ, Angulo-Sánchez JL (2002). *Aloe vera* Response to Plastic Mulch and Nitrogen. In: Janick J, Whipkey A (eds) *Trends in new crops and new uses*: ASHS Press, Alexandria, VA.
- MSTAT-C (1990). A microcomputer program for the design management and analysis of agronomic research experiments. MSTAT, Michigan States University, East Lansing.
- Nobel PS, Quero E, Linares H (1991). Root versus shoot biomass: Responses to water, nitrogen and phosphorus interaction application for *Agave lechuguilla*. *Hort Abst.*, 61: 513.
- Okamura N, Asal M, Hine N, Yago A (1996). Highperformance liquid chromatographic determination of phenolic compounds in *Aloe* species. *J. Chromatogr. A.*, 746: 225-231.
- Pichgram N (1987). Investigation on plant spacing and fertilizer effect on growth of *Aloe (Aloe barbadensis Mill.)*. Ph.D. Thesis, University of Kasetsart, Bangkok, Thailand.
- Saha R, Palit S, Ghosh BC, Mitra BN (2005). Performance of *Aloe vera* as influenced by organic and inorganic sources of fertilizer supplied through fertigation. *Acta Hort.*, 676: 171-175.
- Sakai R (1989). Epidemiologic survey on lung cancer with respect to cigarette smoking and plant diet. *Jpn. J. Cancer Res.*, 80: 513-520.
- Tisdall JM, Oades JM (1982). Organic matter and water-stable aggregates in soils. *J. Soil Sci.*, 33: 141-163.
- Uyanoz R, Cetin U, Zengin M, Gur K (2002). Effect of different organic wastes on nitrogen mineralization and organic carbon contents of soil: International Conference at Canakkale, Turkey. *Sustainable Land Use and Management International*, pp. 223-228.
- Van Schaik AH, Struik PC, Damian TG (1997). Effects of irrigation and N on the vegetative growth of *Aloe barbadensis Mill.* in Aruba. *Tropical-Agric.*, 74: 104-109.