

Full Length Research Paper

Bone tumors at Addis Ababa University, Ethiopia: Agreement between radiological and histopathological diagnoses, a -5-year analysis at Black-Lion Teaching Hospital

Bayush E. Negash¹, Daniel Admasie¹, Biruk L. Wamisho^{2*} and Mihiret W. Tinsay³

¹Radiology, Addis Ababa University, Medical Faculty (AAU-MF), Ethiopia.

²Orthopedic Surgery, AAU-MF, Orthopedic Department, Ethiopia.

³Pathology, Addis Ababa University, Ethiopia.

Accepted 17 March, 2019

The study aims at looking into clinical, radiological and histopathological agreement in diagnosing bone tumors and elaborates the spectrum of bone tumors referred to Black-Lion Hospital, BLH. All bone tumor patients, who presented to all the three departments at BLH between the study periods of December 2003 – March 2008 were recruited. The radiological and histopathological diagnoses made were categorized separately using WHO classification of bone tumors. Strength of agreement was measured using Cohen's Kappa test. From the total of 216 bone tumor patients presented and biopsied in the five year period, complete information could not be gathered only for 11 (5%) of the patients. Hence there were 111(51.3%) males and 94(48.7%) females with histopathological tissue diagnosis of the bone tumor. Male to female ratio was nearly 1. Commonest age range generally affected was 15-29 (7-55 years). Of the tumors histopathologically diagnosed, 36%(74/205) of the tumors were malignant. The commonest primary malignant tumor from bone biopsy was osteosarcoma (52/182, 28.5%). It was also the single most common clinical, radiological and histological diagnosis made. Exostosis was second. Considering all bone tumors together, radiological diagnosis was confirmed by similar histological diagnosis in 172 out 205 cases (84%) and the corresponding Cohen's kappa value (0.82) showed excellent agreement between radiological and histological diagnoses of all bone tumors. In set-ups where there is limited or no histopathological service, joint clinical and radiological decision could lead to a higher degree of accuracy in diagnosis of bone tumors.

Key words: Bone tumors, diagnostic agreement, radiological, orthopedic, histopathology.

INTRODUCTION

Bone tumor is an abnormal growth arising from bone. It could be benign or malignant but most commonly the term is used for primary tumors; it is less applied to secondary or metastatic tumors found in bone. Bone tumors may be classified as "primary tumors" which originate in the bone, and "secondary tumors" which originate elsewhere and involve the bone secondarily. Primary tumors of bone can be divided into benign tumors and malignant

tumors. Common benign bone tumors may be neoplastic, developmental, traumatic, infectious, or inflammatory in etiology. Primary bone tumor accounts for 0.2% of all tumors in humans, whereas involvement of skeletal tissue by metastatic disease is much more common (Schulthess et al., 2005). Bone tumors are mostly mesenchymal in origin, though, for example Ewing's sarcoma is thought to have neuroectodermal precursor cells. Primary bone tumors are uncommon malignancy, but they are important causes of cancer morbidity and mortality, especially among young people. Although relatively rare in childhood, primary bone tumor represents the sixth most common neoplasm in children, while in adolescents and

*Corresponding author. E-mail: lbiruklw@yahoo.com. Tel: (+00251) 911 232 507.

young adults; they are the third most frequent, exceeded only by leukemia and lymphoma (Schulthess et al., 2005). Examples of benign bone tumors include osteoma, osteochondroma, aneurysmal bone cyst, and fibrous dysplasia. Malignant primary bone tumors include osteosarcoma, chondrosarcoma, Ewing's sarcoma, and other sarcoma types. Multiple myeloma is a hematologic malignant tumor which also frequently presents as one or more bone tumors.

Secondary bone tumors include metastatic tumors which have spread from other organs, such as the breast, lung, and prostate. Metastatic tumors more frequently involve the axial skeleton than the appendicular skeleton. Tumors which originate in the soft tissues may secondarily involve bones through direct invasion. A review of literature reveals that malignant bone lesions were demonstrated in bones of ancient Egyptian mummies (Hamada et al. 1972). For a long time, osteosarcoma has been associated with a low social-economic stratum (Dodge et al., 1973). This is a major problem in the developing countries where limb-sparing surgery is not yet possible. Amputation is therefore often the mode of treatment making the patient disabled, non productive and a burden to the family (Ssebagala, 1997; Ahuka et al., 1995; Kiwanuka et al., 1983).

Literature on bone tumors in Africa is scarce. In a study carried out in Uganda between 1964 and 1968, osteosarcoma was found to be the commonest primary malignant bone tumour with a peak age of 10 - 19 years. The commonest site was the lower femur (Dodge et al., 1973). This is similar to what has been reported elsewhere (Parkin et al., 1997). It has been suggested that major histocompatibility complex linked genes may determine susceptibility to some types of tumors (Turo et al., 1990). Other epidemiological factors mentioned to be linked are mechanical trauma, ionizing radiation and chronic osteomyelitis (Fornasier et al., 1998; Bascoulergue et al., 1996; Hasbini et al., 1998; Adil et al., 1996; Maeda et al., 1996).

Radiological diagnosis of any bone tumor takes into account the clinical history, site of lesion, borders of the lesion, type of matrix, type of bone destruction, type of periosteal reaction, nature and extent of soft tissue involvement and number of lesions. A systematic approach to clinical history, radiographic evaluation and histopathology is necessary for accurate diagnosis all three are important. As many bone lesions overlap, an experienced clinician systematically integrates the radiological and histopathological results to plan and offer the best possible management to the patient.

OBJECTIVES

This study is the first of its kind in the East African region to address 32 types of bone tumors and aims at:

1. Assessing the level of agreement between radiological

diagnoses of bone tumors at AAU-MF and

2. Profiling the spectrum of bone tumors presenting to BLH.

PATIENTS AND METHODS

All bone tumor patients, who presented to all the three departments at BLH between the study periods of December, 2003 – March, 2008 were recruited for the study. Histopathological and radiological diagnoses of the tumors were identified and their clinical profile looked in detail. All files related to the patients were reviewed. There was no complete information only for 11 patients and were not considered in this study. Hematologic tumors including malignant myeloma were also excluded from the study, as they are mainly diagnosed by bone marrow cytology and not bone tissue biopsy. All patients had a clinical examination, plain radiographs and biopsies of the affected part of the extremity. Radiographs of few selected difficult cases were usually discussed at the weekly joint orthopedic and radiologic sessions. The radiological and histopathological diagnoses made were categorized separately using WHO classification of bone tumors. The tumors were then classified according to the recent WHO classification of bone tumors (Dorfman et al., 2002; Schajowicz et al., 1993). Locations were divided into nine groups namely: bones of the limbs, bones of the limbs and spine, clavicle, cranio-facial bones, pelvic bones, ribs, scapula, spine, and sternum. The collected information was then analyzed using SPSS (statistical package for social sciences) and Microsoft Excel. Cohen's Kappa test to determine the level of radiological and histopathological diagnostic agreement was calculated online (kappa).

RESULTS

After exclusions by criteria, a total of 205 bone tumor patients were enrolled. By far the majority, 182 (182/205, 89%) of the biopsied bone lesions were neoplastic whereas 23 (23/205, 11%) of the lesions were non neoplastic conditions. Primary tumors accounted for 94.5% (172/182) of the neoplastic lesions. Metastasis was diagnosed in 10 (10/182, 5.5%) of the neoplasms. Benign lesions accounted for 57% (74/172) and malignant lesions for 43% (74/172) of the primary neoplasms. The histological diagnosis shows over one third (74/205, 36.1%) of the tumors are malignant: 22%(45/205) osteosarcoma, 5.36 % (11/205) Ewing's sarcoma, 0.5% leiomyosarcoma, 2.0% lymphoma, 1% malignant fibrous histiocytoma and 0.5% spindle cell neoplasm. The age ranged from 4 months to 65 years with median age 19 years. The majority of bone lesion occurred below 30 years of age (n=158, 77.1%); among these 55.1% (42.4% of the total) were in the age range of 10 to 19 years. 111 patients were male making male to female ratio 1.08:1. Bone tumor in children below the age of 15 occurred in 52 cases out of which the highest is osteosarcoma (n=18, 34.6 %), followed by exostosis (n=10, 19.2 %), Ewing's sarcoma (n=7, 13.5 %).

Table 1 shows the categorical distribution of radiological and histopathological diagnoses of all bone tumors in each sex. The table clearly shows that the distribution of bone lesion in each sex is varying from one type to other , for example; the male to female ratio is 1:1 in the case

Table 1. Categorical distribution of Radiological and Histopathological diagnoses of all bone tumors in each sex from December, 2003 – March, 2008, in Black-Lion Hospital, Addis Ababa University, Ethiopia.

Diagnoses	Males		Females		Total No. of Patients		Percent	
	(Rad.)	(Histopath.)	(Rad.)	(Histopath.)	(Rad.)	(Histopath.)	(Rad.)	(Histopath.)
ABC	1	1	3	3	4	4	1.95	1.95
Ameloblastoma	14	12	7	5	21	17	10.22	8.27
Bone Islands	0	0	1	0	1	0	0.49	0
Bony dysplasia	0	0	1	0	1	0	0.49	0
Brodie Abscess	0	0	1	0	1	0	0.49	0
Cementifying fibroma	0	0	0	2	0	2	0	0.98
Chondromyxoid fibroma	1	2	1	1	2	3	0.98	1.46
Dentigerous cyst	1	1	0	0	1	1	0.49	0.49
Enchondroma	2	2	2	3	4	5	1.95	2.44
Eosinophilic granuloma	0	0	2	0	2	0	0.98	0
Ewing's sarcoma	10	8	2	3	12	11	5.85	5.37
Exostosis	26	27	17	18	43	45	20.98	21.95
Fibrous dysplasia	3	8	5	6	8	14	3.9	6.83
GCT	13	14	8	8	21	22	10.24	10.73
Haemangioma	1	1	3	4	4	5	1.95	2.44
Leiomyosarcoma	0	1	0	0	0	1	0	0.49
Lymphoma	1	1	5	3	6	4	2.93	1.95
Malig fibrous histocyt.	0	1	0	1	0	2	0	0.98
Maxillary Sarcoma	1	0	0	0	1	0	0.49	0
Metastasis	3	3	8	7	11	10	5.37	4.88
Myositis ossificans	1	0	0	0	1	0	0.49	0
Non-ossifying fibroma	1	1	0	0	1	1	0.49	0.49
Normal report	1	0	0	0	1	0	0.49	0
Ossifying fibroma	0	0	1	2	1	2	0.49	0.98
Osteoblastoma	0	1	0	1	0	2	0	0.98
Osteoid osteoma	1	1	0	0	1	1	0.49	0.49
Osteoma	1	1	2	2	3	3	1.46	1.46
Osteosarcoma	27	23	23	22	50	45	24.39	21.95
Pathological fracture	1	0	0	0	1	0	0.49	0
Pleomorphic adenoma	0	1	0	0	0	1	0	0.49
Simple one Cyst	1	1	2	2	3	3	1.46	1.46
Spindle Cell Neoplasm	0	0	0	1	0	1	0	0.49
Total	111	111	94	94	205	205	100	100

of osteosarcoma, 3:2 in exostosis, 2:1 in ameloblastoma, 2:1 in, and 8:3 in Ewing's sarcoma.

The three most common bone lesions histopathologically proven were: osteosarcoma and exostosis equally first (n=45, 21.95 % each), giant cell tumor, GCT (n=22, 10.73%) and ameloblastoma (17, 8.29%), respectively. Radiologically, the commonest tumor singled out is osteosarcoma (24.5%, 50/205). The distribution of bone tumor groups is indicated in Table 2. Table 2 shows the distribution of bone tumor groups based on histopathological matrix. As clearly seen, the most common group of tumors were cartilaginous in origin (n=53, 25.9%) followed by osteogenic tumors (n=51, 24.9%). Table 3 shows the site/location of the tumor based on WHO classification. The majority of specimen were from bones of

the limbs (n=135, 65.8%), followed by craniofacial bones (n=41, 20.0%), pelvic bones (n=10, 4.9%), ribs (n=6, 2.9%), and scapulae (n=5, 2.4%).

Considering all bone tumors together, the study indicated that radiological diagnosis was confirmed by histological diagnosis in 172 out 205 cases (84%) and the corresponding Cohen's kappa value (0.82) calculated online, showed excellent agreement between radiological and histological diagnoses of all bone tumors (Cohen et al., 1990). Only one patient was reported radiologically normal and histopathologically diagnosed to have osteoblastoma. There was no patient reported normal after biopsy. The main diagnostic disagreement was noticed in diagnosing osteosarcoma. Radiological and histopathological diagnoses were not in agreement in eight (15.7%,

Table 2. Distribution of bone tumor groups based on histopathological matrix in December, 2003 – March, 2008 at Black-Lion Hospital. Addis Ababa University, Ethiopia.

Tumor group	Histological diagnosis	tumor (% in group)	Total	
			n	%
Cartilaginous tumors			53	25.85
	Exostosis	84.9	45	21.95
	Enchondroma	19.43	3	2.44
	Chondromyxoid fibroma	5.66		1.46
Osteogenic tumors			51	24.88
	Osteoma osteoid	5.88	3	1.46
	Osteoma	1.96	1	0.49
	Osteoblastoma	3.92	2	0.98
	Osteosarcoma	88.24		21.95
Fibrogenic tumors			4	1.95
	Cementifying fibroma malig	50.00	2	0.98
	Fibrous histocyt.	50.00	2	0.98
Ewing's sarcoma	Ewing's Sarcoma		11	5.37
Hematologic neoplasm	Lymphoma		4	1.95
GCT	Gct		22	10.73
Vascular tumor	Haemangioma		5	2.44
Miscellaneous tumors			32	15.61
	Ameloblastoma	53.13	17	8.29
	Metastasis	31.25	10	4.88
	Leiomyosarcoma	3.13		0.49
	Spindle Cell Neoplasm	3.13	1	0.49
	Non-Ossifying Fibroma	3.13	1	0.49
	Ossifying Fibroma	6.25	2	0.98
Miscellaneous lesions			23	11.22
	ABC	17.39	4	1.95
	Dentigerous Cyst	4.35	1	0.49
	Simple Bone Cyst	13.04	3	1.46
	Pleomorphic Adenoma	4.35	1	0.49
	Fibrous Dysplasia	60.87	14	6.83
Total			205	

8/51) of the osteosarcoma patients. While the lesion was finally histopathologically proven to be benign, radiological diagnosis of malignant bone tumor was made only in six ((6/51, 11.8%) of the cases. These are shown on Table 4.

DISCUSSION

This comparative study of bone tumors as a spectrum of lesions is the first of its kind from Ethiopia, if not from Africa. We have noted that the documentation of bone tumor patients at Black-Lion Hospital is excellent, only 5% (11/216) files were lost. This could be due to the fact that the analysis focused only on the last five years, which are very recent. We also noted that pathology department has reported spectrum of 32 different types of histopathological bone lesions. This shows the departments long

years of experience in detecting different bone lesions and it also indicates the presence of many types of bone tumors in a referral hospital.

In general terms, the age, sex, site and distribution of the tumors is comparable to most reports from elsewhere (Jongkolnee et al., 1990) we found the same groups of common tumors like osteosarcoma, osteochondroma, chondrosarcoma and giant cell tumors. Hematologic malignancies, like multiple myeloma, seem to be lower in our study. This is due to the fact that these tumors are diagnosed based on bone=marrow smear and not bone by biopsy. Excluding ameloblastoma of the jaw, 39.7% (n=60) of the primary non-hematologic bone tumors were malignant-a lower figure when compared to 63% in other series (Unni et al., 1996). As the most common primary non-hematologic malignant tumors of the bone, the frequency of osteosarcoma in the present study is relatively

Table 3. WHO distribution of site of bone tumor in December, 2003 – March, 2008 at Black-Lion Hospital. Addis Ababa University, Ethiopia.

Histopathological Diagnosis	WHO Site of bone tumor									
	BL	BL +S	C	CFB	PB	Rib	Scapula	Spine	Sternum	No
ABC	2	-	-	-	2	-	-	-	-	4
Ameloblastoma	-	-	-	17	-	-	-	-	-	17
Cementifying fibroma	-	-	-	2	-	-	-	-	-	2
Chondromyxoid fibroma	2	-	-	-	1	-	-	-	-	3
Dentigerous cyst	-	-	-	1	-	-	-	-	-	1
Enchondroma	2	-	1	-	-	2	-	-	-	5
Ewing's sarcoma	9	-	1	-	-	-	1	-	-	11
Exostosis	35	-	-	1	2	3	4	-	-	45
Fibrous dysplasia	4	-	-	9	-	1	-	-	-	14
Gct	21	-	-	-	1	-	-	-	-	22
Haemangioma	3	-	1	1	-	-	-	-	-	5
Leiomyosarcoma	1	-	-	-	-	-	-	-	-	1
Malig fibrous histo	2	-	-	-	-	-	-	-	-	2
Metastasis	5	1	-	-	1	-	-	2	1	10
Lymphoma	-	-	-	3	-	-	-	-	1	4
Non-Ossifying fibrous	1	-	-	-	-	-	-	-	-	1
Ossifying fibroma	-	-	-	2	-	-	-	-	-	2
Osteoblastoma	2	-	-	-	-	-	-	-	-	2
Osteoid osteoma	1	-	-	-	-	-	-	-	-	1
Osteoma	-	-	-	3	-	-	-	-	-	3
Osteosarcoma	42	-	-	-	3	-	-	-	-	45
Pleomorphic adenoma	-	-	-	1	-	-	-	-	-	1
Simple bone cyst	2	-	-	1	-	-	-	-	-	3
Spindle cell neoplasm	1	-	-	-	-	-	-	-	-	1
Total	135	1	3	41	10	6	5	2	2	205

Key: BL, Bone of the Limb; S, Spine; C, Clavicle; CFB, Craniofacial bone; PB, Pelvic bone

Table 4. Disagreements between radiological and histopathological diagnosis of osteosarcoma of the limbs at BLH, Dec. 2003-Mar. 2008.

Radiological diagnosis	Pathological diagnosis	Frequency
Osteosarcoma	*MFH	2
Osteosarcoma	#GCT	2
Lymphoma	Osteosarcoma	1
Osteosarcoma	Lieomyosarcoma	1
Osteosarcoma	Ossifying fibroma	1
Osteosarcoma	Fibrous dysplasia	1

*MFH, Malignant fibrous histiocytoma; #GCT, Giant cell tumor.

higher (n=45, 75%) compared to those studies conducted in Chiang Mai University (67.4%) (Jongkolnee et al., 2006) ; this could be explained by the fact that most patients with malignancy are referred to this hospital as it is the country's highest referral institution. The most common primary malignant bone tumor is osteosarcoma and its frequency is similar to the frequency in other study (Dorfman et al., 2002). The second most common malig-

nant tumor of the bone seen in children and adolescent was Ewing's sarcoma (Rosen et al., 1997). Osteosarcoma typically is a long bone tumor and rarely affects flat bones of the axial or appendicular skeleton. The cases examined in this study show 42 out of 45 cases of osteosarcoma are in long bone and 3 out of 45 in the pelvic bones. This observation agrees very well with the study conducted at La Timone adults University Hospital for a

16 years period (Duffaud et al., 2000). The present study showed that exostosis is the most common benign bone tumor accounted for 49.5% of all benign bone tumors. This is higher when compared with the study done in Chiang Mai University (23.58%) and we noticed low incidence of GCT (24.2%) compared to Chiang Mai University (34.9%). This may reflect the fact that most of these tumors are handled at lower level institutions as they are clinically benign; hence not referred to our hospital.

Specific types of tumors targeted certain age groups and anatomic sites. As a group, bone neoplasms affect all ages and arise in virtually every bone but most bone tumors presenting during the first decades of life have a propensity to originate in the long bones of the extremities. In our study, approximately 77.1% bone tumors were younger than 30 years and bone of the limb were the major site affected 65.8%. The predilection bone in specific tumor types correlates well with most of the results of previous reports (Dorfman et al., 2002; Rosen et al., 1997).

Cohen's (1960) kappa statistic (K) has long been used to quantify the level of agreement between two raters in placing persons, items, or other elements into two or more categories (Gwinn et al., 1986). Hence this test – statistic can be used to measure the level/strength of agreement between different raters (Orthopedists, radiologists and pathologists) in placing/diagnosing bone tumors into the different WHO categories. Kappa values are easily calculated online. Our corrected Cohen's kappa value of 82% (0.82) is an excellent level of agreement ($k > 0.75$) between radiological and histological diagnoses of all bone tumors. But as still some histologically benign bone tumors which are easily treated by simple surgery were diagnosed as radiologically malignant lesions which usually demands radical surgery, radiotherapy and chemotherapy; histopathological confirmation must always accompany bone tumor diagnosis whenever possible. Therefore bone tumor diagnosis should be made by joint consideration of clinical presentation, radiological appearance and histopathological results whenever possible.

Conclusion

Patients with a wide spectrum of bone tumors presented to Black Lion Hospital. There is an excellent/high level of agreement between radiological and histopathological diagnoses of bone tumors at Black Lion Hospital. In setups where there is limited or no histopathological service, joint clinical and radiological decisions could lead to a higher degree of accuracy in diagnosis of bone tumors.

Recommendations

Based on the findings of this study, we recommend the following:

i) As exact diagnosis of bone tumors is at times difficult, joint approach integrating clinical, radiological and histo-

pathological findings is recommended to increase accuracy. This is an excellent learning point to be adopted from BLH, AAU-MF.

ii) Where there is no or limited histopathological service, the surgeon could decide on patient's management based on radiologist's report and his clinical findings.

iii) Wherever available, histopathological diagnosis helps the surgeon in planning limb-salvaging surgery for early malignant and all benign bone lesions.

ACKNOWLEDGMENTS

We thank the Department of Radiology and Pathology in the Medical faculty for letting us access patient details.

We appreciate our patients' co-operation in the study.

REFERENCES

- Adil A, Hoefel, Fikry T (1996). Osteoid Osteoma After a fracture of the radius. *AJR*. 167: 145 -146.
- Ahuka OL, Lusi KM (1995). Amputations in rural areas of another Zaire: an aetiological and epidemiological study. *East and Central Afr. J. Surg.* 1:33-35.
- Bascoulergue G, Gorgeon F, Lecomte-Houcke M, Baviera E, Mazabraud M(1996). Malignant synovial chondromatosis or chondromatous transformation of synovial chondromatosis of the knee: *Bulletin du cancer*. 83(11): 951-956.
- Cohen J (1960). A coefficient of agreement for nominal scales. *Educational and Psychological Measurement*. 20: 37-46.
- Dodge OD (1973). Tumours of bone and jaw. In: Templeton AC, (Ed). *Turnouts in a Tropical country*. Springer - Verlag Berlin - Heidelberg. 14: 222-233.
- Dorfman HD, Czerniak B, Kotz R (2002). WHO classification of tumors bone: In *WHO Classification of tumors of soft tissue and bone*. Lyon: IARC press. p. 227- 232
- Duffaud F, Digue L, Baciuchka-Palmaro M, Volot F, perles Daniel C, Garbe L, Favre R (2000). Osteosarcomas of flat bones in adolescents and adults. *Cancer*. 88(2): 324-332
- Fornasier VL, Protzner K (1998). Radium induced tibial sarcoma in a treated case of hind foot angiomatosis: *Skeletal Radiol*. 27(3): 164-168.
- Gwinn JL, Senac MOJ, Isaacs H (1986). Primary lesions of bone in the first decade of life. *Radiol*. 160:49-495
- Hamada G, Rida A (1972). Orthopaedics and Orthopaedic diseases in Ancient and Modern Egypt: *Clin. Orthop*. 89: 253.
- Hasbini A, Lartigau E, Le Pechoux C, Acharki A, Vanel D, GeninJ, Le Cesne A (1998). Chondrosarcoma in Ollier's disease: a propos of two cases and review of Literature. *Cancer Radiother*. 4: 384-391.
- Jongkolnee (2006). Spectrum of bone tumors in Chiang Mai University. *J Med Assoc Thai*. 89(6): 780- 86.
- Kiwanuka M (1983). Lower limb amputation in Uganda. *Med. Thesis*. 3:40.
- Maeda G, Yokoyama R, Othomo K, TakayamaJ, Beppu Y, Fukuma H, Ohira M (1996). Osteochondroma after total body irradiation in bone marrow transplant patients: Report of 2 cases (Review) (14 Refs), *Jap. Clin. Oncol*. 6: 480-483.
- Online kappa calculation. <http://statpages.org/> and <http://www.mail-archive.com/edstat@lists.ncsu.edu/msg00789.html>
- Parkin W, Raymond Y (1997) Cancer incidence in five Continents IARC scientific publications. 7(143): 70-86.
- Rosen GF, CA, Mankin HJ, Selch MT(1997). Bone tumors. In Pine J W, editor, *Cancer Medicine*, Maryland. p. 2503-2550
- Schajowicz F(1993). Histological typing of bone tumors. World Health Organization International Histological Classification of Tumors. Berlin: Springer-Verlag.
- Ssebagala SJM (1998a). The clinical presentations of patients with Osteosarcoma in Mulago Hospital 1987-Jan 1997. *Med.Thesis*. 1:1-2

Ssebagala SJM(1998b). The clinical presentations of patients with Osteosarcoma in Mulago Hospital 1987- Jan 1997. Med.Thesis. 2: 4-10.

Turo S, Masaki C, Mitsuo N, Nideomi W, Eiichi U(1990). HLA Phenotypes in patients who have Osteosarcoma. J. Bone Joint Surg. 72A: 68.

Unni K (1996). Dahlin's bone tumors. general aspects and data on 11,087 cases. Lippincott-Raven, 5 edition,

Von Schulthess Ch, Zollikofer L(2005). In G.K. von Schulthess Ch.L. Zollikofer, editor, MUSCULOSKELETAL DISEASES. ISBN 88-470-08-Springer Milan Berlin Heidelberg, New York,