

Full Length Research Paper

Identification of Two Distinct Virulent Genotypes of *Xanthomonas oryzae* pv. *oryzae* in West Africa

Onasanya Amos^{1*}, M. M. Ekperigin², A. Afolabi⁴, R. O. Onasanya², Abiodun A. Ojo¹ and I. Ingelbrecht³

¹Department of Chemical Sciences, Afe Babalola University, Ado-Ekiti, Ekiti State, Nigeria.

²Federal University of Technology, Akure, PMB 704, Akure, Ondo State, Nigeria.

³Institute of Plant Biotechnology for Developing Countries, Ghent University, K. L. Ledeganckstraat 35, 9000 Ghent, Belgium.

⁴Biotechnology and Genetic Engineering Advanced Laboratory, Sheda Science and Technology Complex, PMB 186, Garki, Abuja, Nigeria

Accepted 6 April, 2024

Bacterial leaf blight (BLB) caused by *Xanthomonas oryzae* pv. *oryzae* (*Xoo*), is a very destructive rice disease worldwide. The aim of the present study was to examine if the *Xoo* virulence pathotypes obtained using phenotypic pathotyping could be confirmed using molecular approach. After screening of 60 Operon primers with genomic DNA of two *Xoo* isolates (virulent pathotype, *Vr* and mildly virulent pathotype, *MVr*), 12 Operon primers that gave reproducible and useful genetic information were selected and used to analyze 50 *Xoo* isolates from 7 West African countries. Genetic analysis revealed two major *Xoo* virulence molecular type (Mt) which were *Mta* and *Mtb* with *Mta* having two subgroups (*Mta1* and *Mta2*). *Mta1* (*Vr1*) subgroup genotype has occurrence in six countries and *Mta2* (*Vr2*) in three countries while *Mtb* genotype characterized mildly virulence (*MVr*) *Xoo* isolates present in five countries. The study revealed possible linkage and correlation between phenotypic pathotyping and molecular typing of *Xoo* virulence. Durable resistance rice cultivars would need to overcome both *Mta* and *Mtb* *Xoo* virulence genotypes in order to survive after their deployment into different rice ecologies in West Africa.

Key words: Bacterial leaf blight, *Xanthomonas oryzae* pv. *oryzae* (*Xoo*), *Xoo* virulence pathotype, molecular typing, genomic DNA, Operon primer, *Xoo* virulence genotype, *Xoo* pathogen migration, West Africa.

INTRODUCTION

Rice is one of the most widely cultivated food crop worldwide, but its production is constrained by fungal, bacterial and viral diseases. Bacterial leaf blight (BLB) caused by *Xanthomonas oryzae* pv. *oryzae* (*Xoo*), is a very destructive rice disease and its incidence has been reported from different parts of Asia, northern Australia, Africa and USA (Adhikari et al., 1995; Sere et al., 2005; Jiang et al., 2006). In West Africa, BLB disease incidence ranged from 70 to 85% and yield loss ranged from 50 to 90%, indicating a wide spread of BLB disease in farmers'

fields (Sere et al., 2005). Some selected *Xoo* isolates have shown high level of pathogenicity and virulence on the cultivated rice varieties (Sere et al., 2005; Onasanya et al., 2009; Dewa et al., 2011). Crop loss assessment studies have revealed that this disease reduces grain yield to varying levels, depending on the stage of the crop, degree of cultivar susceptibility and to a great extent, the conduciveness of the environment in which it occurs (Savary et al., 2006). The severity and significance of damage caused by infection have

*Corresponding author. Email: onasanyaa@abuad.edu.ng. Tel: +2348074289553.

necessitated the development of strategies to control and manage the disease, so as to reduce crop loss and to avert an epidemic. The identification and characterization of major genes for qualitative resistance and polygenic factors controlling quantitative resistance have contributed a great deal to the success in breeding resistant cultivars and their deployment (Chen et al., 2002). Recent research has provided considerable evidence that the deployment of bacterial antagonists to *Xoo* might be an effective strategy, bringing about disease suppression by biological control (Gnanamanickam, 2009). To understand the epidemiology and ecology of *Xoo* pathogens and their potential for virulence change, various phenotypic characters as well as molecular markers have been used in studies of *Xoo* pathogen population structure (Jiang et al., 2006). Identification and classification of bacteria are normally carried out by morphology, nutritional requirements, antibiotic resistance, isozyme comparisons, phage sensitivity (Akanji et al., 2011; Chaudhary et al., 2012) and more recently by DNA based methods, particularly rRNA sequences (Anzai et al., 2002; Chandrashekar et al., 2012), strain-specific fluorescent oligonucleotides (Zhao et al., 2007) and the polymerase chain reaction (PCR) (Akanji et al., 2011). Several repetitive elements found in the *Xoo* pathogen have been used as probes in restriction fragment length polymorphism (RFLP) analysis (Gonzalez et al., 2007). However, for the large number of samples needed for ecological and virulence studies, a simpler and cheaper technology is required. PCR is increasingly becoming an important tool in population biology, because of its simplicity and potential to rapidly screen a large number of samples with a minimal amount of DNA.

In West Africa, several *Xoo* genetic studies have been conducted and different *Xoo* pathotypes identified but little information is available on *Xoo* virulence genotypes population structure and distribution (Basso et al., 2011). The virulence pathotypes of several *Xoo* isolates from West African countries based on cultivars reactions has been determined (Basso et al., 2011; Dewa et al., 2011). The main goal of this study was to determine *Xoo* virulence genotypes using the characterized *Xoo* isolates virulence pathotypes identified by Onasanya et al. (2009) using random amplified polymorphic DNA polymerase chain reaction (RAPD-PCR) assays. The identification and differentiation of different *Xoo* virulence genotypes and distribution in West Africa would greatly help in rice breeding improvement programs aiming at the effective development of rice cultivars with durable resistance to BLB disease.

MATERIALS AND METHODS

Bacterial isolates

Fifty *X. oryzae* pv. *oryzae* (*Xoo*) isolates (Table 1) used in this study

were from Onasanya et al. (2009). The identity of all the fifty *Xoo* isolates had been confirmed by oxidative biochemical test as well as their virulence pathotypes (Onasanya et al., 2009).

Isolates propagation

BLB isolates were first propagated using a modified procedure developed by Akanji et al. (2011). Nutrient broth (75 ml; pH 7.5) was prepared inside a 100 ml conical flask. Each *Xoo* isolate (100 μ l) from storage was transferred into 50 ml of nutrient broth and kept under constant shaking at 30°C for 24 h for bacterial growth. The bacterial cell was removed by centrifugation, washed with 0.1

mM Tris-EDTA (pH 8.0) and kept at -20°C for DNA extraction.

Genomic DNA extraction

DNA extraction was according to Onasanya et al. (2003) with some modification. 0.3 g of washed bacterial cell was suspended in 200 μ l of cetyl trimethylammonium bromide (CTAB) buffer (50 mM Tris, pH 8.0; 0.7 mM NaCl; 10 mM EDTA; 2% hexadecyltrimethylammonium bromide; 0.1% 2-mercaptoethanol), followed by 100 μ l of 20% sodium dodecyl sulfate and incubated at 65°C for 20 min. DNA was purified by two extractions with chloroform and precipitated with -20°C absolute ethanol. After washed with 70% ethanol, the DNA was dried and resuspended in 200 μ l of sterilized distilled water. DNA concentration was measured using DU-65UV spectrophotometer (Beckman Instruments Inc., Fullerton CA, USA) at 260 nm. DNA quality was checked on a 1% agarose gel in Tris-Acetate-EDTA (TAE) buffer (45 mM Tris-acetate, 1 mM EDTA, pH 8.0) after electrophoreses.

RAPD-PCR analysis

This analysis was performed according to Akanji et al. (2011). DNA primers used were purchased from Operon Technologies (Alameda, CA, USA) and each was ten nucleotides long. Two concentrations of each DNA (25 and 95 ng per reaction) were used to test reproducibility and eliminate sporadic amplification products from the analysis. Sixty primers (OPP, OPQ, OPR, OPS, OPT, OPV, OPX and OPY series) were screened with DNA of two *Xoo* isolates (virulence, *Vr*, and mildly virulence, *MVr*, isolates) for their ability to amplify the *Xoo* genomic DNA. Primers that gave useful polymorphisms were selected and used in amplifying the DNA from all *Xoo* isolates. Amplifications was performed in 25 μ l reaction mixture consisting of genomic DNA, reaction buffer (Promega), 100 μ M each of dATP, dCTP, dGTP and dTTP, 0.2 μ M Operon random primer, 2.5 μ M MgCl₂ and 1U of Taq polymerase (Boehringer, Germany). A single primer was used in each reaction. Amplification was performed in a thermowell microtiter plate (Costa Corporation) using a MJ Research programmable thermal controller. The cycling program was (i) 1 cycle of 94°C for 3 min; (ii) 45 cycles of 94°C for 1 min for denaturation, 40°C for 1 min for annealing of primer and 72°C for 2 min for extension; and (iii) a final extension at 72°C for 7 min. Amplification products were maintained at 4°C until electrophoresis.

Electrophoresis of PCR products

The amplification products were resolved by electrophoresis in a

Table 1. Identity of *X. oryzae* pv. *oryzae* isolates used for the study.

S/N	Isolates code*	Host plant	Country
1	XN-1	D52-37	Niger
2	XN-2	D52-37	Niger
3	XN-3	IR15296829	Niger
4	XN-4	IR15296829	Niger
5	XN-5	WITA 8	Niger
6	XN-6	WITA 8	Niger
7	XB-7	Local	Benin
8	XB-8	Local	Benin
9	XB-9	Local	Benin
10	XB-10	Local	Benin
11	XB-11	Local	Benin
12	XNG-12	WITA9	Nigeria
13	XNG-13	WITA9	Nigeria
14	XNG-14	WITA 4	Nigeria
15	XNG-15	WITA 4	Nigeria
16	XNG-16	WITA 8	Nigeria
17	XBF-17	TS2	Burkina Faso
18	XBF-18	TS2	Burkina Faso
19	XBF-19	FKR14	Burkina Faso
20	XBF-20	FKR19	Burkina Faso
21	XBF-21	FKR14	Burkina Faso
22	XBF-22	Chinese	Burkina Faso
23	XM-23	Adventices	Mali
24	XM-24	Kogoni	Mali
25	XM-25	Kogoni	Mali
26	XM-26	Kogoni	Mali
27	XM-27	Kogoni	Mali
28	XM-28	Kogoni	Mali
29	XM-29	Jamajigi	Mali
30	XM-30	Nionoka	Mali
31	XG-31	Weed	Guinea
32	XG-32	Weed	Guinea
33	XG-33	Weed	Guinea
34	XG-34	Local	Guinea
35	XG-35	Local	Guinea
36	XG-36	Local	Guinea
37	XG-37	Local	Guinea
38	XG-38	Local	Guinea
39	XG-39	Local	Guinea
40	XG-40	Local	Guinea
41	XTG-41	Local	The Gambia
42	XTG-42	Local	The Gambia
43	XTG-43	Local	The Gambia
44	XTG-44	Local	The Gambia
45	XTG-45	Local	The Gambia
46	XTG-46	Local	The Gambia

Table 1. Cont.

47	XTG-47	Local	The Gambia
48	XTG-48	Local	The Gambia
49	XTG-49	Weed	The Gambia
50	XTG-50	Weed	The Gambia

* = *X. oryzae* pv. *oryzae* isolates obtained from Onasanya et al. (2009).

1.4% agarose gel using Tris-Acetate-EDTA (TAE) buffer (45 mM Tris-acetate, 1 mM EDTA, pH 8.0) at 100 V for 2 h. A 1 kb ladder (Life Technologies, Gaithersburg, MD, USA) was included as molecular size marker. Gels were visualized by staining with ethidium bromide solution (0.5 µg/ml) and banding patterns were photographed over UV light using UVP-computerized gel photo documentation system.

Cluster analysis

Positions of scorable amplified DNA bands were transformed into a binary character matrix ("1" for the presence and "0" for the absence of a band at a particular position). Pairwise distance matrices were compiled by the Numerical Taxonomy System (NTSYS) 2.0 software (Rohlf, 2000) using the Jaccard coefficient of similarity (Ivchenko and Honov, 1998). Cluster dendrogram was created by unweighted pair-group method arithmetic (UPGMA) cluster analysis (Éena et al., 2009). Principal component analysis with GGEbiplot was carried out on 50 *Xoo* isolates using genetic data generated from twelve Operon primers (Ebdon and Gauch, 2002).

RESULTS

Genetic analysis of fifty *X. oryzae* pv *oryzae* (*Xoo*) isolates from West Africa have been carried out. After screening of 60 Operon primers with genomic DNA of two *Xoo* isolates (virulent pathotype, *Vr* and mildly virulent pathotype, *MVr*), only 12 primers gave reproducible polymorphism and useful genetic information that differentiated the fifty *Xoo* isolates. Amplification with the 12 primers generated 210 bands from which 136 (64.8%) of them were polymorphic (Table 2) with sizes ranging between 0.5 and 4.0 kb (Figure 1). Using the 136 RAPD markers (Table 2) in cluster and principal component analyses revealed two major (*Mta* and *Mtb*) molecular typing (Mt) virulence genotypes among fifty *Xoo* isolates (Figures 2 and 3). *Mta* genotype was made up of 42 virulence (*Vr*) *Xoo* isolates with two subgroup genotypes (*Mta1* and *Mta2*). *Mta1* (*Vr1*) subgroup genotype was typical of 25 *Xoo* isolates with 50% occurrence in six countries (Niger, Benin Republic, Nigeria, Burkina Faso, Mali and Guinea) (Table 3). *Mta2* (*Vr2*) subgroup genotype was typical of 17 *Xoo* isolates with 34% occurrence in three countries (Mali, Guinea and The Gambia) (Table 3). *Mtb* genotype characterized 8 mildly virulence (*MVr*) *Xoo* isolates with 16% occurrence in five

Table 2. Oligonucleotide primers that showed genetic polymorphism among the *X. oryzae* pv. *oryzae* isolates using random amplified polymorphic DNA polymerase chain reaction analysis.

Operon primer	Nucleotide sequence 5' to 3'	No. of fragments amplified	No. of polymorphic bands	Polymorphism (%)
OPP-17	TGACCCGCCT	18	16	88.9
OPP-18	GGCTTGGCCT	14	11	78.6
OPR-07	ACTGGCCTGA	20	11	55.0
OPS-08	TTCAGGGTGG	23	13	56.5
OPS-10	ACCGTTCCAG	20	13	65.0
OPS-13	GTCGTTCCCTG	16	9	56.3
OPT-09	CACCCCTGAG	16	10	62.5
OPT-12	GGGTGTGTAG	13	7	53.8
OPT-15	GGATGCCACT	18	10	55.6
OPV-05	TCCGAGAGGG	19	12	63.2
OPY-06	AAGGCTCACC	16	11	68.8
OPY-08	AGGCAGAGCA	17	13	76.5
Total		210	136	64.8




Figure 1. DNA fingerprinting patterns of 50 *X. oryzae* pv. *oryzae* (*Xoo*) isolates using OPS-08 random amplified polymorphic DNA primer. M: 1kb molecular size marker; kb: kilobase pair. *Xoo* isolate: 1 = XN-1; 2 = XN-2; 3 = XN-3; 4 = XN-4; 5 = XN-5; 6 = XN-6; 7 = XB-7; 8 = XB-8; 9 = XB-9; 10 = XB-10; 11 = XB-11; 12 = XNG-12; 13 = XNG-13; 14 = XNG-14; 15 = XNG-15; 16 = XNG-16; 17 = XBF-17; 18 = XBF-18; 19 = XBF-19; 20 = XBF-20; 21 = XBF-21; 22 = XBF-22; 23 = XM-23; 24 = XM-24; 25 = XM-25; 26 = XM-26; 27 = XM-27; 28 = XM-28; 29 = XM-29; 30 = XM-30; 31 = XG-31; 32 = XG-32; 33 = XG-33; 34 = XG-34; 35 = XG-35; 36 = XG-36; 37 = XG-37; 38 = XG-38; 39 = XG-39; 40 = XG-40; 41 = XTG-41; 42 = XTG-42; 43 = XTG-43; 44 = XTG-44; 45 = XTG-45; 46 = XTG-46; 47 = XTG-47; 48 = XTG-48; 49 = XTG-49; 50 = XTG-50.




Figure 2. Molecular typing of 50 *Xanthomonas oryzae* pv. *oryzae* (*Xoo*) virulence as revealed by 136 random amplified polymorphic DNA markers. *Mta* = Molecular type a; *Mtb* = Molecular type b; *Vr* = Virulence; *MVr* = Mildly virulence. *Xoo* isolate: 1 = XN-1; 2 = XN-2; 3 = XN-3; 4 = XN-4; 5 = XN-5; 6 = XN-6; 7 = XB-7; 8 = XB-8; 9 = XB-9; 10 = XB-10; 11 = XB-11; 12 = XNG-12; 13 = XNG-13; 14 = XNG-14; 15 = XNG-15; 16 = XNG-16; 17 = XBF-17; 18 = XBF-18; 19 = XBF-19; 20 = XBF-20; 21 = XBF-21; 22 = XBF-22; 23 = XM-23; 24 = XM-24; 25 = XM-25; 26 = XM-26; 27 = XM-27; 28 = XM-28; 29 = XM-29; 30 = XM-30; 31 = XG-31; 32 = XG-32; 33 = XG-33; 34 = XG-34; 35 = XG-35; 36 = XG-36; 37 = XG-37; 38 = XG-38; 39 = XG-39; 40 = XG-40; 41 = XTG-41; 42 = XTG-42; 43 = XTG-43; 44 = XTG-44; 45 = XTG-45; 46 = XTG-46; 47 = XTG-47; 48 = XTG-48; 49 = XTG-49; 50 = XTG-50.

countries (Niger, Benin Republic, Nigeria, Burkina Faso, and Guinea) (Table 3). Thus in Niger, Benin Republic, Nigeria and Burkina Faso molecular typing revealed the

presence of *Mta1 (Vr1)* and *Mtb (MVr)* *Xoo* genotypes, *Mta1 (Vr1)* and *Mta2 (Vr2)* genotypes in Mali, *Mta1 (Vr1)*, *Mta2 (Vr2)* and *Mtb (MVr)* genotypes in Guinea, and




Figure 3. Principal component analysis that revealed subgroup virulence genotypes among 50 *X. oryzae* pv. *oryzae* isolates using genetic data generated from twelve random amplified polymorphic DNA (RAPD) primers.

Mta2 (Vr2) genotype in The Gambia (Figure 4 and Table 3).

DISCUSSION

Molecular basis for African *Xoo* virulence identification is a prerequisite into understanding the genetics of *Xoo* virulence population structure in West Africa and deployment of durable resistance cultivars (Sere et al., 2005; Adhikari et al., 1999; Adhikari et al., 1995). The present study examined if the two *Xoo* virulence pathotypes (*Pta* and *Ptb*) obtained using phenotypic pathotyping by Onasanya et al. (2009) could be confirmed using molecular approach. Molecular typing using random amplified polymorphic (RAPD) markers has revealed two major (*Mta* and *Mtb*) virulence genotypes

among the 50 *Xoo* isolates in which *Mta* was virulence (*Vr*) and *Mtb* mildly virulence (*MVr*). This report supports recent isozyme fingerprints of 30 *Xoo* isolates from 5 countries (Mali, Burkina Faso, Niger, Benin Republic and Nigeria) in West Africa that revealed two major genetic groups (Onasanya et al., 2008). These two genotypes of *Xoo* virulence identified by molecular typing were very identical to *Xoo* virulence pathotypes (*Pta* and *Ptb*) obtained using phenotypic pathotyping indicating possible linkage and correlation between phenotypic pathotyping and molecular typing of *Xoo* virulence (Adhikari et al., 1999; Lalitha et al., 2010).

The high distinction pattern of each isolates in this study suggests possible high level of genetic variation among *Xoo* isolates in different host cells (Innes et al., 2001; Mongkolsuk et al., 2000). The genetic analyses revealed that *Mta* virulence genotype might cover about 84% of BLB population across Niger, Benin Republic,

Table 3. *X. oryzae* pv. *oryzae* isolate group, virulence and distribution relative to country of origin.

Typing	Main group	Subgroup	Virulence	Isolate origin and distribution						Occurrence (%)	
				Niger	Benin	Nigeria	Burkina Faso	Mali	Guinea		The Gambia
Pathotype*	<i>Pta</i>	<i>Pta1</i>		-	-	-	4	1	4	1	20
		<i>Pta2</i>	<i>Vr</i>	3	-	-	-	-	1	4	16
		<i>Pta3</i>		-	2	3	1	2	3	-	22
	<i>Ptb</i>	<i>Ptb1</i>		2	2	1	1	1	1	2	20
		<i>Ptb2</i>	<i>MVr</i>	1	1	1	-	4	1	3	22
Molecular type	<i>Mta</i>	<i>Mta1</i>	<i>Vr1</i>	4	3	3	5	7	3	-	50
		<i>Mta2</i>	<i>Vr2</i>	-	-	-	-	1	6	10	34
	<i>Mtb</i>	-	<i>MVr</i>	2	2	2	1	-	1	-	16

* = Onasanya et al., 2009; *Pta* = pathotype a; *Ptb* = pathotype b; *Mta* = molecular type a; *Mtb* = molecular type b; *Vr* = virulence; *MVr* = mildly virulence.




Figure 4. *X. oryzae* pv. *oryzae* isolates virulence genotype population structure status in West Africa. *Mta* = Molecular type a; *Mtb* = molecular type b; *Vr* = virulence; *MVr* = mildly virulence. Country: C1 = Niger; C2 = Benin Republic; C3 = Nigeria; C4 = Burkina Faso; C5 = Mali; C6 = Guinea; C7 = The Gambia.

Nigeria, Burkina Faso, Mali, the Gambia and Guinea and possibly be responsible for most sporadic cultivars infestation and epidemics in these countries. Also, the existence of *Mta1* and *Mta2* subgroups were likely due to interactions among isolates and strains that originally constituted *Mta* genotype (Innes et al., 2001; Chisholm et al., 2006). *Mtb* genotype existed in over 16% of BLB population across Niger, Benin Republic, Nigeria, Burkina Faso and Guinea, and might be responsible for most sporadic cultivars infestation and epidemics in these countries. *Mta1* (*Vr1*) and *Mtb* (*MVr*) genotypes were found to exist in Niger, Benin Republic, Nigeria, and Burkina Faso, *Mta1* and *Mta2* in Mali, *Mta1*, *Mta2* and *Mtb* in Guinea, and *Mta2* in The Gambia suggesting possible *Xoo* pathogen migration between these countries and long-term *Xoo* pathogen survival (Adhikari et al., 1995; Dewa et al., 2011).

Distinct phenotypes usually consist of isolates that are genetically less related and such identification of isolates using cultural and morphological techniques often lack consistency and precision (Onasanya et al., 2009). Molecular typing of *Xoo* virulence has proven particularly useful in situations where it is necessary to differentiate virulence among two or more bacterial pathogens (Lalitha et al., 2010). In the current study, it was discovered that identification of virulence in *Xoo* depends on different host origins and occurrence of mutants. For instance, *Mta* virulence genotype might cover about 84% of BLB population across Niger, Benin Republic, Nigeria, Burkina Faso, Mali, The Gambia and Guinea and *Mtb* genotype existed in over 16% of BLB population across Niger, Benin Republic, Nigeria, Burkina Faso and Guinea but isolates virulence distributions vary within subgroups. Based on phylogenetic study, it was discovered that after prolonged season-to-season interactions among isolates of *Mta* or *Mtb* genotype in different cultivated rice and weed hosts, different subgroup virulence genotypes (*Mta1* and *Mta2*) may emerge as a result of mutation (Mongkolsuk et al., 2000). The emerged subgroup virulence genotypes might result in occurrence of highly virulent isolates and strains with very broad interaction and pathogenicity across wide range of cultivated rice varieties across West African countries.

Conclusion

The present molecular study of *Xoo* virulence identified two major *Xoo* virulence genotypes (*Mta* and *Mtb*) and two subgroups (*Mta1* and *Mta2*). Existence of different *Xoo* virulence genotypes suggests high level of *Xoo* pathogen interaction with host cells and mutation. The study revealed possible linkage between *Xoo* virulence pathotype and *Xoo* virulence genotype. Different *Xoo* virulence genotypes were known to exist within country and there was evidence of *Xoo* pathogen migration between countries. Durable resistance rice cultivars

would need to overcome both *Mta* and *Mtb* *Xoo* virulence genotypes in order to survive after their deployment into different rice ecologies in West Africa.

ACKNOWLEDGEMENT

We are very grateful to the Central Biotechnology Laboratory, International Institute of Tropical Agriculture (IITA) Nigeria for funding this research in its laboratory.

REFERENCES

- Adhikari TB, Vera Cruz CM, Zhang Q, Nelson RJ, Skinner DZ, Mew TW, Leach JE (1995). Genetic diversity of *Xanthomonas oryzae* pv. *oryzae* in Asia. *Appl. Environ. Microbiol.* 61: 966-971.
- Adhikari TB, Basnyat RC, Mew TW (1999). Virulence of *Xanthomonas oryzae* pv. *oryzae* on rice lines containing single resistance genes and gene combinations. *Plant Dis.* 83(1): 46-50.
- Akanji BO, Ajele JO, Onasanya A, Oyelakin O (2011). Genetic Fingerprinting of *Pseudomonas aeruginosa* Involved in Nosocomial Infection as Revealed by RAPD-PCR Markers. *Biotechnol.* 10(1): 70-77.
- Anzai Y, Kim H, Park JY, Wakabayashi H, Oyaizu H (2002). Phylogenetic affiliation of the Pseudomonads based on 16S rRNA sequence. *Int. J. Syst. Evol. Microbiol.* 50(4): 1563-1589.
- Basso A, Onasanya A, Issaka S, Sido AY, Haougui A, Adam T, Sere Y, Saadou M (2011). Bacterial leaf blight of rice in Niger: Pathological diversity of isolates collected on irrigated lands. *J. Applied Biosci.* 38: 2551-2563.
- Chandrashekar S, Umesha S, Chandan S (2012). Molecular detection of plant pathogenic bacteria using polymerase chain reaction single-strand conformation polymorphism. *Acta. Biochim. Biophys. Sin.* 44(3): 217-223.
- Chaudhary SU, Javed I, Muzzammil H (2012). Effectiveness of different fungicides and antibiotics against bacterial leaf blight in rice. *J. Agric. Res.* (2012). 50(1): 109-117.
- Chen H, Wang S, Zhang Q (2002). New gene for bacterial blight resistance in rice located on chromosome 12 identified from Minghui 63, an elite restorer line. *Phytopathology* 92: 750-754.
- Chisholm ST, Coaker G, Day B, Staskawicz BJ (2006). Host-microbe interactions: Shaping the evolution of the plant immune response. *Cell* 124: 803-814.
- Dewa KMK, Banito A, Onasanya A, Kpemoua KE, Sere Y (2011). Rice Bacterial Blight in Togo: Importance of the Disease and Virulence of the Pathogen. *Current Research in Bacteriology* 4: 94-100.
- Ebdon JS, Gauch HG (2002). AMMI analysis of national turfgrass performance trials: I. Interpretation of genotype by environment interaction. *Crop Sci.* 42:489-496.
- Éena J, Eszter A, Péter I, Arnold H, János P (2009). BOOL-AN: A method for comparative sequence analysis and phylogenetic reconstruction. *Mol. Phylogenet. Evol.* 52(3): 887-897.
- Gnanamanickam SS (2009). An Overview of Progress in Biological Control. *Biomedical and Life Sciences* 8: 43-51.
- Gonzalez C, Szurek B, Manceau C, Mathieu T, Sere Y, Verdier V (2007). Molecular and pathotypic characterization of new *Xanthomonas oryzae* strains from West Africa. *Mol. Plant Microbe Interact.* 20(5): 534-546.
- Innes D, Beacham IR, Beven CA, Douglas M, Laird MW, Joly JC, Burns DM (2001). The cryptic *ushA* gene (*ushAc*) in natural isolates of *Salmonella enterica* (serotype Typhimurium) has been inactivated by a single missense mutation. *Microbiology* 147: 1887-1896.
- Ivchenko GI, Honov SA (1998). On The Jaccard Similarity Test. *J. Mathematical Sci.* 88(6): 789-794.
- Jiang GH, Xia ZH, Zhou YL, Wan J, Li DY, Chen RS, Zhai WX, Zhu LH (2006). Testifying the rice bacterial blight resistance gene xa5 by genetic complementation and further analyzing xa5 (*Xa5*) in comparison with its homolog TFLIAY1. *Mol. Gen. Genomics* 275: 354-

366.

- Lalitha SM., Varma CMK, Premalatha P, Devi GL, Zehr U, Freeman W (2010). Understanding the Bacterial Blight Pathogen-Combining Pathotyping and Molecular Marker Studies. *International J. Plant Pathol.* 1 (2): 58 – 68.
- Mongkolsuk S, Whangsuk W, Fuangthong M, Loprasert S (2000). Mutations in *oxyR* resulting in peroxide resistance in *Xanthomonas campestris*. *J. Bacteriol.* 182: 3846- 3849.
- Onasanya A, Mignouna HD, Thottappilly G (2003). Genetic fingerprinting and phylogenetic diversity of isolates of *Staphylococcus aureus* from Nigeria. *Afri. J. Biotechnol.* 2(8): 246 – 250.
- Onasanya A, Ekperigin MM, Nwilene FE, Sere Y, Onasanya RO (2009). Two pathotypes of *Xanthomonas oryzae* pv. *oryzae* virulence identified in West Africa. *Current Res. Bacteriol.* 2(2): 22-35.
- Onasanya A, Ekperigin MM, Sere Y, Nwilene FE, Ajele JO (2008). Enzyme polymorphism and genetic diversity in *Xanthomonas oryzae* pv. *oryzae* isolates causing rice bacterial leaf blight disease in West Africa. *Int. J. Agric. Res.* 3(3): 227-236.
- Rohlf FJ (2000). *NTSys pc*, Version 2.02j, Exeter Software, Setauket, New York.
- Savary S, Teng PS, Willcoquet L, Jr. Nutter FW (2006). Quantification and modeling of 25 crop losses: a review of purposes. *Ann. Rev. Phytopathol.* 44:89-112.
- Sere Y, Onasanya A, Verdier V, Akator K, Ouédraogo LS, Segda Z, Mbare MM, Sido AY, Basso A (2005). Rice Bacterial Leaf Blight in West Africa: Preliminary Studies on Disease in Farmers' Fields and Screening Released Varieties for Resistance to the Bacteria. *Asian J. Plant Sci.* 4: 577-579.
- Zhao WJ, Zhu SF, Liao XL, Chen HY, Tan TW (2007). Detection of *Xanthomonas oryzae* pv. *oryzae* in seeds using a specific TaqMan probe. *Molecular Biotechnology* 35(2): 119-127.