

Full Length Research Paper

## ***In vitro* rat lymphocyte proliferation induced by *Ocimum basilicum*, *Persea americana*, *Plantago virginica*, and *Rosa* spp. extracts**

R. Gomez-Flores<sup>1\*</sup>, L.Verástegui-Rodríguez<sup>1</sup>, R. Quintanilla-Licea<sup>2</sup>, P. Tamez-Guerra<sup>1</sup>, R. Tamez-Guerra<sup>1</sup> and C. Rodríguez-Padilla<sup>1</sup>

<sup>1</sup>Departamento de Microbiología e Inmunología, Universidad Autónoma de Nuevo León, San Nicolás de los Garza, NL, México.

<sup>2</sup>Departamento de Química, Facultad de Ciencias Biológicas Universidad Autónoma de Nuevo León, San Nicolás de los Garza, NL. México.

Accepted 16 January, 2019

Medicinal plants have been used for centuries and have become part of complementary medicine worldwide because of their potential health benefits. Some of their metabolites have been successfully used directly in the treatment and prevention of infectious diseases and cancer, or indirectly by stimulating the immune system. In the present study, we investigated the effects of methanol and aqueous extracts of the Mexican plants *Ocimum basilicum*, *Persea americana*, *Plantago virginica*, and *Rosa* spp. on *in vitro* rat lymphocyte proliferation. Methanol extracts of *O. basilicum*, *P. americana*, *P. virginica*, and *Rosa* spp. stimulated up to 80, 16, 69 and 66% lymphoproliferation, respectively, whereas their respective aqueous extracts induced up to 83, 48, 31 and 83% lymphoproliferation, as compared with untreated controls. The effect of *O. basilicum* aqueous extract at concentrations of 31.25, 62.5, 125 and 250 µg/ml on lymphoproliferation was significantly different ( $p < 0.05$ ) than the effects of *P. americana* and *P. virginica* at the same concentrations. We also observed that the lymphoproliferative effect of *Rosa* spp. aqueous extract was significantly higher ( $p < 0.05$ ) than that of the methanol extract. Methanol and aqueous vehicles did not affect lymphocyte viability nor proliferation activity. The observed immunostimulatory effect may be of benefit in increasing the pool of lymphocytes in immunodeficiency patients.

**Key words:** *Ocimum basilicum*, *Persea americana*, *Plantago virginica*, *Rosa* spp., immunoregulation, lymphocyte proliferation, phytomitogens, mexican plants.

### INTRODUCTION

In the last three decades there has been a significant increase in the use of natural products in health care, and their potential applications in agriculture, pharmaceutical and food industry are being investigated. In fact about 30% of drugs used in industrialized countries are derived from plants (Kaufman et al., 1999). Plants play an essential role in the health care needs of native populations in America not only for the treatment of diseases, but also

to improve the immunological response against many pathologies (Borchers et al., 2000). The use of preparations and infusions of plants to treat diseases has been practiced widely in Mexico and Central America for several centuries (Gomez-Flores et al., 2000; Kinghorn et al., 1998), but their effectiveness should be scientifically validated to increase the credibility of their use.

In the present study, we evaluated the effects of methanol and aqueous extracts of the widely distributed plants *Ocimum basilicum*, *Persea americana*, *Plantago virginica*, and *Rosa* spp. on *in vitro* rat lymphoproliferation, an activity that has not been previously studied or reported for these plants. *O. basilicum* L. (Lamiaceae)

\*Corresponding author. E-mail: [rgomez60@hotmail.com](mailto:rgomez60@hotmail.com).

(basil, sweet basil) is mostly used in cosmetics, liqueurs, medicines, and perfumes. Its oil has antimicrobial and anti-inflammatory properties (Singh, 1999). Similarly, *P. americana* Mill (Lauraceae) (avocado) seeds are rich in tannins, and are used to treat diarrhoea; carotenoids and tocopherols from the fruit were shown to inhibit the *in vitro* growth of prostate cancer cell lines (Lu et al., 2005). Additionally, *P. americana* leaves have been reported to possess anti-inflammatory and analgesic activities (Adeyemi et al., 2002). *Plantago* species, on the other hand, have been reported to possess anti-inflammatory and antitumor properties (Samuelson, 2000), and induce rat lymphocyte proliferation and nitric oxide and tumor necrosis factor- $\alpha$  production by rat peritoneal macrophages (*Plantago major*) (Gomez-Flores et al., 2000). Finally, people have used *Rosa spp.* (Rosaceae) (Rose wild) leaves for their laxative and astringent properties.

## MATERIAL AND METHODS

### Reagents and culture media

Penicillin-streptomycin solution, L-glutamine, and RPMI 1640 and AIM-V media were obtained from Life Technologies (Grand Island, NY). Fetal bovine serum (FBS), sodium dodecyl sulfate (SDS), N,N-dimethylformamide (DMF), PBS, and 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide (MTT) were purchased from Sigma-Aldrich (St. Louis, MO). Extraction buffer was prepared by dissolving 20% (wt/vol) SDS at 37°C in a solution of 50% each DMF and demineralized water, and the pH was adjusted to 4.7.

### Animals

Sprague-Dawley male rats (200 – 220 g) were purchased from Harlan Sprague-Dawley, Inc. (Indianapolis, IN). They were kept in a pathogen- and stress-free environment at 24°C, under a light-dark cycle (light phase, 06:00 - 18:00 h), and given water and food *ad libitum*. Animals were killed by asphyxiation in a 100% CO<sub>2</sub> chamber.

### Collection and processing of plants

Plants were collected during summer in San Nicolás de los Garza N. L., México. *O. basilicum*, *P. virginica*, *P. americana* and *Rosa spp.* used in this study were identified by M.Sci. María del Consuelo González de la Rosa, Chief of the Herbarium of the Biological Sciences College at Autonomous University of Nuevo Leon, with voucher specimen numbers 024780 for *O. basilicum*, 024783 for *P. virginica*, 024782 for *P. americana*, and 024781 for *Rosa*. Leaves were rinsed, dried at 37°C for 36 h, and macerated in a coffee grinder (Proctor Silex E160B). Methanol extracts were prepared by placing 4.3 g of the leaf powder in 80 ml absolute methanol under shaking for 24 h at room temperature. The resulting extracts were centrifuged at 2800 rpm for 15 min, and supernatants were placed in 1 ml Eppendorf tubes, previously weighted. Then this material was dried under vacuum using a speed-vac concentrator (Savant Instruments Inc., Hicksville, NY), and dissolved in 300  $\mu$ l methanol and 700  $\mu$ l RPMI 1640 medium supplemented with 10% FBS, 1% L-glutamine, and 0.5% penicillin-streptomycin solution (referred as complete RPMI medium). The aqueous extracts were prepared by placing 1g of the plant powder in 50 ml boiling distilled water for 10

min. The resulting extracts were then centrifuged at 2800 rpm for 15 min, and supernatants were placed in 1 ml glass tubes, previously weighted. Then this material was dried by lyophilizing, and dissolved in 300  $\mu$ l methanol and 700  $\mu$ l complete RPMI 1640 medium. Stock solutions of both methanol and aqueous extracts were then prepared at 0.5 mg/ml in complete RPMI 1640 medium, and sterilized by filtering through a 0.22  $\mu$ -membrane (Millipore, Bedford, MA).

### Phytochemical analysis

The phytochemical analyses were done using standard procedures (Trease and Evans, 1996; Mojab et al., 2003; Olaleye, 2007).

### Cell preparation and culture

Thymus was removed immediately after rat death (see above), and a single cell-suspension was prepared by disrupting the organ in RPMI 1640 medium as previously reported (Nowak et al., 1998). The cell suspension was washed three times in this medium, and suspended and adjusted to 1 X 10<sup>7</sup> cells/ml in complete RPMI 1640 medium.

### T cell proliferation assay

T cell proliferation was determined by a colorimetric technique using MTT (Franco-Molina et al., 2003). Thymus cell suspensions (100  $\mu$ l of 1 X 10<sup>7</sup> cells/ml) were added to flat-bottomed 96-well plates (Becton Dickinson) containing triplicate cultures (100  $\mu$ l) of RPMI 1640 medium supplemented with 5% fetal bovine serum (unstimulated control), plant extracts at various concentrations, or vehicles (similarly processed as with plant methanol and aqueous extractions, but without plant material) for 48 h at 37°C in 95% air-5% CO<sub>2</sub> atmosphere. After incubation, MTT (0.5 mg/ml, final concentration) was added, and cultures were additionally incubated for 4 h. Cell cultures were then incubated for 16 h with extraction buffer (100  $\mu$ l), and optical densities, resulting from dissolved formazan crystals, were then read in a microplate reader (Bio-Tek Instruments, Inc., Winooski, VT) at 540 nm (absorbance or optical density at 540 nm wavelength). The lymphocyte proliferation index (LPI) was calculated as follows:

$$LPI = \frac{A_{540} \text{ in plant extract treated}}{\text{cells } A_{540} \text{ in untreated cells}}$$

Percentages of proliferation, provided in the Results section, were calculated by using the lymphocyte proliferation indexes as compared with untreated controls (that is, an LPI of 1.6 indicates 60% increase in proliferation).

### Statistical analysis

The results were expressed as mean  $\pm$  SEM of the response of 3 replicate determinations from three independent experiments. Level of significance was assessed by Student's *t* test and ANOVA.

## RESULTS


### Phytochemical analysis

Phytochemical screening procedures were carried out to

**Table 1.** Phytochemical analysis of the Mexican plant methanolic and aqueous extracts.

Plant	Extract	Alk*	Flav	Sap	Terp	Tann	Sac
<i>OB (basil)</i>	Me	+	-	-	+	+	+
	Aq	-	-	-	-	-	+
<i>PA (avocado)</i>	Me	+	+	-	+	+	+
	Aq	+	+	-	+	+	+
<i>PV (hoary plantain)</i>	Me	+	-	+	+	+	+
	Aq	-	-	+	+	-	+
<i>Rose spp.</i>	Me	+/-	+	+/-	+	+	+
	Aq	-	+	+	-	+	+

\*Alk =, alkaloids; Flav = flavonoids; Sap = saponins; Terp = terpenes; Tann = tannins; Sac = saccharides.



**Figure 1.** Lymphoproliferation induced by *Ocimum basilicum* methanolic and aqueous extracts. Rat thymic cell suspensions were incubated in the presence or absence of various concentrations of *O. basilicum* methanolic and aqueous extracts, after which lymphoproliferation was measured colorimetrically, as explained in the text. Data represent means  $\pm$  SEM of triplicate determinations from three independent experiments. \*\* $p < 0.01$ , \* $p < 0.05$  compared with *O. basilicum* extract-untreated control. Optical densities at 540 nm for untreated cells were  $0.95 \pm .01$  and  $1.1 \pm .04$  for methanolic and aqueous extract treatments, respectively. A PI of 1 indicates no alteration on proliferation.

determine the potential active compounds that may be involved in the observed biological activities. Table 1 revealed that alkaloids were present in all methanolic extracts. The aqueous extracts did not contain alkaloids. Furthermore, flavonoids were only detected in *P. americana* and *Rosa* methanolic and aqueous extracts; saponins were detected in *P. virginica* and *Rosa* methanolic and aqueous extracts; terpenes were present in all but aqueous *O. basilicum* and *Rosa* extracts; tannins were present in all but aqueous *O. basilicum* and *P. virginica* extracts; and saccharides were present in both methanolic and aqueous extracts (Table 1).


#### Effect of plant extracts on T cell proliferation

*O. basilicum* methanolic extract significantly ( $p < 0.05$ ) sti-


mulated 36 and 80% proliferation (from LPI data) of resident lymphocytes at 125 and 250  $\mu\text{g/ml}$  respectively, whereas *O. basilicum* aqueous extract stimulated 15, 35, 44 and 83% proliferation of resident lymphocytes at 31.25, 62.5, 125 and 250  $\mu\text{g/ml}$  respectively, as compared with untreated control (Figure 1); there was no difference between the effect of the aqueous and methanolic extracts. The effect of *O. basilicum* aqueous extract at concentrations of 31.25, 62.5, 125 and 250  $\mu\text{g/ml}$  on lymphoproliferation was significantly different ( $p < 0.05$ ) than the effects of *P. americana* and *P. virginica* at the same concentrations. Optical densities at 540 nm for untreated control cells were  $0.95 \pm .01$  and  $1.1 \pm .04$  for methanolic and aqueous extract treatments, respectively; these were used to calculate the LPIs.

In addition, *P. americana* methanolic extract significantly ( $p < 0.05$ ) stimulated 17% proliferation (from LPI data) of resident lymphocytes at 250  $\mu\text{g/ml}$ , whereas the aqueous extract stimulated 12, 20 and 48% lymphoproliferation at 62.5, 125 and 250  $\mu\text{g/ml}$  respectively, as compared with untreated control; there was no difference between the effect of the aqueous and methanolic extracts (Figure 2). The effect of *P. americana* aqueous extract at 250  $\mu\text{g/ml}$  on lymphoproliferation was significantly different ( $p < 0.05$ ) than the effects of *P. virginica* and *Rosa* at the same concentration. Optical densities at 540 nm for untreated cells were  $0.73 \pm .10$  and  $1.06 \pm .04$  for methanolic and aqueous extract treatments, respectively; these were used to calculate the LPIs. Similarly, *P. virginica* methanolic extract significantly ( $p < 0.05$ ) stimulated 29, 30 and 69% proliferation (from LPI data) of lymphocytes at 62.5, 125 and 250  $\mu\text{g/ml}$  respectively, whereas the aqueous extract stimulated 11 and 31% lymphoproliferation at 125 and 250  $\mu\text{g/ml}$  respectively, as compared with untreated control; there was no difference between the effect of the aqueous and methanolic extracts (Figure 3). Optical densities at 540 nm for untreated cells were  $0.84 \pm .17$  and  $0.60 \pm .08$  for methanolic and aqueous extract treatments, respectively; these were used to calculate the LPIs.


Furthermore, *Rosa* methanolic extract significantly ( $p < 0.01$ ) stimulated 66% lymphoproliferation (from LPI data)



**Figure 2.** Lymphoproliferation induced by *Persea americana* methanol and aqueous extracts. Rat thymic cell suspensions were incubated in the presence or absence of various concentrations of *P. americana* methanol and aqueous extracts, after which lymphoproliferation was measured colorimetrically, as explained in the text. Data represent means  $\pm$  SEM of triplicate determinations from three independent experiments. \*\* $p < 0.01$ , \* $p < 0.05$  compared with *P. americana* extract-untreated control. Optical densities at 540 nm for untreated cells were  $0.73 \pm .10$  and  $1.06 \pm .04$  for methanol and aqueous extract treatments, respectively. A PI of 1 indicates no alteration on proliferation.



**Figure 4.** Lymphoproliferation induced by *Rosa* methanol and aqueous extracts. Rat thymic cell suspensions were incubated in the presence or absence of various concentrations of *Rosa* methanol and aqueous extracts, after which lymphoproliferation was measured colorimetrically, as explained in the text. Data represent means  $\pm$  SEM of triplicate determinations from three independent experiments. \*\* $p < 0.01$ , \* $p < 0.05$  compared with *Rosa* extract-untreated control. Optical densities at 540 nm for untreated cells were  $0.92 \pm .06$  and  $0.75 \pm .02$  for methanol and aqueous extract treatments, respectively. A PI of 1 indicates no alteration on proliferation.



**Figure 3.** Lymphoproliferation induced by *Plantago virginica* methanol and aqueous extracts. Rat thymic cell suspensions were incubated in the presence or absence of various concentrations of *P. virginica* methanol and aqueous extracts, after which lymphoproliferation was measured colorimetrically, as explained in the text. Data represent means  $\pm$  SEM of triplicate determinations from three independent experiments. \*\* $p < 0.01$ , \* $p < 0.05$  compared with *P. virginica* extract-untreated control. Optical densities at 540 nm for untreated cells were  $0.84 \pm .17$  and  $0.60 \pm .08$  for methanol and aqueous extract treatments, respectively. A PI of 1 indicates no alteration on proliferation.

at 250  $\mu\text{g/ml}$ , whereas the aqueous extract stimulated 36 and 83% lymphocyte proliferative response at 125 and 250  $\mu\text{g/ml}$  respectively, as compared with untreated control; the aqueous extract significantly ( $p < 0.001$ ) induced higher lymphoproliferative activity than the methanol extract (Figure 4). Optical densities at

540 nm for untreated cells were  $0.92 \pm .06$  and  $0.75 \pm .02$  for methanol and aqueous extract treatments, respectively; these were used to calculate the LPs.


Vehicles for the methanol and aqueous extracts did not alter lymphocyte proliferation (Figure 5).

## DISCUSSION

Natural products play an important role in the field of new drugs research and development (Jin-Ming et al., 2003). Knowledge of plant biologically active compounds and their mechanism (s) of action are desirable, not only for the discovery of novel therapeutic agents that would validate folkloric remedies, but also for the design of new active molecules or modification of current drugs against diverse maladies. One important application area is the immunotherapy; plant products have been shown to modulate the immune system (Spelman et al., 2006).

In the present study, we focused our study on some Mexican plants and demonstrated their potential to stimulate lymphocyte proliferation. Lymphocytes respond to antigen challenge by proliferating, expanding the antigen-specific lymphocyte clones and producing lymphokines, thus amplifying immune responses (Mackay, 1993). This effect can be mimicked by the use of polyclonal phyto-mitogens such as concanavalin A and phytohemagglutinin, which bind to certain sugar residues on T cell surface glycoproteins, including the T cell receptor and CD3 protein, and stimulate T cell proliferative response (Gajewski et al., 1989).

We found that all the tested extracts stimulated lym-



**Figure 5.** Effect of methanol and aqueous extract vehicles on lymphocyte proliferation. Rat thymic cell suspensions were incubated in the presence or absence of various concentrations of vehicles for the methanol and aqueous extracts (prepared as explained in the text), after which lymphoproliferation was measured colorimetrically. Data represent means  $\pm$  SEM of triplicate determinations from three independent experiments. Optical densities at 540 nm for untreated cells were  $0.59 \pm .05$  and  $0.67 \pm .09$  for methanol and aqueous extract vehicle treatments, respectively.

lymphocyte proliferation. In the case of *O. basilicum*, such an activity was not due to the presence of alkaloids, flavonoids, terpenes, saponins, and tannins, because only saccharides were detected in these extracts (Table 1). However, it has been reported that *O. basilicum* leaves contain flavonoids and terpenoids (Grayer et al., 1996; Lemberkovics et al., 1998); in fact, a terpenoid called eugenol was shown to possess anti-inflammatory and antibiotic properties (Miele et al., 2001). In our study, *O. basilicum* extracts were found to induce the highest lymphoproliferative response (up to 80 and 83% for the methanol and aqueous extracts, respectively), as compared with untreated controls. Although, *O. basilicum*'s main reported activity is anti-inflammatory (Singh et al., 1999), its essential oil did not show significant effect on proliferation of murine lymphocytes stimulated with LPS and Con A as reported by others (Courreges and Benencia, 2002), but appeared to modulate both humoral and cell-mediated immune responses (Mediratta et al., 2002). In *P. americana* extract we detected the presence of flavonoids, terpenes, tannins, and saccharides (Table 1), as reported by others (Antia et al., 2005). Although, the aqueous extract of this plant was shown to possess anti-inflammatory and analgesic activities (Adeyemi et al., 2002), lectins from its seeds were not mitogenic for mouse lymphocytes (Meade et al., 1980), and "persin" from avocado leaves was shown to have antifungal properties and to be toxic to silkworms (Oelrichs et al., 1995), immune enhancing properties of this plant has not been previously reported. In addition, we observed the presence of alkaloids, saponins, terpenes, tannins, and saccharides in *P. virginica* extracts; the presence of such

compounds has been reported in *Plantago* species (Samuelson, 2000). The species *P. major* has been broadly used to treat diverse medical conditions ranging from pain to infectious diseases and cancer (Matev et al., 1982; Lithander, 1992; Grigorescu et al., 1973); particularly, anti-inflammatory properties were reported for this species (Herold et al., 2003), whereas Gomez-Flores et al. (2000) showed increased lymphoproliferative response of Con A-treated rat lymphocytes by *P. major* leaf extracts. Furthermore, Biringanine et al. (2005) reported *P. palmate*-mediated stimulation of nitric oxide and tumor necrosis factor-alpha production by activated macrophages.

Finally, the *Rosa* spp. aqueous extract was observed to significantly ( $p < 0.05$ ) induce higher lymphoproliferative activity than the methanol extract; this extract mainly contained flavonoids, saponins, and saccharides (Table 1). Although *Rosa* flowers were demonstrated to inhibit human immunodeficiency virus type-1 reverse transcriptase in-vitro (Fu et al., 2006), and Boyanova and Neshev (1999) showed inhibitory effect of rose oil products on *Helicobacter pylori* growth in vitro, immunomodulation by *Rosa* leaf extracts has not been previously reported.

Plant extracts are potentially curative. Some of these extracts can boost the humoral (Rehman et al., 1999) and cell-mediated immunity (Upadhyay et al., 1992) against viruses (Calixto et al., 1998), bacteria (Boyanova and Neshev, 1999), fungi (Ali et al., 1999), protozoa (Sharma et al., 1998), and cancer (Wong et al., 1994). In the present study, we demonstrated that various Mexican plant extracts stimulated murine lymphoproliferation. However, further studies are necessary to isolate the responsible active compound (s) and elucidate its mechanism of action. The regulation of immune parameters induced by plant extracts may be clinically relevant in numerous disease processes including chronic viral infections, tuberculosis, AIDS, and cancer.

## ACKNOWLEDGEMENTS

This study was supported by grants CN-285-00 from CONACYT and PAICYT 04 to RGF.

## REFERENCES

- Adeyemi OO, Okpo SO, Ogunti OO (2002). Analgesic and anti-inflammatory effect of the aqueous extract of leaves of *Persea americana* Mill (Lauraceae). *Fitoterapia* 73: 375-380.
- Ali MI, Shalaby NM, Elgamil MH, Mousa AS. (1999). Antifungal effects of different plant extracts and their major components of selected aloe species. *Phytother. Res.* 13: 401-407.
- Antia BS, Okokon JE, Okon PA. (2005). Hypoglycemic activity of aqueous leaf extract of *Persea americana* Mill. *Indian J. Pharmacol.* 37: 325-326.
- Biringanine G, Vray B, Vercruyse V, Vanhaelen-Fastre R, Vanhaelen M, Duez P (2005). Polysaccharides extracted from the leaves of *Plantago palmata* Hook.f. induce nitric oxide and tumor necrosis factor-alpha production by interferon-gamma-activated macrophages. *Nitric Oxide* 12: 1-8.
- Borchers AT, Sakai S, Henderson GL, Harkey MR, Keen CL, Stern JS,

- Terasawa K, Gershwin ME (2000). Shosaiko-to and other Kampo (Japanese herbal) medicines: a review of their immunomodulatory activities. *J. Ethnopharmacol.* 73: 1-13.
- Boyanova L, Neshev G. (1999). Inhibitory effect of rose oil products on *Helicobacter pylori* growth *in vitro*: preliminary report. *J. Med. Microbiol.* 48: 705-706.
- Calixto JB, Santos AR, Cechinel Filho V, Yunes RA. (1998). A review of the plants of the genus *Phyllanthus*: their chemistry, pharmacology, and therapeutic potential *Med. Res. Rev.* 18: 225-258.
- Courreges MC, Benencia F (2002). *In vitro* antiphagocytic effect of basil oil on mouse macrophages. *Fitoterapia* 73: 369-374.
- Franco-Molina M, Gomez-Flores R, Tamez-Guerra P, Tamez-Guerra R, Castillo-Leon L, Rodriguez-Padilla C. (2003). *In vitro* immunopotentiating properties and tumour cell toxicity induced by *Lophophora williamsii* (peyote) cactus methanol extract. *Phytother. Res.* 17:1076-1081.
- Fu M, Ng TB, Jiang Y, Pi ZF, Liu ZK, Li L, Liu F. (2006). Compounds from rose (*Rosa rugosa*) flowers with human immunodeficiency virus type 1 reverse transcriptase inhibitory activity. *J. Pharm. Pharmacol.* 58: 1275-1280.
- Gajewski TF, Schell SR, Nau G, Fitch FW (1989). Regulation of T-cell activation: differences among T-cell subsets. *Immunol. Rev.* 111: 79-110.
- Gomez-Flores R, Calderon CL, Scheibel LW, Tamez-Guerra P, Rodriguez Padilla C, Tamez-Guerra R, Weber RJ. (2000). Immunoenhancing properties of *Plantago major* leaf extract. *Phytoter. Res.* 14: 617-622.
- Grayer RJ, Kite GC, Goldstone FJ, Bryan SE, Paton A, Putievsky E. (1996). Intraspecific taxonomy and essential oil chemotypes in sweet basil, *Ocimum basilicum*. *Phytochemistry* 43: 1033-1039.
- Grigorescu E, Stanescu U, Basceanu V, Aur MM (1973). Phytochemical and microbiological control of some plant species used in folk medicine. II. *Plantago lanceolata* L., *Plantago media* L., *Plantago major* L. *Rev. Med. Chir. Soc. Med. Nat. Iasi.* 77: 835-841.
- Herold A, Cremer L, C Iugaru A, Tama V, Ionescu F, Manea S, Szegli G. (2003). Antioxidant properties of some hydroalcoholic plant extracts with antiinflammatory activity. *Roum. Arch. Microbiol. Immunol.* 62: 217-227.
- Jin-Ming K, Ngoh-Khang G, Lian-Sai Ch, Tet-Fatt Ch. (2003). Recent advances in traditional plant drugs and orchids *Acta Pharmacol. Sin.* 24: 7-21.
- Kaufman PB, Cseke LJ, Waber S, Duke JA, Briemann HL. (1999). Natural products from plants. Boca Raton, Florida: CRC Press.
- Kinghorn AD, Ito A, Kennelly EJ, Kim NC, Westenburg HE (1998). Studies on some edible and medicinal plants of Mesoamerica. *Proc. West Pharmacol. Soc.* 41: 253-258.
- Lemberkovics E, Kery A, Marczal G, Simandi B, Szoke E (1998). Phytochemical evaluation of essential oils, medicinal plants and their preparations. *Acta Pharm. Hung.* 68: 141-149.
- Lithander A. (1992). Intracellular fluid of waybread (*Plantago major*) as a prophylactic for mammary cancer in mice. *Tumour Biol.* 13: 138-141.
- Lu QY, Arteaga JR, Zhang Q, Huerta S, Go VL, Heber D (2005). Inhibition of prostate cancer cell growth by an avocado extract: role of lipid-soluble bioactive substances. *J. Nutr. Biochem.* 16: 23-30.
- Mackay CR (1993). Immunological memory. *Adv. Immunol.* 53: 217-265.
- Matev M, Angelova I, Koichev A, Leseva M, Stefanov G. (1982). Clinical trial of *Plantago major* preparation in the treatment of chronic bronchitis. *Vutr. Boles* 21: 133-137.
- Meade NA, Staat RH, Langley SD, Doyle RJ. (1980). Lectin-like activity from *Persea americana*. *Carbohydr. Res.* 78: 349-363
- Mediratta PK, Sharma KK, Singh S (2002). Evaluation of immunomodulatory potential of *Ocimum sanctum* seed oil and its possible mechanism of action. *J. Ethnopharmacol.* 80: 15-20.
- Miele M, Dondero R, Ciarallo G, Mazzei M. (2001). Methyleugenol in *Ocimum basilicum* L. Cv. genovese gigante. *J. Agric. Food Chem.* 49: 517-521.
- Mojab F, Kamalinejad M, Ghaderi N, Vahidipour HR (2003). Phytochemical screening of some species of iranian plants. *Iranian J. Pharm. Res.* 2: 77-82.
- Nowak JE, Gomez-Flores R, Calderon SN, Rice KC, Weber RJ. (1989). Rat natural killer cell, T cell and macrophage functions after intracerebroventricular injection of SNC 80. *J. Pharmacol. Exp. Ther.* 286: 931-937.
- Oelrichs PB, Ng JC, Seawright AA, Ward A, Schaffeler L, MacLeod JK (1995). Isolation and identification of a compound from avocado (*Persea americana*) leaves which causes necrosis of the acinar epithelium of the lactating mammary gland and the myocardium. *Nat. Toxins* 3: 344-349.
- Olaleye MT (2007). Cytotoxicity and antibacterial activity of methanol extract of *Hibiscus sabdariffa*. *J. Med. Plants Res* 1: 9-13.
- Rehman J, Dillow, JM, Carter SM, Chou J, Le BB, Maisel AS (1999). Increased production of antigen-specific immunoglobulins G and M following *in vivo* treatment with the medicinal plants *Echinacea angustifolia* and *Hydrastis canadensis*. *Immunol. Lett.* 68: 391-396.
- Samuelson AB (2000). The traditional uses, chemical constituents and biological activities of *Plantago major* L. A review. *J. Ethnopharmacol.* 71: 1-21.
- Sharma P, Sharma JD (1998). Plants showing antiplasmodial activity--from crude extracts to isolated compounds. *Indian J. Malariol.* 35: 57-110.
- Singh S (1999). Mechanism of action of antiinflammatory effect of fixed oil of *Ocimum basilicum* Linn. *Indian J. Exp. Biol.* 37: 248-252.
- Spelman K, Burns JJ, Nichols D, Winters N, Ottersberg S, Tenborg M (2006). Modulation of cytokine expression by traditional medicines: a review of herbal immunomodulators. *Altern. Med. Rev.* 11: 128-150.
- Trease GE, Evans WC (1966). A Textbook of Pharmacognosy. 9th edn. London: Baillière, Tindall and Cassell.
- Upadhyay S, Dhawan S, Garg S, Talwar GP (1992). Immunomodulatory effects of neem (*Azadirachta indica*) oil. *Int. J. Immunopharmacol.* 14: 1187-1193.
- Wong CK, Leung KN, Fung KP, Choy YM. (1994). Immunomodulatory and anti-tumour polysaccharides from medicinal plants. *J. Int. Med. Res.* 22: 299-312.