

Full Length Research Paper

Fruit performances of the selected fig types in Turkey

Mikdat Simsek

Department of Horticulture, Faculty of Agriculture, Bingöl University, Bingöl, 12000, Turkey. E-mail: miksimsek2001@yahoo.com. Tel: +90-426-213 25 50 – 51. Fax: +90-426-213 28 66.

Accepted 19 September, 2019

In this study, fruit performances of some types of fig (*Ficus carica* L.) were determined in Dara and Yardere villages of Mardin, Turkey. Seven fig types were evaluated in the study for two years. A lot of pomological characteristics to determine the fruit performances of the these types were examined during the 2004 and 2005. According to the averages in two years, fruit weight ranged between 56.29 and 30.88 g, fruit length ranged from 41.80 to 31.97 mm, ostium width ranged between 3.90 and 2.44 mm, total soluble solids content (TSS) ranged between 23.40 and 18.25% and acidity ranged between 0.29 and 0.14%. 47-00-4, 47-00-6 and 47-00-7 types scored the highest in overall quality according to the results of the weighted ranked method.

Key words: *Ficus carica*, Southeast Anatolia region climate, fruit performance, table fig, selection.

INTRODUCTION

The world fig production was 1,056,820 tons according to Food and Agriculture Organization of the United Nations' (FAO) statistics. The world's largest fig producing country is Turkey. The fig production of Turkey was 285,000 tons, which was 27% of the world's total production. The fig export of Turkey was 177,900 tons, which represents 52% of total world fig exports (Anonymous, 2005). About 70% of Turkey's total fig production was for dry consumption (Aksoy et al., 2003). Because of environmental effects on the characteristics of the fruit performances, it was generally believed that the highest quality dried figs were grown in limited areas of the Big and Small Meander valleys where temperature, relative humidity and wind conditions were determined to be optimum for production of high quality dried figs (Ozbek, 1978). Fresh fig production, although still environmentally sensitive, appears to be less demanding in terms of climate characteristics. The figs are well adapted to some regions of the Southeast Anatolia region.

Bursa are the largest fresh fig region in Turkey with extensive fresh fig exports (Aksoy et al., 1992; Ozeker and Isfandiyarolu, 1998). However, Turkey's fresh fig production has not yet fulfilled its export potential. Recently, the fresh fig trade, confined primarily to local markets, has gained international importance (Sahin, 1998). Turkey's fresh fig exports only increased from 150 tons in 1980 to 6123 tons in 1999 and to 10,376 tons in 2004 (Anonymous, 2004). To increase fig production, it will be necessary to extend the harvesting period by planting in diverse ecological regions to utilize cultivars

that differ in their harvest period, and finally to improve quality and increase diversity. In Dara and Yardere villages, none of the fig trees is in solid orchards but is inter-planted with almond trees. The objective of this study was to determine the performance and important characteristics of the selected types to increase the fig production and the export of Turkey. In this context, the Southeast Anatolia region and the Mediterranean shore of Turkey have suitable conditions for the fig production (Kaska et al., 1990). Because of increasing demand for fresh consumption types and cultivars, any method to increase the fig production would be of value (Aksoy et al., 1992).

MATERIALS AND METHODS

This study was carried out 2004 – 2005 in Dara and Yardere villages of Mardin province, which is located in the Southeast Anatolia region of Turkey. 7 fig types were included in the study. The fruit performances of the selected fig types were determined. In this context, 30 fruits were randomly selected from the each fig tree in each year. Harvested fruits were immediately transferred to ice boxes and then stored at 0°C. After that, they were analysed with 3 replication and 10 fruits in each replication for the each year. To provide positive contributions to the production and export of the fig, the quality evaluation of the types was performed according to a weighted ranked method (Table 1). The data were subjected to analysis of variance using JMP 5.0.1. The means were separated by Tukey's test at 0.05.

Fruit weight was measured with a scale sensitive to 0.01 g. The fruit length and width, neck length, ostiole width were measured with a digital compass. Total soluble solids were determined with a hand-held refractometer and pH of fruit juice was determined with

Table 1. Quality evaluation of the selected fig types according to the weighted ranked method.

Characteristics	Weighting factor (coefficient)	Classification and points			
Fruit weight	40	<20.0 g	0	20.1 -30.0 g	2
		30.1 -40.0 g	4	40.1 -50.0 g	6
		50.1 -60.0 g	8	> 60.0 g	10
Fruit shape index	9	l<0.9	8	l=0.9-1.1	10
		l>1.1	6		
Neck length	6	<5.0 mm	0	5.1-10.0 mm	10
		10.1-15.0 mm	6	>15.0 mm	2
Fruit skin cracks	10	None-little	10	Medium	6
		High	0		
Peeling of skin	10	Easy	10	Medium	6
		Difficult	0		
Ostium width	5	0.0-2.0 mm	10	2.1-4.0 mm	8
		4.1-6.0 mm	6	>6.1 mm	2
Total soluble solid content	10	< 13.0%	2	13.1-16.0%	4
		16.1-20.0%	10	20.1-25.1%	8
		> 25.1%	6		
Titrable acidity	10	< 0.050%	0	0.051-0.125%	6
		0.126-0.225%	8	0.226-0.300%	10
		> 0.301%	4		
Total	100				

a pH-meter. Titrable acidity was determined by titrating with 0.1 N NaOH to an endpoint of pH 8.10. The total soluble solid/titrable acidity ratio was calculated. The fruit shape index was calculated by dividing the width by length. In addition to the these characteristics, ease of peeling and fruit skin cracks were evaluated according to the fig descriptor of Aksoy (1991). The coordinates and altitudes of the types were determined with CPS tool.

Beginning of maturation, full maturity, harvesting period, yield, fruit shape, fruit stalk shape, abscission of stalk from the twig, plant growth habit, tree vigor, number of lobes, leaves per shoot, tendency to form suckers, shoot colour, terminal bud colour, number of lobes per leaf, number of leaf per shoot, leaf shape, leaf margin dentation, leaf venation, uniformity of fruit size, abnormal fruit formation, apical dormancy, density of hairs on leaf upper and lower surface, petiole colour, fruit cavity, fruit skin color, shape of the fruit stalk, pulp internal colour, colour of scales around the ostium, adhesion of scales around the ostium, scale size around the ostium, drop at the eye and colour of liquid drop at the ostiole were determined based on the fig descriptor developed by Anonymous (2003).

RESULTS

Pomological variables

To determine the fruit performances in this study, some pomological characteristics of seven fig types were presented in Table 2. All pomological characteristics were significant at 5% level considering 2 years mean results (2004 and 2005).

According to the averages in the 2 years, 47-00-6 had

the highest fruit weight (56.29 g) and 47-00-3 had the lowest fruit weight (30.88 g). Similarly, 47-00-6 had the highest fruit width (55.89 mm) and 47-00-3 had the lowest fruit width (41.05 mm). 47-00-7 had the highest fruit length (41.80 mm) and 47-00-3 had the lowest fruit length (31.97 mm). No neck was observed in 1 type (47-00-8), while the others had necks and the lengths of the neck were changed between 7.68 mm in 47-00- 7 type and 3.93 mm in 47-00-3 type. The ostium width of the selected fig types ranged between 3.90 mm in 47-00-1 and 2.44 mm in 47-00-8.

The total soluble solids were highest for 47-00-6 (23.43%) and lowest for 47-00-1 (18.25%) . The pH of the fruit juice was highest for 47-00-4 (6.04) and lowest for 47-00-9 (4.67). The titrable acidity was highest for 47-00-3 (0.29%) and lowest for 47-00-7 (0.14%). The TSS/titrable acidity was highest for 47-00-7 (137.03) and lowest for 47-00-03 (63.11).

Weighted characterization

When all the characteristics were evaluated together using the weighted analysis '47- 00-4, 47-00-6 and 47-00-7' was calculated as having the highest fruit quality (754 points). According to the weighed ranked method, it was determined that all the fig types had the fruit shape index 54 score, the skin cracks 100 score and the ostium width 40 score (average of years 2004-2005).

Table 2. Some pomological characteristics of the selected fig types in the Southeast Anatolia region (Average of years 2004-2005).

Accession no.	Fruit weight (g)	Fruit length (mm)	Fruit width (mm)	Neck length (mm)	Ostiolum width (mm)	Titrable acidity		pH	TSS/ Titrable acidity
						TSS (%)	(%)		
47-00-1	49.37 bc	38.42 a	51.51 b	5.40 c	3.90 a	18.25 b	0.27 ab	5.29 ab	68.89 c
47-00-3	30.88 e	31.97 b	41.05 e	3.93 d	3.29 b	18.40 b	0.29 a	5.79 a	63.11 c
47-00-4	52.17 ab	38.99 a	50.60 bc	5.36 c	3.78 a	20.20 b	0.24 b	6.04 a	84.53 bc
47-00-6	56.29 a	41.43 a	55.89 a	5.88 b	3.61 ab	23.43 a	0.23 b	5.70 a	100.70 b
47-00-7	50.58 b	41.80 a	52.06 ab	7.68 a	3.76 a	18.38 b	0.14 c	5.26 ab	137.03 a
47-00-8	44.24 c	34.22 b	47.52 cd	0.00 e	2.44 c	19.73 b	0.23 b	5.27 ab	87.19 bc
47-00-9	36.77 d	34.10 b	44.73 de	5.29 c	2.48 c	19.92 b	0.24 b	4.67 b	84.67 bc

Mean separation within columns by Tukey's test at 0.05 level.




Figure 1. The scores of the neck length and the ease of peeling of all the fig types (average of years 2004-2005).

It was showed that the scores of the neck length and the ease of peeling off all the fig types were in Figure 1; the scores of the TSS and titrable asidity of them were Figure 2 and the scores of the fruit weight and the total points of them were Figure 3.

Fruit and plant characteristics

In this study, it was determined that all the fig types had the tendency to form suckers high, the shoot colour brown, the fruit cavity none, the pulp internal colour white, the colour of scales around the ostiolum different from skin, the uniformity of fruit size variable, the abnormal fruit formation scarce, the leaf venation apparent, the abscission of the stalk from the twig hard and the fruit skin cracks none.

Table 3 shows the other fruit and plant characteristics. In this context, the beginning of maturation of the fig types studied ranged between July 20 (early) and August 15 (mid-season). Full maturity of all types was in August (early or mid-season). The harvesting period was continued for at least 15-20 days (short) and at longest 41-60

days (long) . In addition, because the selected fig types were in the gardens of the producers, the yield values of the types were not successfully obtained. Therefore, the yield values were determined as subjective with 1-5 assessment. In the result, it was observed to be medium efficiency (3) of 2 types, and good efficiency (4) of 3 types and very efficiency (5) of 2 types in the selected figs.

The fruit shape of all the fig types was oblate because the index was changed from 1.39 to 1.25. The fruit stalk shape was short and thick (J) for 3 types, long and slender (I) for 3 types and variously enlarged (A) for 1 type. The ease of peeling of the selected fig types were easy in 47-00-3 type and hard in others.

The plant growth habits were obtained as open for 47-00-1, weeping for 47-00- 8, semi-erect for 47-00-7 and spreading for other types. The tree vigors were found low for 47-00-4, intermediate for 47-00-1 and 47-00-9 and high for other types.

The terminal bud colours were light green for 47- 00-7 and green for other types. The number of lobes per leaf was evaluated with five lobes for 47-00- 3, 47-00-4 and47-00-6. Other types were determined with seven lobes. The




Figure 2. The scores of the titrable acidity and the TSS of all the fig types (average of years 2004-2005).




Figure 3. The scores of the fruit weight and the total points of all the fig types (average of years 2004-2005).

lobes. The number of leaf per shoot was ranged from 11.4 to 4.8. Highest number of leaf per shoot for 47-00-6 was 11.4. The lowest number of leaf per shoot were observed as 4.8 for 47-00-1. The leaf shapes was found as base calcarate, lobes latate (D) for 47-00-3 and base calcarate, lobes latate (C) for 47-00-4 and 47-00-6. Other types were found as base calcarate, lobes linear (A). The leaf margin dentations were observed as lobes sides completely dented for 47-00-1 and 47-00-6. Other

types were determined with only upper margins dented.

The apical dormancy was present for 47-00-7. Other types were determined with absent. The density of hairs on leaf upper surface was intermediate for 47-00-7 and sparse for other types. The density of hairs on leaf lower surface was intermediate for 47-00-8, none for 47-00-4 and 47-00-6 and sparse for other types. The petiole colour was green for 47-00-1, 47-00-4 and 47-00-7 and light green for other types. The fruit skin colour was yellow

Table 4. Names of fig types, origins, coordinates and altitudes of the selected fig types in the Southeast Anatolia Region (Average of years 2004-2005).

Accession no.	Names of fig types	Origins	Coordinates	Altitudes (m)
47-00-1	Pete i	Mardin-Dara	37673672 E-4118488 N	599
47-00-3	Zeytuni	Mardin-Dara	37673616 E-4118439 N	598
47-00-4	Pete i	Mardin-Dara	37673647 E-4118458 N	601
47-00-6	Pete i	Mardin-Dara	37673634 E-4118449 N	590
47-00-7	Hejir	Mardin-Yardere	37674259 E-4120096 N	622
47-00-8	Hejir	Mardin-Yardere	37642252 E-4120106 N	630
47-00-9	Zekaki	Mardin-Yardere	37674252 E-4120117 N	642

for 3 types and light green for other types. The shape of the fruit stalk was long and slender (I) in 47-00-1 and 47-00-8, short and thick (J) for 47-00-4, 47-00-7 and 47-00-9 and variously enlarged (A) for 47-00-3.

The adhesion of scales around the ostiolum was detached for 47-00-4 and adhered for other types. The scale size around the ostiolum was medium for 47-00-1, 47-00-6 and 47-00-7 and small for other types. The drop at the eye was present for 47-00-3, 47-00-4 and 47-00-8 and absent for other types. The colour of liquid drop at the ostiole was transparent for 47-00-3, 47-00-4 and 47-00-8 absent for other types.

Names, Origins, Coordinates and Altitudes

Names, origins, coordinates and altitudes of all the fig types were showed in Table 4. All the these types was selected in Dara and Yardere villages of Mardin. The local names of the types were Pete i, Zeytuni, Hejir and Zekaki. The coordinates of 47-00-1 were 37673672 E-4118488 N and the coordinates of 47-00-9 were 37674252 E-4120117 N. 47-00-6 had lowest altitude (590 m) and 47-00-9 had highest altitude (642 m). In addition, Map of Turkey and its areas of some fig types or the cultivars were showed in Figure 4.

DISCUSSION

Fig is grown in nearly all subtropical climates. In Turkey, a number of wild and cultivated forms of fig can be found with a great diversity of shape, flavor and color, primarily for fresh consumption (Ozbek, 1978). The Southeast Anatolia region and mediterranean shore have suitable conditions for fresh fig production (Ka ka et al., 1990).

Because of increasing demand for fresh consumption types and cultivars, any method to increase fig production would be of value (Aksoy et al., 1992).

In this study, the fruit weight is very important for fresh consumption in figs (Aksoy et al., 1992). The fruit weight values ranged from 56.29 to 30.88 g. In similar studies of fig types or cultivars, the fruit weight ranged from 30 to 90 g (Ozeker and Isfandiyaroglu, 1998), 24 to 92 g (Sahin et al., 2001), and 28 to 107 g (Polat and Ozkaya, 2005).

The fruit width values ranged from 55.89 to 41.05 mm. In addition, the fruit length values ranged from 41.80 to 31.97 mm. For precise quantification, the length, width, and fruit size were measured as well as the fruit weight (Eisen, 1901; Condit, 1941; Aksoy et al., 1992). In this study, results concerning the fruit width were higher than those of Polat and Ozkaya (2005) but the fruit length were lower than those of them. The fruit shape index (width/length) is of great importance in packing and transportation (Condit, 1941).

The neck length was longest on 47-00-7 (7.68 mm) and 47-00-6 (5.88 mm). In other studies, the longest neck was 8.86 mm (Calı kan and Polat, 2008), 14.5 mm (Ilgın, 1995), 8.70 mm (Ozeker and Isfandiyaroglu, 1998) and 8.01 mm (Polat and Ozkaya, 2005). In this study, however, the neck lengths of the selected types were lower than those of other researches.

The ostiolum width values ranked from 3.90 to 2.44 mm. The ostiole width was reported as 0.6–9.1 mm (Aksoy et al., 1992), 1.5–4.0 mm (Koyuncu et al., 1998), and 1.0–9.4 mm (Polat and Ozkaya, 2005). In this study, the ostiole width was partly different from the results of the other researchers. A large ostiole on the fig is an undesirable characteristic as pests and pathogens enter the fruit (Can, 1993).

In this study, the total soluble solid contents were ranged from 23.40 to 18.25. These results were lower than those of Calı kan and Polat (2008). But, the results in the study were better than those of them because high quality table figs in term of the TSS contents are better if they are between 13.0 and 25.1% (Aksoy et al., 1992). The titrable acidity ranged from 0.29 to 0.14%. These results were different from those (0.480-0.177%) of Küden et al. (2008). The pH of the fruit juice ranged from 6.04 to 4.67. The TSS/titrable acidity ratio is one of the most important factors in fruit taste (Karaçalı, 2002). Preferred ratio will vary with the use of fig fruits, but ratios will provide guidance in selecting types and cultivars for specific uses (Can, 1993). TSS/titrable acidity ranged from 137.03 to 63.11. These results were good for table figs.

In this study, the fruit stalk shapes of the selected fig types were determined as in Ilgın (1995) and Simsek (2008). In this context, the fruit stalk shape can change

- Eisen G (1901). The Fig. Its history, culture and curing. USDA Bull. 9: 201–283.
- Ilgin M (1995). The investigation of fertilization biology of some fig genotypes selected from Kahramanmaraş, region. PhD Thesis, Cukurova University, Adana, Turkey (in Turkish, with English abstract).
- Karaçalı (2002). Storage and marketing of horticultural products. Ege Uni., Agric. Fac., Publication no: 494 (in Turkish).
- Ka ka N, Küden AB, Küden A, Cetiner S (1990). The investigation of adaptation of Aegean figs and figs selected from Cukurova region. J. Cukurova Uni. Agric. Fac. (in Turkish, with English abstract). 5: 77–86
- Koyuncu MA, Bostan SZ, Islam A, Koyuncu F (1998). Investigation on physical and chemical characteristics in fig cultivars grown in Ordu. Acta Hortic. 480: 87–89.
- Küden AB, Beyazit S, Cömlekçio lu S (2008). Morphological and pomological characteristics of fig genotypes selected from Mediterranean and South East Anatolia Regions. Proceedings of the Third Int. Symposium on Fig. Acta Hortic. ISHS 798: 95-102.
- Ozbek S (1978). Özel Meyvecilik. Cukurova Üniversitesi, Ziraat Fakültesi Yayınları, Adana, (in Turkish). 128: 486p.
- Ozeker E, Isfandiyaroglu M (1998). Evaluation of table fig cultivars in Cesme Peninsula. Acta Hortic. 480: 55–60.
- Polat AA, Ozkaya M (2005). Selection studies on fig in the Mediterranean Region of Turkey. Pak. J. Bot. 37 (3): 567–574.
- Sahin N (1998). Fig adaptation studies in western Turkey. Acta Hortic. 480: 61–70.
- Sahin N, Cabanoglu F, Sahin B (2001). Fig report. T.R. Prime ministry state planning organization. Planning of development with five years. Plant production (fruits). Report of special committee. Ankara (in Turkish) 548p.
- im ek M (2008). Diyarbakır ko ulla rında incir genetik materyalinin seleksiyonu ve tanımlanması. Çukurova Üniversitesi, Science Institute, PhD. thesis, Adana, Türkiye.